



# Irreducible Metacarpal Neck Fracture due to Interposition of the Extensor Tendons

## *Fratura irreductível do colo do metacarpo devido a interposição dos tendões extensores*

Sathya Vamsi Krishna<sup>1</sup> Rajendra Kumbar<sup>1</sup> Sagar Vishwanath Kura<sup>1</sup> Soumen Das De<sup>2</sup>

<sup>1</sup> Department of Hand Surgery, Sanjay Gandhi Institute of Orthopaedics and Trauma, Byrasandra, Jayanagar, Bangalore, India

<sup>2</sup> Department of Hand & Reconstructive Microsurgery, National University Health System, Singapore

Address for correspondence Sathya Vamsi Krishna, MBBS, MS, Department of Hand Surgery, Sanjay Gandhi Institute of Orthopaedics and Trauma, Byrasandra, Jayanagar, Bangalore, India (e-mail: sathyavamsi@yahoo.com).

Rev Bras Ortop 2024;59(Suppl S1):e9–e11.

### Abstract

#### Keywords

- fractures, bone
- fracture fixation, internal
- metacarpal bones
- tendons

A 15-year-old male presented with multiple right-hand fractures, including a severely angulated small finger metacarpal neck fracture. Multiple closed attempts failed, and open reduction was performed. At surgery, the extensor tendon was found to be interposed within the fracture, thereby preventing closed reduction. The tendon was extricated from the fracture site, adequate reduction was obtained, and the fracture was stabilized using K-wires. The fracture united well, with good return of motion and strength. Extensor tendon interposition is a rare scenario associated with metacarpal neck fractures and should be suspected when there is complete loss of contact between the fracture ends and multiple attempts at closed reduction have failed.

### Resumo

#### Palavras-chave

- fixação interna de fraturas
- fraturas ósseas
- ossos metacárpico
- tendões

Um jovem do sexo masculino com 15 anos de idade apresentou múltiplas fraturas da mão direita, incluindo fratura do colo do quinto metacarpo com grande angulação. As múltiplas tentativas de redução fechada falharam, sendo então realizada a redução aberta. Foi encontrado durante a cirurgia o tendão extensor interposto dentro da fratura, impedindo desta forma a redução fechada. O tendão foi liberado do sítio da fratura, de modo que foi obtida a redução adequada e a fratura foi estabilizada com fios de Kirschner. A fratura foi bem consolidada, com um bom retorno do movimento e da força. A interposição do tendão extensor é um cenário raro associado a fraturas do colo metacárpico, devendo se suspeitar dela quando houver perda completa de contato entre as extremidades da fratura e as várias tentativas de redução fechada falharem.

*Study carried out at the Department of Hand Surgery, Sanjay Gandhi Institute of Orthopaedics and Trauma, Byrasandra, Jayanagar, Bangalore, India.*

received  
November 23, 2020  
accepted  
April 7, 2021

DOI <https://doi.org/10.1055/s-0041-1731656>.  
ISSN 0102-3616.

© 2021. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (<https://creativecommons.org/licenses/by/4.0/>).

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

## Introduction

Metacarpal fractures comprise 30% of all hand fractures, and the small and ring metacarpals are the most commonly affected.<sup>1</sup> Fractures of the small finger metacarpal neck – also known as “boxer’s fracture” – are frequently caused by a direct impact on the fully flexed metacarpophalangeal joint (MPJ).<sup>2</sup> Although boxer’s fractures have good healing potential, the relative indications for surgery include rotational deformity, angulation  $> 60^\circ$ , and pseudoclawing of the digit. Although it is unusual, the inability to obtain closed reduction or the occurrence of nonunion in these fractures should always raise suspicion of interposition of surrounding structures within the fracture site, mandating intervention. In the present report, we present a rare case of irreducible 5<sup>th</sup> metacarpal neck fracture due to an interposed extensor tendon that required open reduction.

## Case Report

A 15-year-old male presented with multiple hand fractures after a motor vehicle accident. On examination, there was loss of the metacarpophalangeal joint (MPJ) (“knuckle”) prominences of the small and ring fingers with swelling and tenderness over these areas. There was a puncture wound over the dorsum of the small finger proximal interphalangeal joint (PIPJ). Radiographs showed fractures of the ring and small finger metacarpal necks and an intra-articular fracture of the head of the proximal phalanx of the small finger (►Fig. 1). The small finger metacarpal was angulated  $> 60^\circ$ , with minimal contact between the bone ends. Surgical treatment of the injuries was performed. First, the small finger wound was debrided, and the head of proximal phalanx was fixed with a single 0.045-inch Kirschner wire (K-wire). The small finger metacarpal neck fracture was manipulated multiple times using the Jahss maneuver, but closed reduction was not possible, and the fracture site was exposed via a dorsal midline incision. We found that the proximal fragment was displaced dorsally and that the extensor digiti quinti minimi (EDQM) and the extensor

digitorum communis of the small finger (EDC-IV) were interposed between the fracture fragment, concealing the distal fragment (►Fig. 2). The juncturae tendinum was divided on the radial side, and the ulnar sagittal band was incised so that the tendons could be extricated. The head of the metacarpal was reduced and fixed with 2 crossed 0.045-inch K-wires (►Fig. 1). The ring finger metacarpal neck fracture was reduced using a closed technique and was stabilized using percutaneous K-wires. The hand was immobilized in a thermoplastic splint for 4 weeks in an intrinsic-plus position. After this period, the K-wires were removed and intermittent active motion was initiated. The fractures united in a timely manner and the patient returned to all pre-injury activities at 3 months (►Fig. 1).

## Discussion

In children, metacarpal neck fractures of the small and ring fingers are common due to the metacarpal geometry. The metacarpal angulates distally toward the MCPJ, and the cortical bone is known to be thin within the subcondylar fossa, making it susceptible to injury. These juxtaaphyseal fractures are analogous to adult boxer’s fracture and have good healing potential.<sup>3</sup> Most 5<sup>th</sup> metacarpal neck fractures with angulation  $< 40^\circ$  can be managed conservatively with ulnar gutter splint by immobilizing the wrist, the MCPJ, and the PIPJ in functional position for 3 weeks followed by gentle range of motion. However, fractures with increased angulation, pseudoclawing and rotational deformities are reduced by the Jahss maneuver, and the fractures can be stabilized by various techniques. Malunion is the most commonly known complication leading to loss of knuckle prominence, and nonunion is rare. Requirement of open reduction is exceptional for these fractures unless there is a difficulty in reduction or trouble in correcting the angulation. In our case, we had to perform open reduction since closed maneuver failed to achieve reduction due to interposition of the EDC-IV and the EDQM between the fracture. The probable mechanism could be due to a circumferential injury of the periosteum at the fracture site and buttonholing of the



**Fig. 1** (A,B) Anteroposterior and oblique radiographs of the affected hand showing juxtaaphyseal fracture on the neck of the fourth and fifth metacarpals; (C) Postoperative radiographs of the affected hand following fracture fixations with multiple Kirschner wires; (D/E) Follow-up radiographs of the hand at 5 months revealing a well-united fracture of the 5<sup>th</sup> metacarpal neck.



**Fig. 2** Intraoperative picture showing fracture of the 5<sup>th</sup> metacarpal with the interposed extensor tendons (marked by white arrow marks).

proximal fragment between the tendons and the sagittal band. If this were neglected, there could have been a consequence of definitive fracture nonunion. Tendon interposition is a rare presentation with very few case reports. Blohm et al<sup>4</sup> described a similar presentation with tendon and juncturae tendinum interposition, and Joshy et al.<sup>5</sup> reported a case of boxer's fracture nonunion due to extensor digitorum tendon hindrance.

Although it is rare, the interposition should be anticipated in a Boxer's fracture whenever there is an obvious gap on a

radiograph with complete loss of contact at the fracture ends and difficulty in getting adequate closed reduction. Surgeons should be prepared for a subsequent open reduction. With these measures, nonunion can be avoided, and early functions of hand can be restored.

**Statement of Informed Consent:** The parents of the patient were informed that data concerning the case would be submitted for publication, with which they completely agreed and consented. The present study is exempted from Ethical Committee approval.

#### Financial Support

No external funding was received for any aspect of the present work.

#### Conflict of Interests

The authors declare that they have no conflict of interests.

#### Acknowledgments

We would like to acknowledge the support of Dr. Avinash in the execution of this article.

#### References

- 1 van Onselen EB, Karim RB, Hage JJ, Ritt MJ. Prevalence and distribution of hand fractures. *J Hand Surg Br* 2003;28(05): 491–495
- 2 Soong M, Got C, Katarincic J. Ring and little finger metacarpal fractures: mechanisms, locations, and radiographic parameters. *J Hand Surg Am* 2010;35(08):1256–1259
- 3 Lightdale-Miric N, Kozin SH. Fractures and Dislocations of the Hand and Carpal Bones in Children. In: Rockwood and Wilkins' Fractures in Children. 8th ed. Wolters Kluwer; 2015:263
- 4 Blohm D, Bojsen B. [Interposition of the extensor tendon in a subcapital metacarpal fracture]. *Ugeskr Laeger* 1996;158(47): 6781
- 5 Joshy S, Ebrahim S, Iossifidis A. Tendon interposition in boxer's fracture, an unusual cause of non-union and extensor dysfunction of the little finger. *Inj Extra* 2005;36(07):237–238