

Nonaccidental Trauma Managed with Open Spinal Fixation and Instrumentation and a Review of the Literature

Michael J. Gigliotti¹ Noa Farou² Sandip Savaliya¹

Elias Rizk¹

¹Department of Neurosurgery, Penn State Health Milton S. Hershey Medical Center, Hershey, Pennsylvania, United States

²School of Medicine, Penn State Health Milton S. Hershey Medical

Center, Hershey, Pennsylvania, United States

Indian J Neurotrauma 2022;19:117–121.

Abstract

Keywords

- ► nonaccidental trauma
- fusion
- ► spinal injury
- ► trauma

Nonaccidental trauma (NAT), causing spinal injury is rare and occurs in up to 3% of cases. Management of these injuries is typically conservative, and thus surgical management is not widely reported in the literature. In this case report, we presented three patients to review the effectiveness of spinal instrumentation and posterior fusion in pediatric patients due to NAT. All patients recovered well and were neurologically intact at last follow-up with no postprocedural complications noted. Spinal arthrodesis is a safe, effective way to manage spinal injuries due to NAT in cases of fracture-dislocation, distraction injuries, as well as cases involving neurologic compromise.

Address for correspondence Michael J. Gigliotti, DO, MS,

Medical Center, 500 University Drive, Hershey, PA 17033,

United States (e-mail: mgigliotti19505@gmail.com).

Department of Neurosurgery, Penn State Health Milton S. Hershey

Background

Nonaccidental trauma (NAT) is estimated to occur in 2.4 to 23.8% of cases involving pediatric blunt trauma. It is often present with concomitant injuries to the head, thorax, abdomen, and musculoskeletal system.¹ Vertebral spinal injuries secondary to NAT are rare, accounting for 0 to 3% of injuries observed.¹⁻⁴ The incidence of spinal injuries and NAT have an inverse relationship to age, but a positive relationship between gender (female), race (Black), and socioeconomic status (lower median household income).^{3,5,6}

Lower thoracic and upper lumbar vertebral body compression fractures are the most common form of spine injury seen in child abuse. However, the cervical spine above C3, the cervicothoracic, and thoracolumbar junction appear to be more susceptible to fracture-dislocation injury.³ These injuries are thought to occur due to violent and forceful hyperflexion and hyperextension of the spinal column. They may result in an asymptomatic to clinically severe picture.³ However, many spine injuries are stable and without neurologic compromise, and spinal cord injuries are

published online July 29, 2021 **DOI** https://doi.org/ 10.1055/s-0041-1732792 **ISSN** 0973-0508 rare unless listhesis is present.^{7,8} Surgical management of these types of injuries is not frequently reported in the literature.⁸⁻¹⁰ We presented three cases to review the efficacy of spinal instrumentation and fixation in pediatric patients presenting as a result of NAT.

Illustrative Cases

Case 1

A 3-year-old male patient presented after jumping on a trampoline as well as a fall over six steps 2 weeks prior and complained of severe pain. The patient initially saw his pediatrician, who obtained imaging after noting a lump on his back. Imaging revealed L1 compression fracture, widening of the L1 and L2 facet joint, rotatory subluxation of L1 on L2, and kyphotic deformation of the spine with hydromyelia in the thoracolumbar region (**- Figs. 1, 2**). Physical examination was significant for bilateral lower extremity clonus. The patient underwent L1-L2 posterior instrumented fusion with open fracture reduction, utilizing pedicle screws. The patient

^{© 2021.} Neurotrauma Society of India.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (https://creativecommons.org/licenses/by-nc-nd/4.0/).

Thieme Medical and Scientific Publishers Pvt. Ltd. A-12, 2nd Floor, Sector 2, Noida-201301 UP, India



Fig. 1 AP view (left) and lateral view (right) X-ray, revealing widening of the L1 and L2 facet joint and rotary subluxation of L1 on L2.



Fig. 2 T2-weighted MRI shows kyphotic deformity of the spine with disc protrusion due to L1 compression fracture (right) and incidentally found hydromyelia (left) at T7 to T12.

Fig. 3 Lateral (left) and AP (right) view X-ray, revealing stable hardware without failure and good placement of L1 and L3 pedicle screws with correction of kyphotic deformity.

tolerated the procedure well, and his hospital course was uncomplicated. Prior to discharge, the child protective team evaluated the patient and concluded that the patient's lumbar injuries were likely nonaccidental. At his last follow-up 5 months postoperatively, the patient was doing well and was neurologically intact. His last X-ray did not have evidence of hardware malfunction or abnormalities (**~ Fig. 3**).

Case 2

A 1-year-old female presented to the hospital with new-onset, painful torticollis. Evaluation revealed bruising on her left cheek, forehead, right cheek, left hand, and left forearm. Additionally, the patient's head had 20 degrees of rotation to the right, and her neck was in a cervical collar. She did not have any focal neurological deficits. Imaging revealed a type II dens fracture with grade 4 anterolisthesis (**– Fig. 4**). The child protective team evaluated the patient at admission and determined that the patient's injuries were highly suggestive of physical abuse.

The patient was taken to the operating room, and her fracture was reduced with a halo. A posterior C1-C2 arthrodesis and fusion were performed using intralaminar wires and an autologous rib graft. Postoperative imaging revealed a slight increase in the dens' anterior displacement compared with immediate postoperative imaging (**~ Fig. 5**). However, this was reduced (grade I/II) in comparison to preoperative films (grade IV). The patient was discharged 5 days after her surgery to inpatient rehabilitation and was neurologically intact at discharge. She was discharged with her halo in place.

The patient represented to the emergency department 2 months after her discharge due to concern for one of her halo pins coming out. The halo pins were cleaned, read-justed, and she was discharged. At 4-month clinic follow-up, her halo was removed and replaced with a cervical collar, which was discontinued after 7 months. At her last follow-up at 8 months after her operation, she was neurologically intact, and her imaging did not reveal hardware failure or abnormalities.

Discussion

While nonsurgical management is commonplace to achieve stabilization, surgical intervention has proven successful in pediatric cases of spinal injury due to NAT.^{1,8-10} Open spinal reduction and fixation may be preferable in patients experiencing marked neurologic dysfunction and radiographic abnormalities, depicting fracture-dislocation, ligamentous disruption causing anterolisthesis, spinal cord signal changes, or facet disruption.



Fig. 4 Sagittal (left) and axial (right) view CT without contrast of the cervical spine, revealing grade IV anterolisthesis of the dens due to a type II dens fracture.



Fig. 5 Flexion (left), extension (right), and neutral (inset) position lateral view X-ray after placement of C1 to C2 cables and autologous rib graft. There is improvement of anterolisthesis of the dens from grade IV to grade I/II.

The two cases detailed above support the utilization of open reduction and fixation with instrumentation in certain pediatric NAT spinal injuries. A review of the available literature also demonstrates that neurologic improvement and successful bony reduction are possible in patients (**-Table 1**).^{1,8-10} All instances found in the literature utilized a posterior approach, and three of the four cases had demonstrable neurologic improvement following surgical intervention. Similar to our cases, all of the operative cases in the literature had a successful spinal fusion and did not have any complications caused by surgical intervention.

Technical challenges exist for pedicle screw fixation, including ensuring that suitably sized hardware for small, developing bony structures while ensuring placement accuracy. However, the accuracy of pedicle screw fixation in the pediatric population has been shown to be as high as 94.9%, exceeding the accuracy rate reported among similarly placed screws in adults (91.5%).¹¹ Further, adjuncts such as intraoperative image guidance is useful to achieve correct placement on the first attempt, preventing additional complications or corrective surgeries for malpositioned screws.¹² There is scant evidence of whether short- versus long-segment instrumentation and fixation is preferable in the pediatric population, and a comparative study by Li et al suggest advantages to either approach.¹³For example, short-segment fixation leads to reduced blood loss and shorter intraoperative times, while long-segment instrumentation and fixation provided relief of low back pain and improved spinal-pelvic parameters.¹³ A risk-benefit analysis in select pediatric patients should be performed to weigh the increased risk for operative complications to improve objective spinal-pelvic parameters in long-segment instrumentation versus less surgical trauma

and the potential for fewer complications in short-segment fixation.

There is a lack of longitudinal follow-up of pediatric patients undergoing spinal fixation and the potential for mechanical alteration that may occur in adulthood with respect to spinal fusion. Abe et al found that 40% of patients undergoing cervical fusion had a reduction in the anteroposterior diameter; however, this exclusively occurred in patients undergoing anterior fixation and autologous bone placement.¹⁴ Another study by Parsini et al found that 66% of patients had a normal postoperative alignment of the cervical spine which remained unchanged at follow-up.¹⁵ Interestingly, the 33% of cases that had subaxial malalignment had spontaneous sagittal improvement during long-term follow-up.¹⁵ Complications of traumatic spinal injury are rare but may include pseudomeningocele from dural tears, syrinx formation, and myelomalacia.^{16,17} These findings are likely related to the inciting traumatic event and not from direct surgical intervention.9,10 The complication rate following spinal arthrodesis is approximately 9%, with surgical site infections and wound dehiscence being the most common.17

Conclusion

Bony injury and alignment disruption, in addition to ligamentous injury, of the spinal column may require surgical fixation. The improved neurological and functional outcomes following spinal instrumentation due to nonaccidental trauma, lack of hardware malfunction, and other complications provide evidence that open reduction and internal fixation should be further studied as treatment options for the pediatric population.

Table 1 Ar	eview of the liter	ature of patients und	dergoing spinal in	istrumentation follo	owing traumatic spir	ie injury due to NAT			
Author	Patient age/ gender	Presenting neurological status	Associated injuries	Location, type of spine injury	Type of intervention	Neurological Outcome	Complications	Fusion Successful	Follow-up
Knox et al'	4-month-old/M	No function	Cervical spine EDH, cerebral edema, TBI, mul- tiple extremity fractures	Combined atlanto-occipital and atlantoaxial ligamentous injury with dens fracture	Occiput to C5 fusion and open reduction	Nonambulatory spastic quadriparesis, blindness, profound mental retardation	None	Yes	4 years
Bode et al ⁸	8-month, 30-day-old/M	Intact	R	T12 on L1 anter- olisthesis, poste- rior ligamentous disruption; T11-12 cord contusion	Open reduction, T12-L1 posterior fixation with pedicle screws and K-wires	Intact	None	Yes	14 months
Holland et al ⁹	3-week-old/F	ASIA A	Right clavicular fracture, rib fractures	C5-6 distraction injury, disrup- tion of anterior and posterior elements, bilat- eral facet joint disruption, C6 intervertebral disk avulsion	Open reduction, C4-7 posterior seg- ment fusion using rib autograft and augmentation with bone morphogenic protein	ASIA C	Posttraumatic bilateral pseu- domeningoceles, myelomalacia at C5–6	Yes	2 years
Rooks et al ¹⁰	3-month-old/F	Intact	Multiple extrem- ity fractures, rib fractures, and sternal depres- sion fracture	C5 compression fracture, anterior subluxation of C4 on C5	Total-body cast followed by C4-6 laminectomy and C3-7 posterior fusion	Bladder dysfunction, decreased left arm movement	Posttraumatic mye- lomalacia and syrinx	Yes	NR
Abbreviations:	EDH, extradural her	natoma; NAT, nonaccid€	ental trauma; TBI, tra	aumatic brain injury.					

, traumatic brain injury.
TBI
trauma;
naccidental
no,
NAT
hematoma;
extradural
EDH,
iations:

Indian Journal of Neurotrauma Vol. 19 No. 2/2022 © 2021. Neurotrauma Society of India.

Funding

None.

Conflict of Interest

None declared.

References

- 1 Knox J, Schneider J, Wimberly RL, Riccio AI. Characteristics of spinal injuries secondary to nonaccidental trauma. J Pediatr Orthop 2014;34(4):376–381
- 2 Brink FW, Gold DL, Adler B, Letson MM. Distraction injury of the thoracic spine with spinal cord transection and vascular injury in a 5-week-old infant boy: a case of child physical abuse. Pediatr Emerg Care 2017;33(3):192–197
- ³ Diamond P, Hansen CM, Christofersen MR. Child abuse presenting as a thoracolumbar spinal fracture dislocation: a case report. Pediatr Emerg Care 1994;10(2):83–86
- 4 Jauregui JJ, Perfetti DC, Cautela FS, Frumberg DB, Naziri Q, Paulino CB. Spine injuries in child abuse. J Pediatr Orthop 2019;39(2):85–89
- 5 Knox JB, Schneider JE, Cage JM, Wimberly RL, Riccio AI. Spine trauma in very young children: a retrospective study of 206 patients presenting to a level 1 pediatric trauma center. J Pediatr Orthop 2014;34(7):698–702
- 6 Zhao C, Starke M, Tompson JD, Sabharwal S. Predictors for nonaccidental trauma in a child with a fracture-A national inpatient database study. J Am Acad Orthop Surg 2020;28(4):e164–e171
- 7 Gunda D, Cornwell BO, Dahmoush HM, Jazbeh S, Alleman AM. Pediatric central nervous system imaging of nonaccidental trauma: beyond subdural hematomas. Radiographics 2019;39(1):213–228
- 8 Bode KS, Newton PO. Pediatric nonaccidental trauma thoracolumbar fracture-dislocation: posterior spinal fusion

with pedicle screw fixation in an 8-month-old boy. Spine 2007;32(14):E388–E393

- 9 Holland CM, Kebriaei MA, Wrubel DM. Posterior cervical spinal fusion in a 3-week-old infant with a severe subaxial distraction injury. J Neurosurg Pediatr 2016;17(3):353–356
- 10 Rooks VJ, Sisler C, Burton B. Cervical spine injury in child abuse: report of two cases. Pediatr Radiol 1998;28(3):193–195
- 11 Ledonio CGT, Polly DW Jr, Vitale MG, Wang Q, Richards BS. Pediatric pedicle screws: comparative effectiveness and safety: a systematic literature review from the Scoliosis Research Society and the Pediatric Orthopaedic Society of North America task force. J Bone Joint Surg Am 2011;93(13):1227–1234
- 12 Karandikar M, Mirza SK, Song K, et al. Complex pediatric cervical spine surgery using smaller nonspinal screws and plates and intraoperative computed tomography. J Neurosurg Pediatr 2012;9(6):594–601
- 13 Li Y, Ou Y, Zhu Y, He B, Xu S, Yu H. Effectiveness of short-segment fixation versus long-segment fixation for degenerative scoliosis with Cobb angle 20°~40°: a retrospective observational study. Med Sci Monit 2020;26:e923656–e1
- 14 Abe Y, Yamada K, Abumi K, Iwasaki N, Sudo H. Long-term changes in vertebral morphology after cervical spinal fusion in adolescent pediatric patients: retrospective case series with up to a minimum 12 years of follow-up. World Neurosurg 2019;122:e765–e772
- 15 Parisini P, Di Silvestre M, Greggi T, Bianchi G. C1-C2 posterior fusion in growing patients: long-term follow-up. Spine 2003;28(6):566–572, discussion 572
- 16 Liu V, Gillis C, Cochrane D, Singhal A, Steinbok P. CSF complications following intradural spinal surgeries in children. Childs Nerv Syst 2014;30(2):299–305
- 17 Abu-Bonsrah N, Goodwin CR, Ortega G, et al. Risk factors associated with short-term complications and mortality after pediatric spinal arthrodesis. Neurosurg Focus 2017;43(4):E7