THIEME OPEN ACCESS

Long-term Survival of Hip Cement Spacer: A Case Report

Sobrevida em longo prazo do espaçador cimentado para quadril: Relato de caso

Roshan Jacob¹⁰ Mackenzie Sowers¹⁰ Kelly Chandler¹⁰ Mihir Patel¹⁰ Ashish B. Shah¹⁰ Sameer Mahadeorao Naranje¹⁰

¹Department of Orthopedic Surgery, University of Alabama at Birmingham, Birmingham, Alabama, United States

Rev Bras Ortop 2024;59(6):e793-e796.

Address for correspondence Sameer Mahadeorao Naranje, MD, Department of Orthopedic Surgery, University of Alabama at Birmingham, 1313 13th Street South, Birminhgham, AL 35205, United States (e-mail: snaranje@uabmc.edu).

Abstract

Keywords

- arthroplasty, replacement, hip
- ► bone cements
- ► hip joint
- hip prosthesis
- prosthesis-related infections

Resumo

Palavras-chave

- articulação do quadril
- artroplastia de quadril
- cimentos ósseos
- infecções relacionadas às próteses
- prótese de quadril

We present a unique case of a 56-year-old male patient who ambulated on a hip cement spacer for 11 years. After hemiarthroplasty after a motor vehicle accident, the patient developed periprosthetic joint infection (PJI) several years later, and underwent stage-1 revision. With the resolution of the infection after stage 1, the patient refused the second stage due to satisfaction with the cement spacer for nearly 11 years. To our knowledge, this is the longest reported case of a cement spacer remaining in an ambulating patient. This case demonstrates the mechanical reliability of metalreinforced cement spacers, which can remain for long periods in selected patients.

Apresentamos um caso único de um homem de 56 anos que deambulou com um espaçador cimentado para quadril durante 11 anos. Após ser submetido a uma hemiartroplastia em virtude de um acidente de carro, o paciente desenvolveu uma infecção articular periprotética (IAP) vários anos depois, sendo submetido à técnica de revisão da etapa 1. Com a resolução da infecção depois da etapa 1, o paciente recusou a segunda etapa, devido à satisfação com o espaçador cimentado por quase 11 anos. Pelo que sabemos, este é o caso mais longo relatado de um espaçador cimentado em um paciente deambulando. Este caso demonstra a confiabilidade mecânica dos espaçadores cimentados reforçados com metal que podem ser mantidos por um prazo longo em pacientes selecionados.

Work carried out at the Department of Orthopedic Surgery, University of Alabama at Birmingham, Birmingham, Alabama, United States.

received December 7, 2020 accepted February 11, 2021 DOI https://doi.org/ 10.1055/s-0041-1736341. ISSN 0102-3616. © 2021. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (https://creativecommons.org/licenses/by/4.0/). Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

Introduction

While total hip arthroplasty (THA) is considered one of the most performed and successful elective surgeries in the United States, periprosthetic joint infection (PJI) is a devastating complication due to the reoperation and readmission necessary to treat the infection.¹ While the most frequently used technique to treat infected hip arthroplasty in the United States is two-stage revision using antibiotic-impregnated cement with an antibiotic spacer, the management of PJI and the treatment options should be individualized according to the needs of each patient.²

The purpose of the antibiotic spacer is to provide a higher concentration of medication directly around the area of infection, and this is achieved by placing a drug-eluting femoral stem with antibiotic-impregnated cement.² This is intended as a temporary measure, as cement hip spacers have been reported to have a dislocation rate of 7% and a fracture rate of 2%.² However, newer, prefabricated metalreinforced spacers have been shown to have much higher mechanical strength, with spacer fracture reports as low as 0% in several studies.^{3–6} The goal of reporting the present case is to describe and discuss the long-term results of an articulating hip spacer placed due to PJI, after which the patient continued to have full function of his hip for over 11 years before the second-stage revision. To our knowledge, this is the longest reported case of a retained spacer in a patient being treated for PJI.

Case Report

A 56-year-old morbidly obese male patient with a history of left hip fracture and hemiarthroplasty who presented with new hip pain. He had been involved in a motor vehicle collision (MVC) 25 years prior, and had undergone left hip hemiarthroplasty at another institution (**- Fig. 1**). The patient lived with pain for 14 years until he developed signs of PJI, presenting with elevated erythrocyte sedimentation rate (ESR, 36 mm/h) and elevated C reactive protein (CRP, 25



Fig. 1 Anteroposterior radiograph of the pelvis showing a hemiarthroplasty 14 years after surgery.



Fig. 2 Anteroposterior radiograph of the pelvis showing a Prostalac spacer immediately following stage-1 revision.

mg/L), and eventual sinus and drainage. After discussion, the patient agreed to undergo two-stage hip revision.

Using a posterior approach, an antibiotic cement and polytheline liner were placed without reaming on the acetabular side after cleaning and debriding it. Extraction of the femoral stem and head was performed with no bone loss, and the femoral canal was reamed and cleared of debris distally. After irrigation and debridement, the femoral canal was broached, and a size 3 Prostalac (Depuy-Synthes, Warsaw, IN, US) antibiotic cement stem and appropriately-sized head was placed. After hip reduction, a Hemovac drain (Zimmer Biomet, Warsaw, IN, US) was placed, and the wound was closed in a standard manner. There were no surgical complications (**>Fig. 2**). A microbiology report of fluid taken during surgery revealed *Staphylococcus lugdunensis* with no organisms isolated from blood samples.

The patient experienced gradual resolution of the infection and pain postoperatively after the appropriate antibiotic treatment. Over the following years, he continued his daily activities, including hunting and fishing. The patient refused the second stage of the revision as he was satisfied with the spacer. Eleven years later, he returned to the clinic complaining of pain in his left hip. He did not complain of any drainage or fevers. An imaging exam revealed loosening of the femoral component, and the patient agreed to second-stage revision of the THA (**-Fig. 3**). ESR (5 mm/h) and CRP (1.62 mg/L) returned to within normal limits before spacer removal, with no elevation in the white-cell count, satisfying the criteria by Parvizi et al.⁷ for absence of PJI.

A classic posterior approach was used. The stem and liner were loose and removed uneventfully. There was no evidence of bone loss (type I in the Paprosky classification). The acetabulum was reamed, and an appropriately-sized revision cup was secured with multiple screws. The remaining bone deficit was filled with antibiotic cement. The femur was prepared using the standard technique for modular revision stem. An appropriately-sized modular stem and metal head were placed and easily implanted eith the use of the standard technique after initially performing a trial reduction. The hip reduced, and the wound was closed using the standard



Fig. 3 Anteroposterior radiograph of the pelvis showing a Prostalac spacer retained for 11 years.



Fig. 4 Anteroposterior radiograph of the pelvis showing a total hip arthroplasty as stage-2 revision.

technique. The microbiology reports showed no organism growth from the cultures obtained intraoperatively. Upon his most recent follow-up, three months postoperatively, the patient was ambulating with no pain, no drainage from his operated hip, and no discrepancy regarding leg length (**-Fig. 4**).

Informed consent was obtained from the patient whose case is herein reported.

Discussion

While two-stage revision is the preferred technique for the management of PJI of the hip, the cement hip spacer is not considered the definitive implant in terms of long-term survival. These spacers are intended for short-term use only, until the infection is controlled. However, in the present case report, we discuss how a temporary cement hip spacer was tolerated by the patient over a long period of time (eleven years).

Mechanical strength and fixation are the greatest weak points of the cement hip spacer. A study⁸ involving 35 patients with a median age of 56, with surgeon-made articulating polymethylmethacrylate (PMMA) spacers, showed a mechanical failure rate (subsidence or fracture) of 45% (n = 14), with an average period of 14.5 weeks between procedures. Spacer dislocation occurred in 19% of the patients (n = 6). The authors also found that failure rates were significantly higher in younger individuals due to the higher demand placed on the spacer.⁸ Recently, surgeons have moved to using prefabricated spacers instead of surgeon-made PMMA spacers due to the superior results.^{3,4,8} Prostalac spacers in particular have shown excellent mechanical stability in several studies, with several reports of these spacers maintaining stability for up to 6 years.^{5,6,9–11} The patient presented here had a Prostalac stem and a polyethylene liner that maintained mechanical stability, and the patient maintained mobility for up to 11 years on the stem, without signs of stem fracture or dislocation.

In a retrospective study¹¹ on unplanned retained spacers, the authors reviewed 11 hips that had undergone stage 1 of the revision. Of these patients, 9 remained free of complications, while 1 patient experienced reinfection at 24 months, and the second patient developed aseptic subsidence at 72 months. The mean follow-up for the remaining 9 patients was of 48 months, during which they had no spacerrelated complaints and wished to delay surgery.¹¹

In a separate study⁶ following 25 patients with 2-year follow-up for Prostalac spacer revision, the authors found no spacer-related complications, with 68% of the patients not having undergone reimplantation at 24 months. The patients that did undergo revision in this period were younger, more physically demanding patients. It appears that a large proportion of patients can remain with a spacer for several years without serious complications. Given that many of these patients have comorbidities and contraindications to surgery, this gives both the patient and surgeon an alternate to reimplantation in high-risk cases.

To our knowledge, this is the longest reported case of a retained, prefabricated hip spacer for PJI. We suggest that these spacers can remain in selected patients with good functional long-term results. This is beneficial, as it gives surgeons a wider scope of options when it comes to treating higher-risk patients who are not suitable or unwilling to have the second-stage procedure.

Financial Support

The authors declare that there was no financial support from public, commercial, or non-profit sources.

Conflict of Interests

The authors have no conflict of interests to declare.

References

- ¹ Gundtoft PH, Pedersen AB, Varnum C, Overgaard S. Increased Mortality After Prosthetic Joint Infection in Primary THA. Clin Orthop Relat Res 2017;475(11):2623–2631
- 2 Charette RS, Melnic CM. Two-Stage Revision Arthroplasty for the Treatment of Prosthetic Joint Infection. Curr Rev Musculoskelet Med 2018;11(03):332–340
- ³ Citak M, Masri BA, Springer B, Argenson J-N, Kendoff DO. Are Preformed Articulating Spacers Superior To Surgeon-Made Articulating Spacers in the Treatment Of PJI in THA? A Literature Review. Open Orthop J 2015;9:255–261
- 4 D'Angelo F, Negri L, Binda T, Zatti G, Cherubino P. The use of a preformed spacer in two-stage revision of infected hip arthroplasties. Musculoskelet Surg 2011;95(02):115–120
- 5 Scharfenberger A, Clark M, Lavoie G, O'Connor G, Masson E, Beaupre LA. Treatment of an infected total hip replacement with the PROSTALAC system. Part 1: Infection resolution. Can J Surg 2007;50(01):24–28
- 6 Beaupre LA, Stampe K, Masson E, et al. Health-related quality of life with long-term retention of the PROSthesis of Antibiotic Loaded Acrylic Cement system following infection resolution in

low demand patients. J Orthop Surg (Hong Kong) 2017;25(02): 2309499017716257

- 7 Parvizi J, Tan TL, Goswami K, et al. The 2018 Definition of Periprosthetic Hip and Knee Infection: An Evidence-Based and Validated Criteria. J Arthroplasty 2018;33(05):1309–1314.e2
- 8 Yang FS, Lu YD, Wu CT, Blevins K, Lee MS, Kuo FC. Mechanical failure of articulating polymethylmethacrylate (PMMA) spacers in twostage revision hip arthroplasty: the risk factors and the impact on interim function. BMC Musculoskelet Disord 2019;20(01):372
- 9 Regis D, Sandri A, Magnan B, Bartolozzi P. Six-year follow-up of a preformed spacer for the management of chronically infected total hip arthroplasty. Arch Orthop Trauma Surg 2010;130(09): 1111–1115
- 10 Luk MH, Ng FY, Fu H, Chan PK, Yan CH, Chiu KY. Retention of prosthetic articulating spacer after infected hip arthroplasty as a semipermanent implant: A case report. J Orthop Trauma Rehabil 2019;26(02):105–107
- 11 Choi H-R, Freiberg AA, Malchau H, Rubash HE, Kwon Y-M. The fate of unplanned retention of prosthetic articulating spacers for infected total hip and total knee arthroplasty. J Arthroplasty 2014;29(04):690–693