



# Successful Revascularization of Left Traumatic Subclavian Artery Transection from Contralateral Thoracoacromial Artery

Harwinder Singh<sup>1</sup> Fatimah Mat Johar<sup>1</sup> Wan Azman Wan Sulaiman<sup>1</sup> Mohd Hanifah Jusoh<sup>2</sup>  
Ahmad Sukari Halim<sup>1</sup>

<sup>1</sup>Reconstructive Sciences Unit, School of Medical Sciences, Universiti Sains Malaysia, Kota Bharu, Kelantan, Malaysia

<sup>2</sup>Hand and Microsurgery Unit, Department of Orthopedics, School of Medical Sciences, Universiti Sains Malaysia, Kota Bharu, Kelantan, Malaysia

Address for correspondence Harwinder Singh, MD, MSurg, Reconstructive Sciences Unit, School of Medical Sciences, Universiti Sains Malaysia, 16150, Kota Bharu, Kelantan, Malaysia (e-mail: drharwin11@yahoo.com).

Indian J Plast Surg 2023;56:185–186.

Management of polytrauma in patients with associated traumatic limb vascular injury represents a major challenge, especially in the presence of vascular injury. Subjecting a patient to limb reperfusion surgery is against the concept of damage control orthopaedics. Life before limb is a common principle used in managing such cases. The authors describe a case of successful immediate revascularization of left subclavian artery from a branch of contralateral thoracoacromial artery, using a reverse saphenous vein graft in a polytrauma patient.

A 21-year-old gentleman with no medical illness was involved in a high-velocity road traffic accident and sustained a left traumatic subclavian artery cut. He underwent exploration after 20 hours due to the delay in transferring him to a tertiary center. Intraoperative findings revealed a complete avulsion of the left brachial plexus and a total left subclavian artery transection over the 2nd part (→ Fig. 1). The proximal stump had retracted deep under the 1st rib. No active bleeding was seen; hence, no further manipulation of the vessel was done; the area was packed with a hemostatic Gel Foam. A 30 cm great saphenous vein graft was harvested from the right lower limb and used for revascularization of the left subclavian artery. A right infraclavicular incision was made, and right subclavian artery with its branches isolated to be used as a donor artery. A subcutaneous pocket was created by blunt dissection to allow the vein graft to travel across the chest wall, and revascularization of the left subclavian artery was performed (→ Fig. 2).

The aims of management in the case of polytrauma associated with traumatic limb vascular injury are rapid,


accurate initial assessment, early and vigorous resuscitation, vascular control, and provisional stabilization of the fracture with external fixator to improve survival rate. Subclavian artery exploration and repair is indicated for all serious blunt subclavian artery trauma, except in very selective cases, such as artery injury with minimal, nonocclusive intimal dissection.<sup>1</sup> Exploration of subclavian artery is indicated in active hemorrhage, expanding hematoma, hand ischaemia and open injuries, providing that there is no indication of distal limb ischemia. Due to the rich collateral blood supply around the shoulder girdle, this is often possible.<sup>2</sup> The time-honored concepts of open arterial injury repair consist of acquiring proximal and distal vascular control, followed by reestablishing distal circulation through primary repair or graft placement restoring. This can pose a significant surgical challenge due to the vessels' position at the thoracic outlet, which penetrates the clavicle and chest wall muscles as well as the close proximity of other vital neighboring structures such as the brachial plexus, thoracic duct, phrenic and recurring laryngeal nerves, adjacent arteries and veins.<sup>3</sup> In this case, proximal stump of left subclavian artery retracted deep to 1st rib (→ Fig. 3), causing total ischemia of the whole arm which required cardiothoracic expertise that would have prolonged the operation duration. The management of polytrauma patients with traumatic limb vascular injury are a diagnostic and therapeutic challenge to the trauma team, requiring a multidisciplinary approach. Subclavian artery bypass using vein graft from contralateral subclavian artery branch is a good option in managing polytrauma

article published online  
February 25, 2022


DOI <https://doi.org/10.1055/s-0041-1740527>.  
ISSN 0970-0358.

© 2022. Association of Plastic Surgeons of India. All rights reserved. This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India



**Fig. 1** (A) Right subclavian artery and its branches identified. (B) Left subclavian artery distal stump that was revascularised with reverse GSV.



**Fig. 2** Post revascularisation with autograft.

patients with compromised circulation to upper limb, in order to increase the chances of limb salvage.

**Consent**


Consent was obtained by all participants in this study.

**Conflicts of Interest**

None declared.

**References**

- 1 McCoy DW, Weiman DS, Pate JW, Fabian TC, Walker WA. Subclavian artery injuries. *Am Surg* 1997;63(09):761-764
- 2 Damschen DD, Cogbill TH, Siegel MJ. Scapulothoracic dissociation caused by blunt trauma. *J Trauma* 1997;42(03):537-540
- 3 Franga DL, Hawkins ML, Mondy JS. Management of subclavian and axillary artery injuries: spanning the range of current therapy. *Am Surg* 2005;71(04):303-307



**Fig. 3** CT scan with 3D reconstruction showing a patent bypass graft and the retracted proximal stump of left subclavian artery.