



Falcotentorial Meningiomas: Optimal Surgical Planning and Intraoperative Challenges - Case Report and a Review

Luiz Severo Bem Junior¹ Pedro Lukas do Rêgo Aquino² Luiz Euripedes Almondes Santana Lemos¹
 Maria Amélia do Rêgo Aquino³ Marcelo Moraes Valença⁴ Hildo Rocha Cirne de Azevedo Filho¹

¹ Department of Neurological Surgery, Hospital Da Restauração, Recife, PE, Brazil

² Faculty of Medical Sciences, Universidade de Pernambuco, Recife, PE, Brazil

³ Faculty of Medical Sciences, Universidade Federal de Pernambuco, Recife, PE, Brazil

⁴ Neurosurgery Department, Hospital das Clínicas, Universidade Federal de Pernambuco, Recife, PE, Brazil

Address for correspondence: Luiz Severo Bem Junior, MD, Neuroscience Post-Graduate Program, Federal University of Pernambuco, Av. Prof. Moraes Rego, 1235 - Cidade Universitária, Recife - PE, Zipcode: 50670-901, Recife, Brazil (e-mail: luizseverobemjunior@gmail.com).

Arq Bras Neurocir 2022;41(4):e368–e374.

Abstract

Keywords

- falcotentorial meningioma
- occipital transtentorial approach
- pineal region meningioma
- outcome

Meningiomas arising from the falcotentorial junction are rare, and the selection of the optimal surgical approach is essential. We report a falcotentorial meningioma (FTM) case approached by occipitotranstentorial resection and subtotal tumor resection presenting a satisfactory clinical outcome. The present review sought to reveal the current knowledge regarding the clinical presentation, radiological imaging, and the microsurgical anatomy of FTMs as a form of improving the surgical approach. The selection of the optimal surgical approach is essential for the safe and effective removal of an FTM. Preoperative imaging analysis should identify the anatomical relations of the tumor and guide toward the least disruptive route that preserves the neurovascular structures.

Introduction

Among pineal region tumors, meningiomas are a rare entity, corresponding to between 2 and 8% of pineal tumors and to 1% of all intracranial meningiomas.^{1–3} Guttmann⁴ described the first pineal meningioma in 1930. In 1937, Araki⁵ published two cases successfully treated with a posterior transcallosal approach. These tumors originate from the posterior portion of the velum interpositum or falcotentorial union.³ Falcotentorial meningiomas (FTMs), as with other pineal region tumors, are prevalent in females.³ These tumors present a different relationship with vital neuroanatomical

structures; therefore, it is essential to decide on the ideal surgical approach.¹ It is often difficult to discriminate between FTM and velum interpositum meningiomas, even after significant advances in neuroimaging.³ However, arterial irrigation is the main difference between these two groups of tumors. The tentorial branches of the meningohypophyseal trunk usually supply FTMs, while branches of the posterior choroidal arteries 4 to 7 irrigate velum interpositum meningiomas. The surgical treatment of these tumors is not well-established in literature, since there are two main controversial issues. First, concerning validating criteria for selection of the optimal surgical approach, and second,

received
October 10, 2020
accepted
July 30, 2021

DOI <https://doi.org/10.1055/s-0041-1740593>.
ISSN 0103-5355.

© 2022. Sociedade Brasileira de Neurocirurgia. All rights reserved. This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)
 Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

whether the main infiltrated venous structures can be sacrificed to obtain a radical resection of the tumor or if they should be preserved.⁶

The present review sought to reveal the current knowledge regarding the clinical presentation and radiological imaging of FTM as a form of improving the surgical approach.

Classification

Falcotentorial meningiomas represent a subgroup of tentorial meningiomas originated from the dura mater junction depending on the posterior cerebral falx and on the convergence of both tentorial folds, projecting anterosuperiorly, inferiorly, or posteriorly depending on their growth pattern.³ Regarding their classification, the pineal region meningioma group included FTM with anterosuperior and inferior extension and velum interpositum meningiomas.¹ Yasargil et al. made one of the first tumor descriptions and eight types of tentorial meningiomas (with types T3 and T8 being falcotentorial) were identified.⁷ In 1995, Asari et al. described the classification systems for FMTs based on tumor projection that classifies the tumors as anterior, superior, posterior, and inferior types.⁸ In 2008, Bassiouni et al. classified FMTs according to tumor base location and included four types (► **Table 1**):

Type I. Type I includes tumors between both dural leaves of the cerebral falx, immediately superior to the junction of the vein of Galen with the straight sinus. In the description by Asari et al., this tumor corresponds to the anterior type.⁷

Type II. Type II originates close to junction of the vein of Galen with the straight sinus, underneath the anterior edge of the tentorium. In the description by Asari et al., this type corresponds to an inferior type.⁷

Type III. Type III is the lateralized type; therefore, it includes tumors with a dural origin in a paramedian location of one of the tentorial leaves and has a medial growing pattern toward the vein of Galen.⁷

Type IV. This type includes tumors from dural adhesion in the falcotentorial junction along the straight sinus with a posterior direction and, as described by Asari et al., it corresponds to the posterior type.⁷

On the other hand, velum interpositum meningiomas are related to the posterior portion of the double pia mater layer that covers the posterior wall of the third ventricle.⁹

Clinical Presentation

Falcotentorial meningiomas occur mostly in patients between 42 and 56 years old, similar to the posterior fossa meningiomas (44.4 years old).^{10,11} Previous studies revealed headaches as the most common symptom in pineal region meningiomas (present in 60 to 100% of the cases), followed by ataxia (43 to 62%), personality changes and bradypsychia (37 to 46%), and homonymous hemianopia (20 to 46%).⁶

Ataxia is the most frequent late symptom in type I and II tumors.⁶ Personality changes and bradypsychia were mostly associated in obstructive hydrocephalus cases.⁶ Nevertheless, bilateral visual acuity deterioration due to progressive papilledema and the subsequent atrophy of the optic nerve is extremely rare.^{8,12} Hearing impairment has been occasionally reported in patients with FTM, which improves after surgical removal.⁶ The clinical symptoms of meningioma have a propensity to occur insidiously due to their slow growth, thereby explaining the diagnosis delay to ~ 25 months, on average.³

Some authors consider the presence of alterations in ocular supraversion (Parinaud syndrome) or oculomotor cranial nerve alterations as an important differential diagnosis because this feature presents a considerably lower rate (10%) in pineal region meningiomas than in other neoplasms in the same location.^{13,14}

Surgical Anatomy

Falcotentorial meningiomas present their vascular blood supply originating from the internal carotid artery (ICA), the external carotid artery (ECA), or the posterior cerebral artery (PCA).¹⁵ ► **Table 2** summarizes the blood supply and their distal branches.

Regarding surgical treatment, understanding the relationship between meningiomas of the pineal region and the deep venous system is decisive for better results. Invasive tumor degree or the permeability of the vein of Galen and the straight sinus and the displacement of these vessels are relevant points in view of the mass effect of the tumor.^{7,15}

Occlusion and invasion of the vein of Galen and of the straight sinus were described in preoperative angiographic studies; consequently, the development of a secondary collateral venous drainage is frequently found.^{8,10,29} It can be explained due to the fact that the posterior half of the vein

Table 1 Bassiouni Falcotentorial Meningioma Classification⁷

Type of Meningioma	Origin	Venous displacement	Surgical approach
FTM type I	Posterior cerebral falx	Inferior	Occipital transtentorial
FTM type II	Inferior margin falcotentorial	Superior	Supracerebellar infratentorial
FTM type III	Paramedian tentorial	Superomedial	Paramedian supracerebellar infratentorial/Occipito transtentorial
		Inferomedial	
FTM type IV	Tentorial toward straight sinus	Contralateral	Occipitotranstentorial

Abbreviation: FTM, falcotentorial meningioma.

Table 2 Arterial supply and branches of falcotentorial meningiomas

Arterial supply	Distal Branches
ICA	Anterior choroidal artery branches
	Posterior branches from meningohipophyseal trunk
	Middle meningeal artery branches from external carotid artery
ECA	Ophthalmic artery branches
	Middle meningeal artery branches
PCA	Medial choroidal branches
	Lateral choroidal branches

Abbreviations: ECA, external carotid artery; ICA, internal carotid artery; PCA, posterior cerebral artery.

of Galen and the anterior half of the straight sinus are mostly the first vessels affected by meningiomas and are not related to the growth direction of the tumor.

Identifying vessel occlusion and the secondary collateral venous drainage is essential to evaluate the optimal surgical planning for FTMs.^{8,10} Many authors have proposed that the displacement of the deep venous system due to the direction of tumor growth is the most important characteristic when deciding the surgical approach.^{1,2,6,7} Therefore, Type I FTMs displace the venous complex inferiorly, type II superiorly, type III medially (supero/inferomedial), and type IV to the contralateral side of the meningioma.⁶

Imaging Assessment

Magnetic Resonance Imaging (MRI): Falcotentorial meningiomas are identified as a homogeneous mass, predominantly oval, with homogeneous enhancement after the administration of gadolinium-T1 contrast.²⁷ In T1 sequences, they are usually hypo/isointense and iso/hyperintense in T2 sequences. Peritumoral edema was observed in 85% of the patients in a case series, whereas obstructive hydrocephalus, mainly revealed in infratentorial lesions, was observed in ~ 32%. In this study, the average diameter of the lesion was 52 mm.³⁰

In the preoperative period, knowledge about the patency of the rectal sinus and the displacement of the Galen vein and internal cerebral veins are fundamental to optimal surgical planning, and MRI can provide this information.⁶

Furthermore, MRI is more accurate in enabling the visualization of the straight sinus and the vein of Galen when compared with conventional angiography, since it was able to identify these structures in ~ 20% of the patients.⁶

Cerebral Angiography: Currently, some authors do not indicate cerebral angiography routinely in FTMs because angiomagnetic resonance has advanced in the last years.⁶ The direction of the displacement of the medial choroidal artery is one of the most important angiographic features that can be used to differentiate FTMs from other pineal

masses.^{5,8,31,32} Therefore, medial choroidal artery displaces anteriorly in FTMs and superior and posteriorly in the other pineal tumors.^{5,32}

Surgical Approach

The objective of the surgical treatment of FTMs is to achieve a macroscopically complete resection of the lesion, to relieve or solve the neurologic/clinical symptoms, and to acquire a tissue sample for a definitive diagnosis. Therefore, the recommendation to perform an intraoperative biopsy is established, paving the way to obtaining a differential diagnosis from other pineal tumors (for example, germinoma) for which the management may be different, aiming a partial resection after adjuvant therapy.³³

Classically, preoperative artery embolization is an important adjuvant treatment for meningiomas. However, most FTMs present a short artery caliber, which can difficult preoperative artery embolization.¹⁵

The anatomical relation between the tumor and the deep venous complex of Galen is the most crucial factor when choosing the surgical approach. Five surgical approaches are described for these tumors:

Transtentorial/Transfalcine Occipital Approach. This is the most frequently used approach for pineal meningiomas. It was first described by Poppen 1968 and was improved afterwards by Jamieson in 1971.^{34,35} This approach is more specifically advised in type I and IV FTMs, which originate from the posterior falx immediately above the junction of vein of Galen and the straight sinus. In this location, the growth of the tumor displaces posterior and inferiorly the deep venous complex (► **Table 2**).

The occipital interhemispheric approach is used to reach tumors with a mostly supratentorial and a smaller infratentorial extension. The transfalcine/transtentorial route is used for the removal of the contralateral supratentorial/infratentorial component. The occipital lobe is also gravity-dependent positioned bearing the largest component of the tumor.

This surgical approach is performed with the patient in a prone or three-quarters prone position with the side of the approach depending on the lateralization of the tumor, avoiding excessive occipital retraction (► **Fig. 1A**).²⁸

However, this surgical approach has some disadvantages, such as an increased risk of visual cortex damage (due to




Fig. 1 A. The three-quarter prone position in the transtentorial/transfalcine occipital approach. B. Surgical view of the supracerebellar infratentorial approach. The schematic illustration belongs to the authors.

cortex retraction), possible trochlear nerve damage at the tentorial opening, and a limitation of the contralateral view of the tumor margin.^{28,34,36–38}

Supracerebellar infra-tentorial approach: This is the second most frequently used approach for pineal meningiomas and was the first surgical approach described for these tumors, being reported by Krause et al. in 1913.³⁹ This approach is recommended for type II and III FTM according to the Bassiouni classification. It uses the space between the upper surface of the cerebellum and the tentacle (► **Fig. 1B**). Regarding the position of the patient, the sitting position promotes greater cerebral relaxation, but is inappropriate for patients with cardiac comorbidities. Thus, surgery with the patient in a semiseated position seems to be promising in recent studies.^{19,33,40,41} A suboccipital craniotomy is performed and centered on the torcula, with a consequent exposure of the two transverse sinuses and retraction of the cerebellum. Currently, some studies suggest that a supracerebellar infratentorial approach could be difficult in patients presenting a steep tentacle and, in these cases, the occipital transtentorial/transfalcine approach seems to be better.^{42,43}

Occipital bitranstentorial/falcine approach: This approach was first described by Sekhar et al. in 2002.^{15,44} This surgical approach is recommended particularly for large pineal tumors with supra- and infratentorial extension or for those with a significant lateral extension. This technique can be described as a modification of the occipital transtentorial approach, with the division of the falx and the bilateral tentorium. An important advantage of this approach is the better exposition of the contralateral quadrigeminal region.⁴⁵

Anterior interhemispheric trans-splenial approach: This approach was recently described by Yağmurlu et al. and examined cadaveric heads. This surgical approach has several advantages and provides an effective surgery, particularly for patients with supratentorial pineal region tumors with superior extension; for tentorial tumors that displace the deep venous system inferiorly; or for those that originate from the splenium of the corpus callosum.²⁶ Hendricks et al. described that the anterior interhemispheric approach provides the safest route for accessing the largest portion of the tumor while protecting the deep venous system; however, in this case, the tumor was subtotally removed because of the adherence along the vein of Galen (► **Table 3**).²⁶

Posterior interhemispheric retrocallosal transfalcine approach: In 2016, Liu et al. reported this surgical approach using endoscopic assistance to resect a superiorly positioned FTM. In this study, the Galen vein complex was preserved.¹⁸ Lopez-Gonzales et al. described two cases using this approach and achieved a subtotal and a total resection in those cases, respectively²⁰ (► **Table 3**). However, further studies about this technique are still required.

Adjuvant Radiosurgery

Adjuvant treatments for FTM are based on histopathological grades and surgical resections. The surgical resection margins are classified using the Simpson classification.¹⁷ Adjuvant

radiosurgery is recommended for high-grade histopathological tumor lesions and tumors without Simpson I resection.⁴³

Complications and Prognosis

Despite surgical techniques and neuroimaging advances, risks and complications are inherent to FTM procedures, in view of the depth of the location and relationships to important vascular structures. Currently, the attempt to perform a radical tumor resection is not well-established in the literature.¹⁶ Considering that aggressive resections are related to high morbidity rates, a conservative strategy, with subtotal resections, seems to be beneficial in most cases, decreasing a severe neurological deficit rate. The adherence or invasion into the brainstem or the deep venous system by the tumor is an essential factor for a subtotal resection.¹⁶ Surgical mortality was reported in up to 23% of patients, and morbidity, with permanent neurological damage, was recorded in up to 50%. Homonymous hemianopia is the most frequent postoperative deficit.¹⁶ In addition, transient deficits related to the deviation from the conjugated gaze (Parinaud syndrome) and diplopia, being justified by manipulation during the resection of the structure of the dorsal midbrain, such as the collicular plate. This deficit is mostly reversible a few days after the surgery, especially if the arachnoid plan is respected. Permanent cortical amaurosis has been described as a possible surgical complication related to prolonged occipital lobe retraction time.¹⁶ Hemiparesis, hemidysesthesia, tetraparesis, neglect syndrome, and postoperative intraventricular hemorrhage were described in the literature.^{16,43} Our patient developed superior temporal homonymous quadrantanopia in the immediate postoperative period, permanently, without other neurological symptoms. ► **Table 3** summarizes the most relevant reports in the literature from the past 10 years, focusing on the surgical approach, outcomes, and complications.

Conclusion

The selection of an optimal surgical approach is essential for the safe and effective removal of an FTM. Preoperative imaging analysis should identify the anatomical relations of the tumor and guide the least disruptive route that preserves the neurovascular structures.

Ethical Approval

For this type of study, formal consent is not required.

Informed Consent

The authors state that the patient has given their written informed consent for publication of data and images.

Previous Presentation

None.

Disclosure of Funding and Financial Support

None.

Table 3 Review of the most relevant reports in literature from the past 10 years

Study	Number of patients	Approaches (%)	Outcome	Morbidity or complication
Okada et al., 2020 ¹⁶	4	Parieto-occipital interhemispheric transfalcine, transbitentorial approach	GTR	none
Hendricks et al., 2019 ¹⁷	1	Anterior interhemispheric	STR	ns
Lopez-Gonzales et al., 2019 ¹⁸	2	Retractorless interhemispheric	GTR in two times (1/2)	none
			STR (1/2)	
Zhao et al., 2019 ¹⁹	17	SCIT (4/17)	GTR (12/17)	Permanent visual field deficits (2/17)
		O-TT (4/17)		Transient visual field deficits (4/17)
		AITS (3/17)	STR (5/17)	Hemiparesis (2/17)
		PVT (1/17)		Hemidysesthesia (1/17)
		Torcular (2/17)		Cerebellar hematoma (1/17)
Garcia et al., 2019 ²⁰	2	SCIT (1/2)	GTR (2/2)	none
		Parieto-occipital interhemispheric (1/2)		
Mendez-Rosito, 2019 ²¹	1	SCIT	GTR	none
Talachchi et al., 2018 ²²	16	Occipital (5/16)	GTR (8/16)	Hematoma (2/16)
		Parieto-occipital (5/16)		Hydrocephalus (1/16)
		Occipitosuboccipital (3/16)	STR (8/16)	Pleural effusion (1/16)
		Suboccipital (3/16)		Tracheostomy (1/16)
		Cranial nerve deficit (1/16)		
Couldwell, 2017 ²³	1	Occipital interhemispheric	GTR	ns
Gomes et al., 2017 ²⁴	1	Posterior interhemispheric	ns	Neglectsyndrome
Hong et al., 2017 ²⁵	11	Bioccipital transtentorial (1/11)	GTR (10/11)	Occipitocerebellar hemorrhage (1/11)
		O-TT (4/11)		
		Occipital (4/11)	STR (1/11)	Transient visualfield deficit (1/11)
		Occipitoparietal (2/11)		
Liu et al., 2016 ²⁶	1	Posterior interhemispheric retrocallosal transfalcine	STR	none
Liu, 2016 ¹⁸	1	Combined bi-occipital suboccipital transsinus transtentorial	STR	ns
Nowak et al., 2014 ²⁷	4	O-TT (4/4)	GTR (3/4)	Temporary homonymous hemianopsia (4/4)
				Upward-gaze palsy (1/4)
			STR (1/4)	Postoperative intraventricular hemorrhage (1/4)
Bahari et al., 2014 ²⁰	2	O-TT (2/2)	STR (2/2)	none
Qiu et al., 2014 ²⁸	15	O-TT (15/15)	GTR (11/15)	Homonymous hemianopia (2/15)
			STR (4/15)	Parinaud syndrome (1/15)

Abbreviations: AITS, anterior interhemispheric trans-splenic; GTR, gross total resection (Simpson I and II); ns, not specified; O-TT, occipital transtentorial/transfalci; PTV, parietal transventricular; STR, subtotal resection (Simpson III and IV).

Contributions of the Authors

1. Bem Junior L. S.: Conceptualization, data curation, formal analysis, visualization, writing, reviewing, and editing.
2. Aquino P. L. R.: Data curation, writing of the original draft, reviewing and editing.
3. Lemos L. E. A. S.: Data curation, writing of the original draft, reviewing and editing.
4. Aquino M. A. R.: Data curation, writing, reviewing, and editing.
5. Valença M. M.: Conceptualization, writing, reviewing, and editing, supervision.
6. Azevedo Filho H. R. C.: Conceptualization, writing, reviewing, and editing, supervision.

Funding

None.

Conflict of Interests

The authors have no conflict of interests to declare.

Acknowledgments

The present research was carried out at the Hospital da Restauração, Recife, state of Pernambuco, Brazil. We thank our colleagues from the Hospital da Restauração who provided insight and expertise that greatly assisted in the research. We are also grateful to the advanced neurosurgery support unit (ANSU) from the Hospital da Restauração for all the care provided in the postoperative period of the patient.

References

1. Behari S, Das KK, Kumar A, et al. Large/giant meningiomas of posterior third ventricular region: falcotentorial or velum interpositum? *Neurol India* 2014;62(03):290–295. Doi: 10.4103/0028-3886.136934
2. Hong CK, Hong JB, Park H, et al. Surgical treatment for falcotentorial meningiomas. *Yonsei Med J* 2016;57(04):1022–1028. Doi: 10.3349/ymj.2016.57.4.1022
3. Yaşar S, Kirik A. Surgical Management of Giant Intracranial Meningiomas. *Eurasian J Med* 2021;53(02):73–78
4. Guttman E. Zur pathologie und Klinik der Meningiome. *Z Gesamte Neurol Psychiatr* 1930;123:606–625
5. Ito J, Kadekaru T, Hayano M, Kurita I, Okada K, Yoshida Y. Meningioma in the tela choroidea of the third ventricle: CT and angiographic correlations. *Neuroradiology* 1981;21(04):207–211. Doi: 10.1007/BF00367342
6. Bassiouni H, Asgari S, König HJ, Stolke D. Meningiomas of the falcotentorial junction: selection of the surgical approach according to the tumor type. *Surg Neurol* 2008;69(04):339–349, discussion 349. Doi: 10.1016/j.surneu.2007.02.029
7. Sekhar LN, Goel A. Combined supratentorial and infratentorial approach to large pineal-region meningioma. *Surg Neurol* 1992;37(03):197–201. Doi: 10.1016/0090-3019(92)90230-k
8. Asari S, Maeshiro T, Tomita S, et al. Meningiomas arising from the falcotentorial junction. Clinical features, neuroimaging studies, and surgical treatment. *J Neurosurg* 1995;82(05):726–738. Doi: 10.3171/jns.1995.82.5.0726
9. Tubbs RS, Louis RG Jr, Wartmann CT, et al. The velum interpositum revisited and redefined. *Surg Radiol Anat* 2008;30(02):131–135. Doi: 10.1007/s00276-007-0293-2
10. Quinones-Hinojosa A, Chang EF, McDermott MW. Falcotentorial meningiomas: clinical, neuroimaging, and surgical features in six patients. *Neurosurg Focus* 2003;14(06):e11. Doi: 10.3171/foc.2003.14.6.11
11. Roberti F, Sekhar LN, Kalavakonda C, Wright DC. Posterior fossa meningiomas: surgical experience in 161 cases. *Surg Neurol* 2001;56(01):8–20, discussion 20–21. Doi: 10.1016/s0090-3019(01)00479-7
12. Raco A, Agrillo A, Ruggeri A, Gagliardi FM, Cantore G. Surgical options in the management of falcotentorial meningiomas: report of 13 cases. *Surg Neurol* 2004;61(02):157–164, discussion 164. Doi: 10.1016/S0090-3019(03)00573-1
13. Sung DI, Harisiadis L, Chang CH. Midline pineal tumors and suprasellar germinomas: highly curable by irradiation. *Radiology* 1978;128(03):745–751. Doi: 10.1148/128.3.745
14. Piatt JH Jr, Campbell GA. Pineal region meningioma: report of two cases and literature review. *Neurosurgery* 1983;12(04):369–376. Doi: 10.1227/00006123-198304000-00002
15. Quiñones-Hinojosa A, Chang EF, Chaichana KL, McDermott MW. Surgical considerations in the management of falcotentorial meningiomas: advantages of the bilateral occipital transtentorial/transfalcine craniotomy for large tumors. *Neurosurgery* 2009;64(5, Suppl 2):260–268, discussion 268. Doi: 10.1227/01.NEU.0000344642.98597.A7
16. Okada T, Miyahara K, Tanino S, et al. Parieto-Occipital Interhemispheric Transfalcine, Trans-Bitentorial Approach for Radical Resection of Falcotentorial Meningiomas. *J Neurol Surg A Cent Eur Neurosurg* 2020;81(04):355–361
17. Hendricks BK, Spetzler RF. Right Anterior Parasagittal Craniotomy and Interhemispheric Approach for Resection of a Falcotentorial Junction Meningioma: 2-Dimensional Operative Video. *Oper Neurosurg (Hagerstown)* 2019;17(06):E237–E238. Doi: 10.1093/ons/onz258
18. Lopez-Gonzalez MA, Jaeger A, Kaplan B, et al. Retractorless interhemispheric transtentorial approach for large lesions in the posterior incisural space. *Surg Neurol Int* 2019;10:130. Doi: 10.25259/SNI-117-2019
19. Zhao X, Belykh E, Przybylowski CJ, et al. Surgical treatment of falcotentorial meningiomas: a retrospective review of a single-institution experience. *J Neurosurg* 2019 (August):1–12. Doi: 10.3171/2019.4.JNS19208
20. Blasco García de Andoain G, Delgado-Fernández J, Penanes Cuesta JR, Gil-Simoes R, Frade-Porto N, Sánchez MP. Meningiomas Originated at the Falcotentorial Region: Analysis of Topographic and Diagnostic Features Guiding an Optimal Surgical Planning. *World Neurosurg* 2019;123:e723–e733. Doi: 10.1016/j.wneu.2018.12.013
21. Mendez-Rosito D. The Supra/Infra Transtentorial Transfalcine Approach for the Removal of a Falcotentorial Meningioma: 2-Dimensional Video. *Oper Neurosurg (Hagerstown)* 2019;16(06):E166–E167
22. Shukla D, Behari S, Jaiswal AK, Banerji D, Tyagi I, Jain VK. Tentorial meningiomas: operative nuances and perioperative management dilemmas. *Acta Neurochir (Wien)* 2009;151(09):1037–1051
23. Couldwell WT. Left occipital craniotomy for resection of falcotentorial meningioma. *Neurosurg Focus* 2017;43(VideoSuppl2):V9
24. Gomes D, Fonseca M, Garrotes M, et al. Corpus callosum and neglect syndrome: Clinical findings after meningioma removal and anatomical review. *J Neurosci Rural Pract. Medknow Publications* 2017;8(01):101–106
25. Larson JJ, Tew JM Jr, Wiot JG, de Courten-Myers GM. Association of meningiomas with dural “tails”; surgical significance. *Acta Neurochir (Wien)* 1992;114(1-2):59–63
26. Liu JK, Cohen MA. Endoscopic-assisted posterior interhemispheric retrocallosal transfalcine approach for microsurgical resection of a pineal region falcotentorial meningioma: operative video and

- technical nuances. *Neurosurg Focus* 2016;40Video(Suppl 1):2016.1, 15453
- 27 Nowak A, Dziedzic T, Czernicki T, Kunert P, Marchel A. Falcotentorial and velum interpositum meningiomas: two distinct entities of the pineal region. *Neurol Neurochir Pol* 2014;48(06):397–402
 - 28 Qiu B, Wang Y, Ou S, Guo Z, Wang Y. The unilateral occipital transtentorial approach for pineal region meningiomas: a report of 15 cases. *Int J Neurosci* 2014;124(10):741–747. Doi: 10.3109/00207454.2013.878341
 - 29 Goto T, Ohata K, Morino M, et al. Falcotentorial meningioma: surgical outcome in 14 patients. *J Neurosurg* 2006;104(01):47–53. Doi: 10.3171/jns.2006.104.1.47
 - 30 Talacchi A, Biroli A, Hasanbelliu A, Locatelli F. Surgical Management of Medial Tentorial Meningioma: Falcotentorial and Torcular. *World Neurosurg* 2018;115:e437–e447. Doi: 10.1016/j.wneu.2018.04.066
 - 31 Okami N, Kawamata T, Hori T, Takakura K. Surgical treatment of falcotentorial meningioma. *J Clin Neurosci* 2001;8(Suppl 1):15–18. Doi: 10.1054/jocn.2001.0870
 - 32 Carey JP, Fisher RG, Pelofsky S. Tentorial meningiomas. *Surg Neurol* 1975;3(01):41–44
 - 33 Lozier AP, Bruce JN. Meningiomas of the velum interpositum: surgical considerations. *Neurosurg Focus* 2003;15(01):E11. Doi: 10.3171/foc.2003.15.1.11
 - 34 Poppen JL. The right occipital approach to a pinealoma. *J Neurosurg* 1966;25(06):706–710. Doi: 10.3171/jns.1966.25.6.0706
 - 35 Jamieson KG. Excision of pineal tumors. *J Neurosurg* 1971;35(05):550–553. Doi: 10.3171/jns.1971.35.5.0550
 - 36 Reid WS, Clark WK. Comparison of the infratentorial and transtentorial approaches to the pineal region. *Neurosurgery* 1978;3(01):1–8. Doi: 10.1227/00006123-197807000-00001
 - 37 Maselli G, De Paulis D, Ricci A, Galzio RJ. Posterior cranial fossa tumors: Results and prognostic factors in a consecutive series of 14 operated patients by occipital transtentorial approach. *Surg Neurol Int* 2012;3(01):85. Doi: 10.4103/2152-7806.99911
 - 38 Zaidi HA, Elhadi AM, Lei T, Preul MC, Little AS, Nakaji P. Minimally Invasive Endoscopic Supracerebellar-Infratentorial Surgery of the Pineal Region: Anatomical Comparison of Four Variant Approaches. *World Neurosurg* 2015;84(02):257–266. Doi: 10.1016/j.wneu.2015.03.009
 - 39 Quest DO, Kleriga E. Microsurgical anatomy of the pineal region. *Neurosurgery* 1980;6(04):385–390. Doi: 10.1227/00006123-198004000-00006
 - 40 Sonabend AM, Bowden S, Bruce JN. Microsurgical resection of pineal region tumors. *J Neurooncol* 2016;130(02):351–366
 - 41 Stein BM. The infratentorial supracerebellar approach to pineal lesions. *J Neurosurg* 1971;35(02):197–202. Doi: 10.3171/jns.1971.35.2.0197
 - 42 Hart MG, Santarius T, Kirollos RW. How I do it–pineal surgery: supracerebellar infratentorial versus occipital transtentorial. *Acta Neurochir (Wien)* 2013;155(03):463–467. Doi: 10.1007/s00701-012-1589-5
 - 43 Soffar H, Alsawy MF. Assessment of necessity of neuronavigation in localization of calvarial extra-axial lesions in the setting of limited resources. *Chin Neurosurg J* 2021;7(01):35
 - 44 Gusmão S, Oliveira MM, Arantes A, Ulhoa TH, Morato EG. Occipital bi-transtentorial/falcine approach for falcotentorial meningioma: case report. *Arq Neuropsiquiatr* 2006;64(01):136–138. Doi: 10.1590/S0004-282X2006000100029
 - 45 Yağmurlu K, Zaidi HA, Kalani MYS, Rhoton AL, Preul MC, Spetzler RF. Anterior interhemispheric transsplenic approach to pineal region tumors: anatomical study and illustrative case. *J Neurosurg* 2018;128(01):182–192. Doi: 10.3171/2016.9.JNS16279