

Systematic Comparison of Different Meta-analyses, Systematic Reviews and HTA Reports on Cervical Cancer Screening based on Cytology or HPV Test

Systematischer Vergleich verschiedener Metaanalysen, systematischer Reviews und HTA-Berichte zum Zervixkarzinomscreening mittels Zytologie oder HPV-Test

Authors

M. Jentschke, P. Hillemanns

Affiliation

Klinik für Frauenheilkunde und Geburtshilfe, Medizinische Hochschule Hannover, Hannover, Germany

Key words

- cervical cancer screening
- HPV
- cytology
- dysplasia

Schlüsselwörter

- Zervixkarzinomscreening
- HPV
- Zytologie
- Dysplasie



Deutsche Version unter:
<http://dx.doi.org/10.1055/s-0042-112457>

received 19. 4. 2016
 revised 17. 6. 2016
 accepted 14. 7. 2016

Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-112457>
 Geburtsh Frauenheilk 2016; 76: 1081–1085 © Georg Thieme Verlag KG Stuttgart · New York · ISSN 0016-5751

Correspondence

Dr. Matthias Jentschke
 Klinik für Frauenheilkunde und Geburtshilfe
 Medizinische Hochschule Hannover
 Carl-Neuberg-Straße 1
 30625 Hannover
 Germany
 Jentschke.Matthias@mh-hannover.de

Abstract



Introduction: Cervical cancer screening programs all over the world are being radically revised; preventive screening is increasingly based on detecting infection with human papillomavirus (HPV). This change was driven by large international studies (comparing cytological and HPV-based screening), which were then summarized in systematic reviews or HTA reports. We carried out a systematic comparison of these meta-analyses, comparing their contents, quality, results and recommendations.

Material and Methods: A systematic search in MEDLINE identified eight meta-analyses. Seven more papers were also included in our analysis. The quality of these publications was reviewed using the AMSTAR criteria and presented in tabular form.

Results: There were significant differences with regard to the primary studies included in the meta-analyses, the quality of the meta-analyses, the endpoints, and the outcomes and recommendations based on these endpoints.

Conclusion: Different meta-analyses on the same or similar issues sometimes came to quite different conclusions because they used different approaches. This is quite significant because the underlying primary data did not change. The data indicated that HPV-based screening was superior. The heterogeneity of the meta-analyses affects the decisions taken by policymakers in the healthcare system.

Zusammenfassung



Einleitung: Das Zervixkarzinomscreening befindet sich weltweit im Umbruch und vollzieht einen Wandel hin zu einer Vorsorge, die auf dem Nachweis einer Infektion mit humanen Papillomviren (HPV) basiert. Dieser Wandel basiert auf großen internationalen Studien (Vergleich von Zytologie- und HPV-Screening), die in der Regel in systematischen Übersichtsarbeiten oder HTA-Berichten zusammengefasst werden. Diese Arbeit stellt einen systematischen Vergleich dieser Metaanalysen dar im Hinblick auf Inhalt, Qualität, Ergebnisse und Empfehlungen.

Material und Methoden: Durch eine systematische Suche in MEDLINE konnten 8 Metaanalysen identifiziert werden. Sieben weitere Arbeiten wurden in die Analyse eingeschlossen. Die Qualität wurde mittels AMSTAR überprüft und tabellarisch dargestellt.

Ergebnisse: Es zeigten sich teilweise deutliche Unterschiede im Hinblick auf die eingeschlossenen Primärstudien, die Qualität der Metaanalysen, die Endpunkte und auch die Ergebnisse und die daraus resultierenden Empfehlungen.

Schlussfolgerung: Unterschiedliche Metaanalysen zu gleichen oder ähnlichen Fragestellungen liefern aufgrund unterschiedlicher Ansätze teilweise auch deutlich unterschiedliche Ergebnisse. Dies ist beachtlich, da sich die zugrunde liegenden primären Daten nicht ändern. Diese sprechen für eine Überlegenheit des HPV-basierten Screenings. Die Entscheidungsträger im Gesundheitssystem werden so durch die heterogene Durchführung der Metaanalysen in ihren Entscheidungen beeinflusst.

Introduction

Opportunistic cervical cancer screening programs based on annual Pap smears were rolled out in Germany in 1971 and led to a significant decrease in the incidence of cervical cancer by 60–70% [1]. Despite this success, this approach has certain weaknesses. For example, it is currently not possible to comprehensively document and evaluate participation rates, although this significantly affects the future incidence of invasive carcinoma [2]. Other disadvantages of the current screening program for cervical cancer include the lack of a uniform workup procedure and the absence of continuous quality controls.

In addition to the proposal to introduce an organized screening program, the current discussion on the need to revise cervical cancer prevention demanded by politicians (National Cancer Plan and the Law on the Early Detection and Registration of Cancer [*Krebsfrüherkennungs- und -registergesetz*]) has also focused on the question whether screening should be primarily based on testing for HPV. The new S3 guideline “Prevention of Cervical Cancer” has also focused on this issue (<http://leitlinienprogramm-onkologie.de/Zervixkarzinom-Prävention.89.0.html>).

These considerations were prompted by several large randomized controlled studies carried out in recent years which appeared to show that screening for HPV was superior to early detection based on cytology (summarized in [3]). In many countries, these studies served as the basis for a revision of cervical cancer screening programs. In a number of countries, healthcare policymakers requested that systematic reviews or meta-analyses be carried out to compare HPV with cytology for the prevention of cervical cancer and subsequently based their healthcare policy recommendations on the outcomes of these analyses.

Interestingly, these meta-analyses often differed significantly from one another; the different analyses included or excluded different primary studies and evaluated and assessed effects and overall results differently. This study aimed to review the methodological differences between the various meta-analyses in a systematic reappraisal of the currently available aggregated evidence on the issue of HPV vs. cytology and to compare the different results of these analyses in order to show the basis for the different recommendations issued in different countries.

Material and Methods

Search strategy

A comprehensive systematic search* for systematic reviews of evidence was carried out in MEDLINE via PubMed (<http://www.ncbi.nlm.nih.gov/pubmed>). The search criteria were, first and foremost, to find systematic reviews which compared cervical cancer screening using HPV test with cytological screening. This study also looked at systematic reviews which investigated individual aspects of HPV-based screening for cervical cancer. Non-systematic reviews, guidelines, best practice recommendations or primary studies were not included.

* Search strategy: PubMed, 24.09.2015. Search string: (cytology OR pap) AND (hpv OR papillomavirus) AND (“cervix neoplasm” OR “cervical intra-epithelial neoplasia” OR “cervix dysplasia” OR “cin” OR “cervical cancer”). Limits: meta-analysis, systematic reviews. Number of hits: 216

Table 1 Formulation of the benefit analysis by the IQWiG.

IQWiG assessment	German guidelines	G-BA level of recommendation
Evidence of benefit	LoE Ia	Strongly recommended
Indications	LoE Ia	Recommended
Signs	LoE Ia–b	Qualified recommendation
No signs	< LoE I	Not recommended

Analysis of the quality of the study

After summarizing the contents (the primary studies included, the method, and the results) the methodological quality of the systematic reviews were evaluated and compared using the so-called AMSTAR criteria (http://amstar.ca/Amstar_Checklist.php) [4].

Results

Literature search

After reviewing the abstracts, nine publications (eight independent studies; one of the studies was published in both a long and a short version) were identified out of the original 216 hits which complied with the search criteria [5–13]. Seven other systematic reviews which were not identified in the search – inter alia, two papers by the Institute for Quality and Efficiency in Healthcare (IQWiG) on the “Benefit assessment of HPV test in primary screening for cervical cancer” published in 2011 and 2014 (Table 1) as well as the analysis by Jos Kleijnen compiled for the S3 guideline on the prevention of cervical cancer – were also included in our review ([14–20]; a detailed summary of their contents is available in the online supplement. Only 13 of the included 15 meta-analyses focused on comparing HPV test with cytology in primary screening. Peirson et al. compared cytology, liquid-based cytology and HPV test with no screening [10], and Luu et al. evaluated the sensitivity and specificity of two different HPV detection methods (Hybrid Capture 2 and PCR) [6].

Study quality

The quality of the studies was compared using the AMSTAR criteria and is summarized in Table 2. The difference in the quality of the studies was sometimes quite significant, although it is important to point out that many methodological aspects may often not find mention in the final publication and the study was therefore evaluated as though these methodological aspects did not exist, even though it is possible that they were taken into consideration. The highest quality studies were the two studies by the IQWiG, along with a few others.

There were also considerable differences in the choice of primary endpoints:

- ▶ histologically verified CIN 3+ [5]
- ▶ relative rate of detection, relative specificity, relative PPV [11]
- ▶ cervical cancer mortality, incidence of cervical cancer [10]
- ▶ CIN 2+, CIN 3+ [9]
- ▶ relative rate of CIN 2, CIN 2+, CIN 3+; cervical cancer mortality, and incidence of cervical cancer [7]
- ▶ overall survival, disease-specific (tumor-specific) survival, incidence of invasive cervical cancer, incidence of CIN 3/CIS, incidence of CIN 3+, injury arising directly or indirectly from screening, health-related quality of life, psychosocial aspects [15]

Table 2 Evaluation of systematic reviews using the AMSTAR criteria (http://amstar.ca/Amstar_Checklist.php).

	Bouchard-Fortier 2014	Pileggi 2014	Peirson 2013 See full text for additional information	Pan 2014	Patanwala 2013	Luu 2013	Vesco 2011 Whitlock 2011	Murphy 2012 (Online Suppl.)	MaHTAS 2011	IQWiG 2012	IQWiG 2014	KCE REPORT 238, 2015	Arbyn 2012	MSAC 2013	Jos Kleijnen 2014
1. Was an "a priori" design provided?	yes	no	yes	no	yes	yes	yes	no	yes	yes	yes	yes (update of older meta-analyses)	yes (update of older meta-analyses)	nein	ja
2. Was there duplicate study selection and data extraction?	yes	yes	yes	no	yes	yes	yes	no	yes	yes	yes	no	no	no	yes
3. Was a comprehensive literature search performed?	yes	yes	yes	no	yes	yes	yes	yes	yes	yes	yes	no	yes	no	yes
4. Was the status of publication (i.e. grey literature) used as an inclusion criterion?	no	no	yes	no	no	no	yes	no	no	yes	no	no	no	no	no
5. Was a list of studies (included and excluded) provided?	no	no	no	no	no	no	yes	no	yes	Yes	yes	no	no	no	yes
6. Were the characteristics of the included studies provided?	yes	yes	yes	yes	yes	yes	yes	yes	yes	Yes	yes	yes	no	no	yes
7. Was the scientific quality of the included studies assessed and documented?	yes	no	yes	no	yes	no	yes	no	yes	Yes	yes	no	no	no	yes
8. Was the scientific quality of the included studies used appropriately in formulating conclusions?	yes	no	yes	no	yes	no	yes	no	yes	Yes	yes	no	no	no	yes
9. Were the methods used to combine the findings of studies appropriate?	yes	yes	yes	yes	yes	yes	yes	yes	yes	Yes	yes	yes	yes	no	yes
10. Was the likelihood of publication bias assessed?	yes	yes	yes	no	no	no	no	no	no	No	no	no	no	no	yes
11. Was the conflict of interest included?	no	no	no	no	no	no	yes	no	no	No	no	no	no	no	no
AMSTAR score	8/11	5/11	9/11	2/11	7/11	5/11	10/11	3/11	8/11	9/11	8/11	3/11	3/11	0/11	9/11

Inclusion of primary studies

It was also very noticeable that reviews often looked at very different primary studies. Although most reviews included the best known European RCTs (NTCC1 and 2, Swedescreen, ARTISTIC, POBASCAM, Finland study), the Indian study by Sankaranarayanan et al. published in 2009 [21] and the Canadian CCCaST [22] were usually not included. The heterogeneity of the primary studies was also interpreted differently by different authors. This heterogeneity prompted some authors of reviews to dispense completely with any meta-analytic pooling of individual trials (e.g. Patanwala 2013 [9]). However, it was not possible to identify a link to the quality of the investigated meta-analyses determined using AMSTAR, e.g. that high-quality meta-analyses tend to be critical of the level of evidence.

Results of the meta-analyses

The different authors of the various meta-analyses also came to different conclusions and made different recommendations. To take one example, the joint study by Vesco (full report in 2011) and Whitlock (short summary in 2011), which served as the basis for the recommendations of the US Preventive Services Task Force, came to the conclusion that HPV testing using the Hybrid Capture 2 Test is significantly more sensitive than cytology only with respect to the detection of CIN 2+ (and of CIN 3+, less data available). Nevertheless the authors concluded that the available data was not yet complete, consistent, or relevant enough to permit a conclusive decision to be made about the role of HPV testing in primary screening.

In their interpretation of the data, the IQWiG 2011 and 2014 studies noted that the data appeared to show (Table 1) that the incidence of CIN 3+ and of invasive cervical cancer was lower with HPV screening and came to the conclusion that the level of evidence of Ia meant that HPV screening should be proposed to the German Federal Joint Committee as a general recommendation – although not as a strong recommendation [15, 16].

In their review and meta-analysis published in 2012 Murphy et al. recommended even more strongly that HPV testing should be introduced in the context of organized screening programs as the primary screening method for women over the age of 30 or 35 years. MaHTAS (Health Technology Assessment Section, Ministry of Health Malaysia) was also of the opinion that there was a high level of evidence showing HPV screening would be able to reduce the incidence and mortality rate of cervical cancer.

Overall, six of the 13 meta-analyses investigating this issue reported a clear benefit of using HPV test (potentially combined with cytology for co-testing) or recommended HPV test as the primary screening method (Arbyn 2015, Bouchard-Fortier, MaHTAS, MSAC, Murphy, and Pileggi). The remaining meta-analyses also reported benefits from HPV screening but were of the opinion that the data was not yet sufficient to support a general recommendation to use HPV testing to screen for cervical cancer. None of the analyses reported that cytology-based screenings offered more benefits compared to HPV screening.

Discussion

In Western countries all over the world prevention programs for cervical cancer are being revised. Currently, numerous countries (USA, Australia, Netherlands, Spain, Portugal, Sweden, Italy, Slovenia, etc.) are switching from previously established programs based on cytological screening to early detection of cervical pre-

cancer based on HPV testing in women above the age of 30–35 years. In the USA recent publications by Ronco et al. [23] and the final results of the ATHENA study [24] have led to primary HPV screening being recommended since the beginning of 2015 as an alternative to cytological screening due to its equal or higher efficacy compared to cytology [25]. This paradigm change is based on data obtained from various high-quality RCTs (see above). When formulating recommendations, policymakers in the healthcare system base their decisions on the results of systematic reviews which summarize RCTs. Depending on the method and analysis used in these reviews, the reviews come to sometimes quite different conclusions – as was shown in this systematic summary.

Meta-analyses can arrive at different conclusions because they interpret the existing evidence differently. The IQWiG has highlighted the “very different screening strategies” and the “limited similarities between studies” [15]. Kleijnen also considers the heterogeneity of studies to be moderate to high and the level of evidence for this to be very low [17], and Patanwala entirely dispensed with meta-analytic pooling of individual studies because of their heterogeneity [9]. In contrast, MaHTAS in Malaysia [14] and the publication by Arbyn et al. in 2015 [18] reported that the level of evidence was good or strong, using the same data. The primary studies included in the reviews also differed significantly. The greatest differences between the different meta-analyses were reflected in the quality as measured using the AMSTAR criteria. Five reviews received only three or fewer points (out of a maximum of eleven) (Table 2); however, this may also be due to the fact that the documentation was not always very transparent (no description of the methodology used).

In the end, around half of all investigated meta-analyses found significant indications that screening based on HPV testing was superior to cytology or recommended HPV-based screening. The remaining studies considered that the data was not yet sufficient to warrant a clear recommendation. The aim of our study was to identify and present the impact of methodological groundwork on decisions taken by healthcare policymakers.

The Lancet article by Ronco et al. published in 2013 [23] was not included in our study. Their publication reviewed and followed up the data of around 180 000 women aged between 20 and 64 years studied in four RCTs (Swedescreen, POBASCAM, ARTISTIC and NTCC) over an average of six-and-a-half years. A total of 107 invasive cervical cancers occurred in that period; the ratio of the incidence rates was 0.60 (95% CI: 0.40–0.89) in favor of the HPV group. According to these data, the protection against invasive cancer was 60–70% higher with HPV-based screening. With regard to the screening intervals, the cumulative cancer incidence was lower five-and-a-half years after a negative HPV test than at three-and-a-half years after an unremarkable cytology-based test, meaning that screening based on testing for HPV every five years in women aged over 30 years promises greater safety than cytology screening carried out every three years [23]. HPV screening carried out every three years was not compared with annual cytology-based screening, as the latter is not done in four European countries and is not recommended by the IARC/WHO. It will be interesting to see which decisions will be taken with regard to future programs for the prevention of cervical cancer in Germany. In Germany, the Joint Federal Committee (G-BA) is responsible for establishing the guidelines. The S3 guideline “Prevention of Cervical Cancer” will be finalized shortly and will be published later this year.

Conclusion

Different meta-analyses which examined the same or similar issues came to quite different conclusions because of the different approaches used for analysis. This is quite significant because the primary data on which the meta-analyses were based did not change. The data indicate that HPV-based screening is superior. The heterogeneity of the meta-analyses affects the decisions taken by policymakers in the healthcare system.

Conflict of Interest

M. Jentschke received lecture fees and partial reimbursement of travel and conference expenses from Abbott Molecular, Wiesbaden. P. Hillemanns received research grants from GSK and Abbott and lecture fees from SPMSD, Roche and Hologic.

References

- 1 Bundesministerium für Gesundheit. Ziel 2a – Weiterentwicklung der Gebärmutterhals-Krebsfrüherkennung. 2012. Online: <http://www.bmg.bund.de/p/praevention/nationaler-krebsplan/was-haben-wir-bisher-erreicht/ziel-2a-weiterentwicklung-der-gebaermutterhals-krebsfrueherkennung.html>; last access: 04.09.2014
- 2 Marquardt K, Buttner HH, Broschewitz U et al. Persistent carcinoma in cervical cancer screening: non-participation is the most significant cause. *Acta Cytol* 2011; 55: 433–437
- 3 Jentschke M, Soergel P, Hillemanns P. Importance of HPV genotyping for the screening, therapy and management of cervical neoplasias. *Geburtsh Frauenheilk* 2012; 72: 507–512
- 4 Shea BJ, Grimshaw JM, Wells GA et al. Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. *BMC Med Res Methodol* 2007; 7: 10
- 5 Bouchard-Fortier G, Hajifathalian K, McKnight MD et al. Co-testing for detection of high-grade cervical intraepithelial neoplasia and cancer compared with cytology alone: a meta-analysis of randomized controlled trials. *J Public Health (Oxf)* 2014; 36: 46–55
- 6 Luu HN, Dahlstrom KR, Mullen PD et al. Comparison of the accuracy of Hybrid Capture II and polymerase chain reaction in detecting clinically important cervical dysplasia: a systematic review and meta-analysis. *Cancer Med* 2013; 2: 367–390
- 7 Murphy J, Kennedy EB, Dunn S et al. HPV testing in primary cervical screening: a systematic review and meta-analysis. *J Obstet Gynaecol Can* 2012; 34: 443–452
- 8 Pan QJ, Hu SY, Guo HQ et al. Liquid-based cytology and human papillomavirus testing: a pooled analysis using the data from 13 population-based cervical cancer screening studies from China. *Gynecol Oncol* 2014; 133: 172–179
- 9 Patanwala IY, Bauer HM, Miyamoto J et al. A systematic review of randomized trials assessing human papillomavirus testing in cervical cancer screening. *Am J Obstet Gynecol* 2013; 208: 343–353
- 10 Peirson L, Fitzpatrick-Lewis D, Ciliska D et al. Screening for cervical cancer: a systematic review and meta-analysis. *Syst Rev* 2013; 2: 35
- 11 Pileggi C, Flotta D, Bianco A et al. Is HPV DNA testing specificity comparable to that of cytological testing in primary cervical cancer screening? Results of a meta-analysis of randomized controlled trials. *Int J Cancer* 2014; 135: 166–177
- 12 Vesco KK, Whitlock EP, Eder M et al. Screening for cervical Cancer: a systematic Evidence Review for the U.S. Preventive Services Task Force. Evidence Synthesis No. 86. AHRQ Publication No. 11-05156-EF-1. 2011/12. 2nd ed. Rockville, MD: Agency for Healthcare Research and Quality; 2011
- 13 Whitlock EP, Vesco KK, Eder M et al. Liquid-based cytology and human papillomavirus testing to screen for cervical cancer: a systematic review for the U.S. Preventive Services Task Force. *Ann Intern Med* 2011; 155: 687–697, W214–W215
- 14 Health Technology Assessment Section (MaHTAS). HPV DNA-BASED SCREENING TEST FOR CERVICAL CANCER. Ministry of Health Malaysia; 2011. Online: <http://www.moh.gov.my/attachments/6726.pdf>; last access: 29.07.2014
- 15 Institut für Qualität und Wirtschaftlichkeit im Gesundheitswesen (IQWiG). Nutzenbewertung eines HPV-Tests im Primärscreening des Zervixkarzinoms. 2011. Online: https://www.iqwig.de/download/S10-01_AB_HPV-Test_im_Primaerscreening_des_Zervixkarzinoms.pdf; last access: 29.07.2014
- 16 Institut für Qualität und Wirtschaftlichkeit im Gesundheitswesen (IQWiG). [S13-03] Nutzenbewertung eines HPV-Tests im Primärscreening des Zervixkarzinoms – Rapid Report. 2014. Online: https://www.iqwig.de/download/S13-03_Rapid-Report_HPV-Test-im-Primaerscreening-des-Zervixkarzinoms.pdf; last access: 29.07.2014
- 17 Birnie R, Wolff R, Hilmer D et al. Evidence Review for the S3 Guideline “Prevention of Cervical Cancer”. York, United Kingdom: Ltd. KSR ed.; 2014
- 18 Arbyn M, Haelens A, Desomer A et al. Cervical cancer screening program and Human Papillomavirus (HPV) testing, part II: update on HPV primary screening. KCE Belgian Healthcare Knowledge Centre; 2015. Online: https://kce.fgov.be/sites/default/files/page_documents/KCE_238Cs_HPV%20DNA%20Testing_Synthesis.pdf; last access: 19.01.2015
- 19 Arbyn M, Ronco G, Anttila A et al. Evidence regarding human papillomavirus testing in secondary prevention of cervical cancer. *Vaccine* 2012; 30 (Suppl. 5): F88–F99
- 20 Medical Services Advisory Committee (MSAC). Application No. 1276 – Renewal of the National Cervical Screening Program. Australian Government, Department of Health; 2014. Online: [http://www.msac.gov.au/internet/msac/publishing.nsf/Content/FD36D6990FFAA639CA25799200058940/\\$File/1276%20-%20Final%20MSAC%20PSD%20-%20NCS%20Renewal.pdf](http://www.msac.gov.au/internet/msac/publishing.nsf/Content/FD36D6990FFAA639CA25799200058940/$File/1276%20-%20Final%20MSAC%20PSD%20-%20NCS%20Renewal.pdf); last access: 20.08.2014
- 21 Sankaranarayanan R, Nene BM, Shastri SS et al. HPV screening for cervical cancer in rural India. *N Engl J Med* 2009; 360: 1385–1394
- 22 Mayrand MH, Duarte-Franco E, Rodrigues I et al. Human papillomavirus DNA versus Papanicolaou screening tests for cervical cancer. *N Engl J Med* 2007; 357: 1579–1588
- 23 Ronco G, Dillner J, Elfstrom KM et al. Efficacy of HPV-based screening for prevention of invasive cervical cancer: follow-up of four European randomised controlled trials. *Lancet* 2013; 383: 524–532
- 24 Wright TC, Stoler MH, Behrens CM et al. Primary cervical cancer screening with human papillomavirus: end of study results from the ATHENA study using HPV as the first-line screening test. *Gynecol Oncol* 2015; 136: 189–197
- 25 Huh WK, Ault KA, Chelmow D et al. Use of primary high-risk human papillomavirus testing for cervical cancer screening: interim clinical guidance. *Gynecol Oncol* 2015; 136: 178–182