Endoscopic abbreviations

Authors

Deborah E. Bowman, Hilary Hamilton-Gibbs, Michael B. Wallace, Peter D. Siersema

Bibliography

DOI http://dx.doi.org/ 10.1055/s-0042-114571 Published online: 30.8.2016 Endoscopy 2016; 48: 876–878 © Georg Thieme Verlag KG Stuttgart - New York ISSN 0013-726X

Corresponding author

Hilary Hamilton-Gibbs Managing Editor, Endoscopy Editorial Office Landwehr Str. 9 80336 Munich Germany Phone: +49-89-9077-936-00 Fax: +49-89-9077-936-20 hhg@hamilton-services.de

This article is published simultaneously in the journals *Endoscopy* and *Gastrointestinal Endoscopy*. Copyright 2016 © by Georg Thieme Verlag KG and © by the American Society for Gastrointestinal Endoscopy We all use abbreviations when we write papers for scientific journals. Abbreviations help us to limit the word count of a research paper, especially when, as is common nowadays, journals stipulate a maximum word count.

Furthermore, in daily clinical practice, some procedures or devices are often referred to by their abbreviation, not only in discussions between colleagues but also when we are talking to our patients. Examples include EMR and ESD for endoscopic mucosal resection and endoscopic submucosal dissection, respectively, both of which are well recognized and easily understood.

However, there are also abbreviations that are less common and that are used in some articles but are written out in full in others. Examples include ER for endoscopic resection and SC for screening colonoscopy. Moreover, some abbreviations can have more than one meaning; for example, IM can stand for intramuscular, intramucosal, or intestinal metaplasia. Other abbreviations could stand for words or phrases not directly related to our field, such as MI for myocardial infarction or mitral valve insufficiency.

For the sake of clarity, *Endoscopy* and *Gastrointestinal Endoscopy* have joined forces to compile a list of generally accepted abbreviations to be used in our journals. As you will see, we also include some terms that we feel should always be written in full. Our expectation is that this compilation will not only assist our authors when they are writing their papers, but will also help our readers avoid getting mired in unfamiliar abbreviations.

Of course, the abbreviations in this list are based on words and phrases that have been used in both journals in the past few years. We realize that procedures are evolving and new innovative products will be introduced in our endoscopy suites. We therefore plan to update the abbreviations regularly. For your reference, this list and future updated versions can be downloaded from the websites of both journals.

We hope this list is helpful to those writing research papers for *Endoscopy* and *Gastrointestinal Endoscopy*.

Deborah E. Bowman, MFA, ELS, Senior Managing Editor of Clinical Publications, *Gastrointestinal Endoscopy*

Hilary Hamilton-Gibbs, Managing Editor, *Endoscopy*

Michael B. Wallace, MD, MPH, Editor-in-Chief, *Gastrointestinal Endoscopy*

Peter D. Siersema, MD, PhD, Editor-in-Chief, *Endoscopy*

| Word or Phrase | Abbreviation | Word or Phrase | Abbreviation |
|--|-------------------|---|---------------------|
| acute biliary pancreatitis | ABP | erythrocyte sedimentation rate | ESR |
| acute necrotic collection | ANC | esophageal adenocarcinoma | EAC |
| acute pancreatitis | Write in full | esophageal introitus | Replace by upper |
| acute peripancreatic fluid collection | APFC | | esophageal sphinc |
| adenoma detection rate | ADR | | ter (UES) |
| analysis of covariance | ANCOVA | esophageal squamous cell carcinoma | ESCC |
| analysis of variance | ANOVA | esophagogastric junction | EGJ |
| anteroposterior | Write in full | esophagogastroduodenoscopy | EGD |
| area under the receiver-operating characteristic | AUC | EUS-guided biliary drainage | EUS-BD |
| curve | | EUS-guided gallbladder drainage | EUS-GBD |
| argon plasma coagulation | APC | extracorporeal shock wave lithotripsy | ESWL |
| artificial neural network | ANN | familial adenomatous polyposis | FAP |
| autofluorescence imaging | AFI | fecal immunochemical test | FIT |
| autoimmune pancreatitis | AIP | fecal occult blood test | FOBT |
| Barrett's esophagus | BE | fine-needle aspiration | FNA |
| benign biliary stricture | BBS | flexible sigmoidoscopy | Write in full |
| biosimilar | BSim | flexible spectral imaging color enhancement | FICE |
| body mass index | BMI | fluorescence in situ hybridization | FISH |
| 5 | | fluorodeoxyglucose | |
| Boston Bowel Preparation Scale | BBPS | ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,, | FDG |
| capsule endoscopy | Use video capsule | fully covered self-expandable metal stent | FCSEMS |
| | endoscopy (VCE) | gastric intestinal metaplasia | GIM |
| carcinoembryonic antigen | CEA | gastric outlet obstruction | GOO |
| celiac plexus neurolysis | CPN | Gastric Outlet Obstruction Scoring System | GOOSS |
| charge-coupled device | CCD | gastroesophageal reflux disease | GERD |
| cholangiocarcinoma | CCA | gastrointestinal | GI |
| choledochoduodenostomy | CDS | GI stromal tumor | GIST |
| chronic pancreatitis | Write in full | hazard ratio | HR |
| chronic radiation proctopathy | CRP | Helicobacter pylori | HP |
| Clinical Outcomes Research Initiative | CORI | hematoxylin and eosin | H&E |
| colorectal cancer | CRC | hepatic abscess | Write in full |
| common bile duct | CBD | hepaticogastrostomy | HPG |
| computed tomographic colonography | CTC | high definition | Use high-definition |
| computed tomography | CT | ingli definition | resolution |
| | CAD | high-definition resolution | HDR |
| computer-aided design | | high-grade dysplasia | HGD |
| confidence interval | CI | | |
| confidence limit | CL | high-grade intra-epithelial neoplasia | HGIN |
| confocal laser endomicroscopy with variants: | CLE; also pCLE, | high-resolution manometry | HRM |
| probe-based CLE, needle-based CLE, endoscope- | nCLE, eCLE | hyperplastic polyp | Write in full |
| based CLE | | image-enhanced endoscopy | IEE |
| contrast enhancement | Write in full | immunoglobulin | lg |
| C-reactive protein | CRP | infected pancreatic necrosis | IPN |
| Crohn's disease | CD | inflammatory bowel disease | IBD |
| cumulative sum | CUSUM | intention-to-treat | ITT |
| diagnostic odds ratio | DOR | international normalized ratio | INR |
| direct endoscopic necrosectomy | DEN | interquartile range | IQR |
| dose area product | DAP | intestinal metaplasia | IM |
| double-balloon enteroscopy | DBE | intraductal papillary mucinous neoplasm | IPMN |
| early gastric cancer | EGC | intraductal ultrasonography | IDUS |
| electrohydraulic lithotripsy | EHL | intraductal diclasofiography | |
| electronic chromoendoscopy | | | Write in full |
| | ECE | laterally spreading tumor | LST |
| endogastric tube | EGT | likelihood ratio | LR |
| endoscopic full-thickness resection | EFTR | lower esophageal sphincter | LES |
| endoscopic mucosal resection | EMR | lower GI bleeding | LGIB |
| endoscopic resection | Write in full | low-grade dysplasia | LGD |
| endoscopic retrograde cholangiography | ERC | low-grade intraepithial neoplasia | LGIN |
| endoscopic retrograde cholangiopancreato- | ERCP | lumen-apposing metal stent | LAMS |
| graphy | | lymph node | LN |
| endoscopic retrograde pancreatography | ERP | lymph node metastasis | LNM |
| endoscopic sphincterotomy | Write in full | magnetic resonance cholagiopancreatography | MRCP |
| endoscopic submucosal dissection | ESD | magnetic resonance enterography | MRE |
| endoscopic ultrasound or endoscopic ultra- | EUS | magnetic resonance imaging | MRI |
| | | magnetic resonance imaging | MRI |
| sonography | | | |
| | EUS-FNA | mean adenomas per proceduro | MAP |
| sonography endoscopic ultrasound-guided fine-needle aspiration | EUS-FNA | mean adenomas per procedure metabolic syndrome | MAP MetS |

| Word or Phrase | Abbreviation | Word or Phrase | Abbreviation |
|--|---------------|---|------------------|
| mucosa-associated lymphoid tissue | MALT | sessile serrated adenoma | SSA |
| mucosal vascular pattern | MVP | sessile serrated polyp | SSP |
| narrow-band imaging | NBI | single-balloon enteroscopy | SBE |
| nasogastric | Write in full | small bowel | Write in full |
| nasogastric tube | NGT | small-bowel capsule endoscopy | SBCE |
| natural orifice transluminal endopscopic surgery | NOTES | spastic esophageal disorder | SED |
| necrotizing pancreatitis | Write in full | sphincter of Oddi dysfunction | SOD |
| needle-based confocal laser endomicroscopy | nCLE | sphincter of Oddi manometry | SOM |
| negative predictive value | NPV | squamous cell carcinoma | SCC |
| neuroendocrine tumor | NET | standard deviation | SD |
| nonsteroidal anti-inflammatory drug | NSAID | standard error of the mean | SEM |
| normal saline solution | NSS | submucosal | Write in full |
| obscure GI bleeding | OGIB | submucosal fluid cushion | SFC |
| odds ratio | OR | submucosal gland | SMG |
| open gastrojejunostomy | OGJ | systemic inflammatory response syndrome (SIRS) | SIRS |
| open-access endoscopy | OAE | Third Eye Retroscope | TER |
| oral sodium sulfate | OSS | total radiation dose | TRD |
| orthotopic liver transplantation | OLT | tracheoesophageal fistula | TEF |
| over-the-scope clip | OTSC | traditional serrated adenoma | TSA |
| pancreatic duct | PD | transhepatic portosystemic shunt | TIPSS |
| pancreatic fluid collection | PFC | ulcerative colitis | UC |
| pancreatic stent placement | PSP | uncovered self-expandable metal stent | USEMS |
| partially covered self-expandable metal stent | PCSEMS | underwater EMR | UEMR |
| peptic ulcer disease | PUD | upper esophageal sphincter | UES |
| percutaneous drainage | PCD | upper GI bleeding | UGIB |
| percutaneous endoscopic gastrojejunostomy | PEG | video capsule endoscopy | VCE |
| percutaneous endoscopic jejunostomy | PEJ | visual analog scale | VAS |
| percutaneous endoscopic necrosectomy | PEN | volatile organic compound | VOC |
| peroral endoscopic myotomy | POEM | walled-off necrosis | WON |
| peroral pancreatoscopy | POPS | walled-off pancreatic necrosis | Use WON |
| photodynamic therapy | PDT | water exchange | Write in full |
| polyethylene glycol | PEG | water immersion | Write in full |
| polyglycolic acid | PGA | white-light colonoscopy | WLC |
| polymerase chain reaction | PCR | white-light endoscopy | WLE |
| polyp detection rate | PDR | white-light imaging | WLI |
| positive predictive value | PPV | wire-guided cannulation | WGC |
| positron emission tomography | PET | Brand Name | Abbreviation |
| post-ERCP pancreatitis | PEP | Barrx-HALO radiofrequency ablation method | RFA-HALO |
| Preservation and Incorporation of Valuable | PIVI | DualKnife | Write in full |
| Endoscopic Innovations | | Hemoclip | Replace with cli |
| primary sclerosing cholangitis | PSC | Hemospray | Replace with he |
| probe-based confocal laser endomicroscopy | pCLE | nemospidy | mostatic spray |
| proton pump inhibitor | PPI | HybridKnife | Write in full |
| radiofrequency ablation | RFA | i-SCAN digital contrast | I-SCAN |
| randomized controlled trial | RCT | SpyGlass, SpyProbe, SpyBite cholangioscopy sys- | Replace with int |
| rapid on-site evaluation | ROSE | tem | ductal cholangi |
| receiver operating characteristic | ROC | | copy (IDC) |
| rectal telangiectasia density | RTD | WavSTAT | Replace with las |
| regions of interest | ROI | | induced fluores |
| relative risk or risk ratio | RR | | cence spectrosc |
| Roux-en-Y gastric bypass | RYGB | | py (LIFS) |
| screening colonoscopy | Write in full | | |
| | | | |
| self-expandable metal stent | SEMS SAE | | |