

Syphilis infection throughout the whole gastrointestinal tract

A 30-year-old Japanese woman presented to our hospital with epigastric pain. A physical examination revealed epigastric tenderness and red eruptions on her face. Laboratory evaluation showed normal results for leukocyte count and serum protein level, but a slightly elevated C-reactive protein (CRP) level of 1.10 mg/dL. The findings on serologic tests for syphilis, including rapid plasma regain and *Treponema pallidum* antibody hemagglutination test, were positive.

Esophagogastroduodenoscopy (EGD) revealed hypertrophy of the rugae and multiple irregular ulcers in the stomach (▶ Fig. 1). A *Helicobacter pylori* culture test was negative. Colonoscopy revealed reddish mucosa in the terminal ileum and slightly elevated lesions with decreased normal vascular networks in the descending colon (▶ Fig. 2). Biopsy specimens were obtained from the gastric ul-

cers, the red ileal mucosa, and the elevated lesions in the descending colon. Pathological analysis of these revealed regenerative epithelium and marked lymphoplasmacytic infiltration with no evidence of malignancy. Warthin–Starry staining revealed infiltration by spirochetes in the epithelium and lamina propria of the ileal and colonic tissues. No spirochetes were detected in the gastric tissues, although a polymerase chain reaction did detect the presence of some genes from *T. pallidum* with 100% similarity.

The patient was treated with penicillin (1500 mg/day) for 7 weeks, following which both her epigastric pain and the endoscopic appearances were improved. This is the first reported case of syphilis infection in which lesions were observed throughout the whole gastrointestinal tract. It has been reported that the rate of gastric involvement is only 1% [1] and

syphilis infections in the colon have been detected in only a few cases [2–5]. When gastroscopic findings such as hypertrophy of the rugae and multiple irregular ulcers are observed, gastric syphilis should be considered. In addition, other sections of the gastrointestinal tract, including the small and large intestines, should be examined.

Endoscopy_UCTN_Code_CCL_1AB_2AG

Competing interests: None

Masami Ijiri¹, Mikihiro Fujiya¹, Nobuhiro Ueno¹, Shin Kashima¹, Tomohisa Watari², Satoshi Fujii³, Toshikatu Okumura¹

¹ Division of Gastroenterology and Hematology/Oncology, Department of Medicine, Asahikawa Medical University, Asahikawa, Hokkaido, Japan

² Department of Medical Laboratory and Blood Center, Asahikawa Medical University Hospital, Asahikawa, Hokkaido, Japan

³ Department of Laboratory Medicine, Asahikawa Medical University, Asahikawa, Hokkaido, Japan



Fig. 1 Esophagogastroduodenoscopy images showing hypertrophy of the rugae and multiple irregular ulcers in the stomach.




Fig. 2 Colonoscopy images showing: **a** reddish mucosa in the terminal ileum; **b, c** slightly elevated lesions with decreased normal vascular networks in the descending colon.

References

- 1 *Atten MJ, Attar BM, Teopengco E* et al. Gastric syphilis: a disease with multiple manifestations. *Am J Gastroenterol* 1994; 89: 2227–2229
- 2 *Song SH, Jang I, Kim BS* et al. A case of primary syphilis in the rectum. *J Korean Med Sci* 2005; 20: 886–887
- 3 *Bassi O, Cosa G, Colavolpe A* et al. Primary syphilis of the rectum – endoscopic and clinical features. Report of a case. *Dis Colon Rectum* 1991; 34: 1024–1026
- 4 *Febbraro I, Manetti G, Balestrieri P* et al. Rectal cancer or rectal chancre? Beware of primary syphilis. *Dig Liver Dis* 2008; 40: 579–581
- 5 *Israel GB, Leonor GR*. Primary syphilitic chancre in the rectum. *Endoscopy* 2014; 46: E533

Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-117715>
Endoscopy 2016; 48: E338–E339
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Mikihiro Fujiya, MD
Division of Gastroenterology
and Hematology/Oncology
Department of Medicine
Asahikawa Medical University
2-1-1-1, Midorigaoka-higashi
Asahikawa
Hokkaido 078-8510
Japan
Fax: +81-166-682469
fjym@asahikawa-med.ac.jp