


## Intra-arterial migration of a fractured endoscopic needle

A 53-year-old man with chronic pancreatitis presented with abdominal pain and sepsis. Imaging revealed a liver abscess secondary to distal biliary obstruction. Following drainage of his abscess, the patient underwent endoscopic retrograde cholangiopancreatography (ERCP) with placement of a fully covered metal biliary stent across a 2-cm suspicious shouldered and irregular low common bile duct stricture. A pancreatic head malignancy was suspected at ERCP and on computed tomography (CT) imaging. The regional specialist hepatobiliary multidisciplinary team (MDT) recommended endoscopic ultrasound with fine needle aspiration (EUS-FNA).

EUS-FNA was difficult because of the changes of severe chronic calcific pancreatitis, duodenal stenosis, increased pancreatic head vascularity, and metal stent artefact. Five needle passes were made with a 22-gauge needle (Boston Scientific, Marlborough, Massachusetts, USA) using standard technique. Cytology was consistent with pancreatitis, with no evidence of malignancy.

A routine chest radiograph 6 months later revealed a new linear density in the heart (► **Fig. 1**). The interim abdominal CT imaging was re-reviewed (► **Fig. 2**). Although not recognized at the time, owing to the highly calcified pancreas, it became clear that a fractured EUS-needle tip had migrated from the duodenal wall into the epigastrium (► **Fig. 3**), then through the diaphragm and into the left ventricle. On a subsequent chest radiograph, the needle had disappeared and a further CT scan revealed that it had migrated to the aortic bifurcation (► **Fig. 4a**).

The needle was retrieved endovascularly via bilateral common femoral artery access. It was first snared from above with a protective occlusion balloon placed below in the left iliac artery (► **Fig. 4b**). The balloon was deflated, the needle was snared from below and was then removed through the left groin



► **Fig. 1** Chest radiograph showing a needle within the left ventricle (linear density within the red ellipse).

sheath (► **Fig. 4c**; ► **Video 1**). The patient made an uneventful recovery after the procedure.

Endoscopic needle fracture has been previously described in the upper gastrointestinal tract [1,2] and in a bronchoscopy setting [3]. Fractured metal sharps such as orthopedic fixation wires have been known to migrate into the arterial circulation, including into the heart [4].


This is the first known case of an endoscopic needle migrating intra-arterially.

Endoscopy\_UCTN\_Code\_CPL\_1AJ\_2AZ


### Competing interests

None


### VIDEO 1



► **Video 1:** Endovascular removal of an intra-arterial fractured needle. The needle was snared from above with a protective occlusion balloon below in the left iliac artery. The balloon was deflated, then the needle was snared from below and removed through the groin sheath.



► **Fig. 2** Axial computed tomography (CT) scan showing the needle (red arrow) that was not identified originally among the pancreatic calcifications.



► **Fig. 3** Obliquely reformatted computed tomography (CT) images showing: **a** the needle extending through the duodenal wall; **b** the needle having moved to the epigastrium.

## The Authors

**Edward Lake<sup>1</sup>, Joanne Puleston<sup>2</sup>,  
Finn Farquharson<sup>1</sup>**

- 1 Department of Radiology, Central Manchester University Hospitals NHS Foundation Trust, Manchester, UK
- 2 Department of Gastroenterology, Central Manchester University Hospitals NHS Foundation Trust, Manchester, UK

## Corresponding author

**Edward Lake, MD**

Department of Radiology, Manchester  
Royal Infirmary, Grafton Street, Manchester,  
M13 9WL, UK  
edwardlake@yahoo.com



► **Fig. 4** Further imaging shortly before and during removal of the needle. **a** A volume-rendered computed tomography (CT) angiogram showing the needle at the aortic bifurcation. **b** The needle was snared from above with balloon occlusion below. **c** The needle was snared from below and was removed through a sheath.

## References

- [1] Rimbaş M, Attili F, Andrade Zurita S et al. Fractured needle during endoscopic ultrasound-guided fine-needle aspiration of a pancreatic head mass. *Endoscopy* 2015; 47 (Suppl. 01): E432
- [2] DeWitt J, Sherman S, Lillemoe KD. Fracture of an EUS-guided FNA needle during an attempted rendezvous for an inaccessible pancreatic duct. *Gastrointest Endosc* 2011; 73: 171 – 173
- [3] Özgül MA, Çetinkaya E, Tutar N et al. An unusual complication of endobronchial ultrasound-guided transbronchial needle aspiration (EBUS-TBNA): the needle breakage. *Ann Thorac Cardiovasc Surg* 2014; 20 Suppl: 567 – 569
- [4] Leonardi F, Rivera F. Intravascular migration of a fractured cerclage wire into the left heart. *Orthopedics* 2014; 37: e932 – e935

## Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-123704>  
*Endoscopy* 2017; 49: E70–E72  
 © Georg Thieme Verlag KG  
 Stuttgart · New York  
 ISSN 0013-726X