



# Comparison of the Causes of Death Identified Using Automated Verbal Autopsy and Complete Autopsy among Brought-in-Dead Cases at a Tertiary Hospital in Sub-Saharan Africa

Yuta Yokobori<sup>1,2</sup> Jun Matsuura<sup>1</sup> Yasuo Sugiura<sup>1</sup> Charles Mutemba<sup>3,4</sup> Peter Julius<sup>3,5</sup>  
Cordelia Himwaze<sup>3,5</sup> Martin Nyahoda<sup>6</sup> Chomba Mwangi<sup>7</sup> Lloyd Kazhumbula<sup>3</sup> Motoyuki Yuasa<sup>2</sup>  
Brian Munkombwe<sup>8</sup> Luchenga Mucheleng'anga<sup>9</sup>

<sup>1</sup>National Center for Global Health and Medicine, Shinjuku-ku, Japan

<sup>2</sup>Department of Public Health, Graduate School of Medicine, Juntendo University, Tokyo, Japan

<sup>3</sup>Ministry of Health, Lusaka, Zambia

<sup>4</sup>Adult Hospital, University Teaching Hospital, Lusaka, Zambia

<sup>5</sup>Department of Pathology and Microbiology, School of Medicine, The University of Zambia, Lusaka, Zambia

<sup>6</sup>Department of National Registration of Home Passport & Citizenship, Ministry Affairs, Lusaka, Zambia

<sup>7</sup>Bloomberg Data for Health Initiative, Lusaka, Zambia

<sup>8</sup>National Center for Health Statistics, Center for Disease Control and Prevention, Atlanta, United States

<sup>9</sup>Office of the State Forensic Pathologist, Ministry of Home Affairs and Internal Security, Lusaka, Zambia

**Address for correspondence** Yuta Yokobori, MD, MPH, MSc, 1-21-1, Toyama, Shinjuku-ku, Tokyo, Japan (e-mail: y-yokobori@it.ncgm.go.jp).

Appl Clin Inform 2022;13:583–591.

## Abstract

**Background** Over one-third of deaths recorded at health facilities in Zambia are brought in dead (BID) and the causes of death (CODs) are not fully analyzed. The use of automated verbal autopsy (VA) has reportedly determined the CODs of more BID cases than the death notification form issued by the hospital. However, the validity of automated VA is yet to be fully investigated.

**Objectives** To compare the CODs identified by automated VA with those by complete autopsy to examine the validity of a VA tool.

**Methods** The study site was the tertiary hospital in the capital city of Zambia. From September 2019 to January 2020, all BID cases aged 13 years and older brought to the hospital during the daytime on weekdays were enrolled in this study. External COD cases were excluded. The deceased's relatives were interviewed using the 2016 World Health Organization VA questionnaire. The data were analyzed using InterVA, an automated VA tool, to determine the CODs, which were compared with the results of complete autopsies.

**Results** A total of 63 cases were included. The CODs of 50 BID cases were determined by both InterVA and complete autopsies. The positive predictive value of InterVA was

## Keywords

- ▶ CRVS
- ▶ Africa
- ▶ ICD
- ▶ automated VA
- ▶ verbal autopsy

received  
August 22, 2021  
accepted  
March 24, 2022

DOI <https://doi.org/10.1055/s-0042-1749118>.  
ISSN 1869-0327.

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

22%. InterVA determined the CODs correctly in 100% cases of maternal CODs, 27.5% cases of noncommunicable disease CODs, and 5.3% cases of communicable disease CODs. Using the three broader disease groups, 56.0% cases were classified in the same groups by both methods.

**Conclusion** While the positive predictive value was low, more than half of the cases were categorized into the same broader categories. However, there are several limitations in this study, including small sample size. More research is required to investigate the factors leading to discrepancies between the CODs determined by both methods to optimize the use of automated VA in Zambia.

## Background and Significance

Accurate and timely cause of death (COD) data are critical for guiding health policies and effectively mobilizing resources based on priority health issues.<sup>1</sup> Ideally, all deaths should be registered, and the CODs should be obtained from accurate death certificates.<sup>2</sup> However, a United Nations Statistics Division report demonstrated that only 55% of countries and regions worldwide achieve death registration rates  $\geq 90\%$ .<sup>3</sup> Approximately 60% of African countries achieved death registration rates  $< 50\%$ .<sup>3</sup> Therefore, improving civil registration and vital statistics (CRVS), including death registration, has become a key topic in global public health. Goal 16 of the Sustainable Development Goals<sup>4,5</sup> includes a target related to CRVS to promote human security. To address these issues and improve coverage of the CRVS, the health sector could play a key role by acting as an entry point, collecting accurate vital data, and utilizing information from CRVS.<sup>6</sup>

The Republic of Zambia is facing substantial challenges in collecting accurate death information. According to the Central Statistics Office, the 2016 death registration rate was 20%.<sup>7</sup> The low death registration rate may be attributed to the large proportion of deaths that occur outside of health facilities. According to the sample vital registration with the verbal autopsy (VA), approximately 50% of deaths occur at home.<sup>8</sup> Further, the information system of the Zambian Ministry of Health in 2016 revealed that more than one-third of deaths in health facilities occurred before arrival at a facility; these deaths are termed brought-in-dead (BID). The CODs of BID cases are not thoroughly analyzed.

VA is a method of gathering information about deceased individuals' symptoms and circumstances to determine their COD. Health information and a description of events prior to death are acquired from conversations or interviews with a person or persons familiar with the deceased.<sup>9</sup> VA is considered a realistic alternative to death certificates to classify the CODs of BID cases.<sup>10,11</sup> The World Health Organization (WHO) has recommended using VA to accumulate data regarding death statistics and to capture CODs to identify trends.<sup>12</sup> While many countries have incorporated VA into the public health information system,<sup>13–16</sup> physician-based VA for all BID cases may not be feasible due to workload and costs and the automation of VA reporting could be useful to improve data collection regarding mortality.<sup>17</sup> Therefore,

automated VA tools have been developed to enable trained personnel other than health care workers to reliably determine the COD of BID cases using a computer program that can assign the most probable COD using background information obtained during standardized interviews.

Based on literature review of previous publications, several automated VA programs such as InterVA, Naïve Bayes Classifier, and InSilicoVA<sup>18–22</sup> have been used to analyze COD in previous studies.<sup>23–27</sup> According to WHO, these automated VA programs have strengths and weaknesses, depending on the setting and the target cause. Currently, there are no recommendations for any particular algorithm. We previously evaluated the automated VA program Tariff Method 2.0 to identify the COD of BID cases from the main health facility in Zambia.<sup>28</sup> We compared COD data obtained using automated VA with the COD data on the death notification form based on insufficient background information and observed that the Tariff Method could determine the COD among more BID cases than the death notification form. Regarding the accuracy of COD data, a previous report indicated that there is a 50.0% chance that the Tariff Method automated VA can correctly identify the COD in adult BID cases.<sup>29</sup> However, any of the automated VA programs' validity has not yet been thoroughly investigated, as it was not compared with the results of complete autopsies.<sup>22,28,30–35</sup> The applicability of automated VA in Zambia to determine the COD of BID cases must be determined by comparing the VA results with those of actual autopsies. Therefore, this study investigates an automated VA tool's validity by comparing the results obtained using an automated VA tool with those obtained using a complete autopsy in BID cases in Zambia.

## Objectives

To compare the COD identified by an automated VA with those by complete autopsy to examine the validity of a VA tool.

## Methods

### Study Design

This cross-sectional study aimed to compare the COD of BID cases identified by an automated VA and complete autopsies at the University Teaching Hospital, a tertiary hospital in Lusaka, the Republic of Zambia.

Lusaka has a population of approximately 2.5 million.<sup>36</sup> As this was the only tertiary hospital in Lusaka with a forensic pathologist when this study began, most BID cases in the city were referred to this hospital. Due to limited resources and a high number of deaths, the cases that died from external causes, such as injuries, suicide, and homicide, were excluded; only BID cases of individuals >13 years of age who were brought to the facility during the daytime hours from Monday to Friday from September 2019 to January 2020 were included. BID was defined as a patient who had died before they arrived at the hospital.

## Data Collection

### Data Collection of Automated VA

The staff who were employed by the Ministry of Home Affairs collected VA data to determine the CODs of the BID cases. First, they asked triage questions to exclude cases who died from external causes and informed consent was obtained from the relatives of each BID case with internal CODs. The staff conducted interviews regarding the death's background using a standardized questionnaire of the 2016 WHO VA instrument (in English).<sup>37</sup> During the interview, the staff used the Android application Open Data Kit (ODK) Collect to convert the collected data to an electronic format.<sup>38</sup> The data stored in ODK Collect were transferred to the Civil Registration Office in Zambia, where they were converted to comma-separated value files. These files were used with InterVA-5 software to identify the CODs as 1 of 53 categories for adults. The Civil Registration Office shared the COD results from InterVA-5 software with our team of researchers. InterVA-5 was chosen as the automated VA tool because the Civil Registration Office already used InterVA-5 to determine CODs among BID cases.

### InterVA-5

The InterVA model<sup>21,39,40</sup> is a Bayesian probability theory-based algorithm developed in 2003 and has since been revised. Bayes' theorem defines the conditional probability of a cause in the presence of a particular indicator.<sup>34,39,41</sup> In building the model, prior probabilities were assigned by expert panels using a semiquantitative scale for each indicator and COD. The likelihood of each COD is calculated by applying the aforementioned theorem, considering each pertinent indicator reported in the VA interview.<sup>42</sup> InterVA can also determine the COD with the highest propensity in a population, resulting in population-level data regarding COD. For each case, InterVA reports single-value point estimates for the three CODs with the highest propensities if they fall above a set threshold. If any COD does not meet the threshold, the death is ruled indeterminate. InterVA-5 has been developed to accommodate the 2016 WHO VA standards and is currently under testing.<sup>22</sup> The product information of InterVA-5 (version 5.1) is available at <http://www.byass-uk/interva/>.

### Data Collection of Autopsies

A forensic pathologist trained in both Zambia and Canada and a surgical pathologist conducted all autopsies, including

the pathological investigations, and recorded the CODs in the study registry. Basic demographic data including sex, age, and research identity were also included. Each COD was categorized into the most appropriate category on the COD list in the 2016 WHO VA instrument by mutual agreement among the three physicians so that the results of CODs by automated VA and complete autopsy could be directly compared.

### Data Verification

We verified that the cases with consent for participation were located in the official civil registry using the patient names, ages, and death dates. If we could not confirm the data's consistency, the case was excluded from the data analysis.

### Statistical Analyses

The 10 most common CODs obtained using the automated VA and complete autopsy were listed and compared. The positive predictive value of automated VA was calculated using four broader disease categories: communicable disease, non-communicable disease (NCD), injury, and maternal CODs. These results were also described by a visual graph. STATA version 16 statistical software (StataCorp., College Station, Texas, United States) and Microsoft Excel 2013 (Microsoft, Redmond, Washington, United States) were used for the data analyses. The concordance of CODs between the two methods was estimated using Cohen's kappa coefficient and was considered excellent if the coefficient was from 1.0 to 0.8; good if the coefficient was from 0.6 to 0.8; moderate if the coefficient was from 0.4 to 0.6; weak if the coefficient was from 0.4 to 0.6, minimal if the coefficient was from 0.2 to 0.4, and none if the coefficient was less than 0.2.<sup>43</sup>


## Results

### Demographic Data

→Fig. 1 shows the flowchart of BID cases enrolled in this study. A total of 152 cases were eligible for this study; 47 cases were excluded as injury cases, the relatives of 33 BID cases chose not to participate, and 9 cases were not verified with the Civil Registration Office. Therefore, the final analysis included 63 BID cases. The demographic data for the BID cases are presented in →Table 1. The median age of the BID cases was 41.0 years (interquartile range: 32.0–52.0 years) (→Fig. 2) and 76.2% were male.

### CODs by Automated VA and Autopsy

Of the 63 BID cases included in this study, the CODs were determined by both automated VA and autopsy in 50 cases (79.4%) (→Table 2). The 10 most common CODs determined by each method are shown in →Table 3. The most common CODs determined by autopsy were different from those determined by automated VA. →Supplementary Table S1 (available in the online version) shows the full lists of the CODs determined by each method. →Table 4 shows the number of CODs in each of the four broad categories used in this study. The positive predictive value of InterVA-5 in the



**Fig. 1** Flowchart of participant enrollment.

**Table 1** Characteristics of eligible participants

Total number	63
Male (%)	76.2%
Age (y) <sup>a</sup>	41 (32.0–52.0)


<sup>a</sup>Age is presented as median and interquartile range.

total population was 20.0% (10/50) and the kappa coefficient was less than 0.2. While the positive predictive value of InterVA-5 for maternal CODs was high (100%; 2/2), it was low for NCD CODs (25.0%; 7/28) and communicable disease CODs

(5.0%; 1/20). The kappa coefficient of each broader category was less than 0.2. The proportions of CODs between InterVA-5 and autopsy in four disease categories are shown in **Table 5** and they indicate that 56.0% of all cases were diagnosed in the same disease categories by both methods. **Fig. 3** also describes these results in a bar graph.

## Discussion

This study compares the CODs determined by complete autopsy with those determined by automated VA in 63 BID cases. As the forensic pathologist who conducted the



**Fig. 2** Age distribution of the participants.

**Table 2** The number of cases whose CODs could not be determined

	Number	Percentage
Undetermined by both VA and autopsy	4	6.3%
Undetermined by VA	6	9.5%
Undetermined by autopsy	3	4.8%
Determined by both VA and autopsy	50	79.4%

Abbreviations: COD, cause of death; VA, verbal autopsy.

autopsies is an expert who received professional training in Zambia and Canada with several years of experience at the target hospital, and another surgical pathologist worked with him for quality control purposes, the autopsy results can be assumed to be reliable sources of CODs. According to the results, the positive predictive values of InterVA-5 on CODs determined by complete autopsy were as low as 22% for the total population.

There are several factors related to weaknesses of automated VA that may have led to the low positive predictive

value observed in our study. Regarding the methodology of VA, there may be challenges in differentiating some diseases with similar symptoms such as acute cardiac disease and stroke, acute cardiac disease and pneumonia, and epilepsy and convulsion from poisoning. To accurately identify the COD of these cases, detailed information of clinical presentation before the deaths needs to be collected. However, in the circumstances where VA is conducted, it may be hard to obtain the accurate data because some important information is limited to the deceased and their closest relations. Therefore, the selection of interviewees may have affected the unfavorable results of this study. Since the surveyors interviewed the persons who brought the deceased to the health facility, the interviewees were not necessarily those who were the most informed regarding the deceased’s health course.

In addition, the categorization of CODs in the 2016 WHO VA instrument may have contributed to the low positive predictive value in this study, as some CODs determined by autopsy could not be represented well in the categories of the instrument. For example, four BID cases identified with disseminated tuberculosis (TB) could not be categorized accurately on the 2016 WHO VA instrument, as it only includes pulmonary TB.<sup>37</sup> As pulmonary TB and

**Table 3** Top CODs identified by InterVA and autopsy corresponding to VA COD list

Automated VA tool (N = 50)				Autopsy (N = 50)			
Rank	Cause of death	No	%	Rank	Cause of death	No	%
1	Acute cardiac disease	18	36.0%	1	Other CVD	10	20.0%
2	HIV/AIDS	9	18.0%	2	Pneumonia	6	12.0%
3	Epilepsy	5	10.0%	3	Acute cardiac disease	6	12.0%
4	Pneumonia	4	8.0%	4	Poisoning	5	10.0%
5	Pulmonary TB	4	8.0%	5	Other infectious disease	5	10.0%
6	Stroke	3	6.0%	6	Unspecified infectious disease	5	10.0%
7	Ectopic pregnancy	2	4.0%	7	Stroke	4	8.0%
8	Diarrheal disease	1	2.0%	8	Meningitis	3	6.0%
8	Other CVD	1	2.0%	9	Other NCD	3	6.0%
8	Meningitis	1	2.0%	10	Ectopic pregnancy	2	4.0%
8	Other infectious disease	1	2.0%	10	Epilepsy	2	4.0%
8	Other NCD	1	2.0%				

Abbreviations: COD: cause of death; CVDs: cardiovascular diseases; NCDs: noncommunicable diseases; TB: tuberculosis; VA: verbal autopsy.

**Table 4** Proportion of cases with CODs as determined by InterVA and autopsy

Disease category	No. of automated VA	No. of matched cases with autopsy	PPR (%)	Kappa coefficient
Communicable diseases	20	1	5.0%	<0.2
NCD	28	7	25.0%	<0.2
Maternal reasons	2	2	100%	NA <sup>a</sup>
Total	50	11	20.0%	<0.2


Abbreviations: NCD, noncommunicable disease; PCOD, cause of death.

<sup>a</sup>NA: Not analyzed due to too few samples

**Table 5** Proportion of cases with disease categories as determined by InterVA and autopsy

Disease category	No. of automated VA	No. of matched cases with autopsy	%
Communicable diseases	20	9	45.0%
NCD	28	17	60.7%
Maternal reasons	2	2	100%
Total	50	28	56.0%

Abbreviations: COD, cause of deaths; NCD, noncommunicable disease.

**Fig. 3** Matched proportion of CODs between InterVA and complete autopsy by InterVA and Broad categories. CODs, causes of death.

disseminated TB correspond to different ICD-10 (International Classification of Diseases 10<sup>th</sup> Revision) codes (A15–16 and A19, respectively<sup>44</sup>), we categorized disseminated TB as “other infectious disease.” Additionally, several BID cases with a ruptured aortic aneurysm were identified on autopsy, including some that were classified as acute cardiac disease by InterVA. According to the 2016 WHO VA instrument, acute cardiac disease does not cover ruptured aortic aneurysm. However, as the two diseases’ presentations are similar, InterVA misclassified the disease as acute cardiac disease instead of other and unspecified cardiac disease, lowering the positive predictive value. These diseases could be classified in the same category since the interventions are also similar.

Furthermore, there could be other factors related to the automation of VA. First, InterVA may have internal problems leading to a low positive predictive value for COD.<sup>45</sup> The propensity used by InterVA to assign CODs is based only on the presence of signs or symptoms and does not account for the absence of signs or symptoms. For example, a recent negative human immunodeficiency virus (HIV) test result is not considered when determining COD, even though it is helpful. In a previous study, InterVA diagnosed HIV/acquired immune deficiency syndrome (AIDS) more frequently than physician-certified VA (PCVA).<sup>46</sup> Moreover, in this study, for

three BID cases with meningitis as the COD, according to the autopsy, InterVA identified HIV/AIDS as the COD. In addition, even under the best circumstances, there is variation in the presentation of symptoms for a given COD. In the context of VAs, this variation is compounded by the added variability that arises from the interviewee’s ability to recall and correctly identify signs or symptoms. Therefore, a broad variability in the ability to correctly answer the interview questions likely exists.

Second, the automated VA was not validated by directly comparing with the CODs determined using complete autopsies. For the validation, some studies have compared results obtained using automated VA with PCVA reports, which is a method of determining COD via reaching a consensus among physicians based on data obtained using the structured VA questionnaire. The reported concordance between the automated VA tool and PCVA in previous studies ranged from 44.0 to 83.1%.<sup>30,33,47–49</sup> By disease categories, there is a Kenyan study reporting that the concordance for noncommunicable disease CODs and communicable disease CODs based on PCVA was 71.5 and 71.7%, respectively.<sup>49</sup> However, the gold standard for determining CODs may not be the PCVA report, but complete autopsy.<sup>50</sup> Although few studies investigated the validity of PCVA based on complete autopsy,<sup>51</sup> Misganaw et al compared PCVA to medical data

from hospital records in Ethiopia and revealed that the concordance between the two sources was 68% for communicable disease CODs and 79% for NCD CODs.<sup>52</sup> Furthermore, CODs determined using hospital data reviews and complete autopsy results had 30 to 63% discrepancy among 18 previous studies.<sup>53</sup> Taken together, multiplication of these data suggests that estimated concordance between automated VA and complete autopsy should be approximately 20 to 45%. However, direct investigation to compare both methods is necessary to estimate the true concordance.

In addition to the factors possibly responsible for the low positive predictive value mentioned above, there are several limitations to the current study. First, the study population may not represent the general population of BID cases due to the small sample size. Especially, there were only two maternal cases. Also, the case selection could be biased as a substantial number of BID cases that arrived outside of the study period were not included and a large proportion of eligible BID cases were not included due to the lack of consent. This may account for the most common CODs in our study compared with the previous study.<sup>28</sup> Therefore, our results may not be fully generalizable. Furthermore, some subgroups were excluded from this study. In Zambia, cases of death due to external causes require legal procedures for forensic autopsy and the manner of death such as accident, suicide, or homicide is recorded in the forensic report. Since these data were confidential, which meant that the CODs of external cases could not be adequately collected within our research framework, this study focused only on BID cases associated with natural causes of death. Pediatric cases were also excluded owing to the potential challenge of obtaining the consent for autopsy from their parents. Therefore, the validity of results obtained through the InterVA-5 tool compared with those obtained from complete autopsy could not be comprehensively evaluated. Lastly, the quality of the interviewers could also cause the low positive predictive value. For example, data were missing in some cases, especially regarding the duration or the severity of symptoms. While all of the interviewers were trained to conduct VA interviews, the training was provided several years before the research period. As the 2016 WHO VA instrument strongly recommends structured training for interviewers before conducting VA interviews,<sup>37</sup> a lack of refreshment training for interviewers prior to the study period may account for the low positive predictive value.

Considering the factors and the limitations mentioned above, more research is required to investigate the discrepancy between the CODs determined by automated VA and complete autopsy; this work should use larger sample sizes and carry out extensive subgroup analysis. Moreover, updated training for interviewers may significantly improve the two methods' positive predictive values. The effects of refreshment training for interviewers on the results of automated VA in Zambia should indicate the improvements that are necessary for the use of this system in routine national information systems. Automated VA should be primarily utilized to replace PCVA where health resources are limited. The objectives of using automated VA are to

provide better COD statistics in places with resource scarcity and implicate the public health interventions required to address CODs. As **Table 5** shows, automated VA successfully assigned CODs among more than half of BID cases into the same broader disease categories as did autopsy. Classifying community deaths using the broader categories can still be very useful when prioritizing public health issues. The strengths and weaknesses of automated VA programs must be recognized before these processes are put into use in the real-world setting.

## Conclusion

This study compares the CODs assigned to BID cases by automated VA with those determined by complete autopsy to validate the results of automated VA. Several reasons for the low agreement may exist, including the validation method of automated VA, the quality of VA interviews, and the limited lists of CODs available in InterVA-5. Meanwhile, since the concordance under the broader categories was more than 50%, automated VA could be still useful to capture the trend of disease burdens for the public health policy formulation. However, there are several limitations in this study such as small sample size, insufficient training of interviewers, and exclusion of particular subgroups. More research is required to investigate the factors leading to discrepancies between the CODs determined by both methods to optimize the use of automated VA in Zambia.

## Clinical Relevance Statement

Automated VA is useful to classify CODs as per disease conditions among people who died in the communities. However, to scrutinize the CODs, the limitations of automated VA need to be understood. Further research should be considered to optimize the use of automated VA in Zambia.

## Multiple Choice Questions

1. Select the answer from a to d, which indicates the sets of events to be collected for vital statistics purpose by the United Nations.

- ① Marriage
- ② Death
- ③ Legitimation
- ④ Annulment

a. All, b. ① and ②, c. ① to ③, d. ①, ②, and ④

**Correct Answer:** The correct answer is option a. The United Nations defines all events for vital statistical purposes. Refer to <https://unstats.un.org/unsd/demographic/standmeth/principles/m19rev3en.pdf>.

2. What percentage of the countries, territories, and areas in the world has at least 90% coverage of death registration?

- a. 34%
- b. 51%

- c. 68%
- d. 85%

**Correct Answer:** The correct answer is option c. According to the United Nations, 68% of countries, territories, and areas in the world have more than 90% of death registration coverage. Refer to <https://unstats.un.org/unsd/demographic-social/crvs/#coverage>.

#### Protection of Human and Animal Subjects

Ethical approval was obtained from the University of Zambia's Biomedical Research Ethics Committee (ref: 018–12–16) and the Ethics Committee of the National Center for Global Health and Medicine in Japan (ref: NCGM-G-003244–00). Written informed consent was obtained from the closest relatives of the deceased.

#### Funding

This research was supported by NCGM Intramural research Fund (19A03) for the study design, collection, analysis, and interpretation of data, and the necessary publication procedures.

#### Conflict of Interest

None declared.

#### Acknowledgments

The authors thank the Ministry of Health in Zambia, Dr. Kennedy Malama, Dr. Alex Makupe, Ms. Alice Zimba, Ms. Gift Izukanji, and Mr. Ernest Matafwali for their help in data collection, data recording, and data analysis.

#### References

- 1 Mathers CD, Fat DM, Inoue M, Rao C, Lopez AD. Counting the dead and what they died from: an assessment of the global status of cause of death data. *Bull World Health Organ* 2005;83(03):171–177
- 2 Mahapatra P, Shibuya K, Lopez AD, et al; Monitoring Vital Events. Civil registration systems and vital statistics: successes and missed opportunities. *Lancet* 2007;370(9599):1653–1663
- 3 United Nations Statistics Division. Demographic and social statistics [updated October 20, 2020]. Accessed April 20, 2022 at: <https://unstats.un.org/unsd/demographic-social/crvs/>
- 4 Open Working Group for the UN General Assembly. Open Working Group Proposal for Sustainable Development Goals. New York, NY: OWG; 2014
- 5 United Nations. The Road to Dignity by 2030: Ending Poverty, Transforming All Lives and Protecting the Planet Synthesis Report of the Secretary-General on the Post-2015 Agenda. New York, NY: UN2014
- 6 Yokobori Y, Obara H, Sugiura Y, Kitamura T. Gaps in the civil registration and vital statistics systems of low- and middle-income countries and the health sector's role in improving the situation. *Glob Health Med* 2021;3(04):243–245
- 7 Central Statistical Office of Zambia. 2016 Vital Statistics Report. Lusaka: Ministry of Home Affairs of Zambia; 2019
- 8 Central Statistical Office of Zambia. 2015/16 Zambia Sample Vital Registration with Verbal Autopsy Report. Lusaka: Ministry of Home Affairs of Zambia; 2018
- 9 Centre for Global Health Research. What is verbal autopsy? Accessed November 16, 2021 at: <http://www.cghr.org/projects/million-death-study-project/what-is-verbal-autopsy/>
- 10 D'Ambruoso L, Boerma T, Byass P, et al. The case for verbal autopsy in health systems strengthening. *Lancet Glob Health* 2017;5(01):e20–e21
- 11 Durrani MIA, Naz T, Atif M, Khalid N, Amelio A. A semantic-based framework for verbal autopsy to identify the cause of maternal death. *Appl Clin Inform* 2021;12(04):910–923
- 12 World Health Organization. Verbal Autopsy Standards: Ascertain-ing and Attributing Cause of Death. Geneva: WHO; 2007
- 13 Baqui AH, Black RE, Arifeen SE, Hill K, Mitra SN, al Sabir A. Causes of childhood deaths in Bangladesh: results of a nationwide verbal autopsy study. *Bull World Health Organ* 1998;76(02):161–171
- 14 Indian Council of Medical Research. Study on Causes of Death by Verbal Autopsy in India. New Delhi: ICMR; 2009
- 15 Dharmaratne SD, Jayasuriya RL, Perera BY, Gunsekera E, Sathasivayyar A. Opportunities and challenges for verbal autopsy in the national Death Registration System in Sri Lanka: past and future. *Popul Health Metr* 2011;9:21
- 16 França E, Campos D, Guimarães MD, Souza MdeF. Use of verbal autopsy in a national health information system: effects of the investigation of ill-defined causes of death on proportional mortality due to injury in small municipalities in Brazil. *Popul Health Metr* 2011;9:39
- 17 Durrani MIA, Qureshi NS, Ahmad N, Naz T, Amelio A. A health informatics reporting system for technology illiterate workforce using mobile phone. *Appl Clin Inform* 2019;10(02):348–357
- 18 Miasnikof P, Giannakeas V, Gomes M, et al. Naive Bayes classifiers for verbal autopsies: comparison to physician-based classification for 21,000 child and adult deaths. *BMC Med* 2015;13:286
- 19 Flaxman AD, Joseph JC, Murray CJL, Riley ID, Lopez AD. Performance of InSilicoVA for assigning causes of death to verbal autopsies: multisite validation study using clinical diagnostic gold standards. *BMC Med* 2018;16(01):56
- 20 Murtaza SS, Kolpak P, Bener A, Jha P. Automated verbal autopsy classification: using one-against-all ensemble method and Naive Bayes classifier. *Gates Open Res* 2019;2:63
- 21 Nichols EK, Byass P, Chandramohan D, et al; WHO Verbal Autopsy Working Group. The WHO 2016 verbal autopsy instrument: an international standard suitable for automated analysis by InterVA, InSilicoVA, and Tariff 2.0. *PLoS Med* 2018;15(01):e1002486
- 22 Byass P, Hussain-Alkhateeb L, D'Ambruoso L, et al. An integrated approach to processing WHO-2016 verbal autopsy data: the InterVA-5 model. *BMC Med* 2019;17(01):102
- 23 Phillips-Howard PA, Laserson KF, Amek N, et al. Deaths ascribed to non-communicable diseases among rural Kenyan adults are proportionately increasing: evidence from a health and demographic surveillance system, 2003–2010. *PLoS One* 2014;9(11):e114010
- 24 Godefay H, Kinsman J, Admasu K, Byass P. Can innovative ambulance transport avert pregnancy-related deaths? One-year operational assessment in Ethiopia. *J Glob Health* 2016;6(01):010410
- 25 Maraba N, Karat AS, McCarthy K, et al. Verbal autopsy-assigned causes of death among adults being investigated for TB in South Africa. *Trans R Soc Trop Med Hyg* 2016;110(09):510–516
- 26 Coates MM, Kamanda M, Kintu A, et al. A comparison of all-cause and cause-specific mortality by household socioeconomic status across seven INDEPTH network health and demographic surveillance systems in sub-Saharan Africa. *Glob Health Action* 2019;12(01):1608013
- 27 Dandona R, Kumar GA, Bhattacharya D, et al. Distinct mortality patterns at 0–2 days versus the remaining neonatal period: results from population-based assessment in the Indian state of Bihar. *BMC Med* 2019;17(01):140
- 28 Yokobori Y, Matsuura J, Sugiura Y, et al. Analysis of causes of death among brought-in-dead cases in a third-level Hospital in Lusaka, Republic of Zambia, using the tariff method 2.0 for verbal autopsy: a cross-sectional study. *BMC Public Health* 2020;20(01):473

- 29 Serina P, Riley I, Stewart A, et al. A shortened verbal autopsy instrument for use in routine mortality surveillance systems. *BMC Med* 2015;13:302
- 30 Ramroth H, Lorenz E, Rankin JC, et al. Cause of death distribution with InterVA and physician coding in a rural area of Burkina Faso. *Trop Med Int Health* 2012;17(07):904–913
- 31 Lozano R, Freeman MK, James SL, et al; Population Health Metrics Research Consortium (PHMRC) Performance of InterVA for assigning causes of death to verbal autopsies: multisite validation study using clinical diagnostic gold standards. *Popul Health Metr* 2011; 9:50
- 32 Bauni E, Ndila C, Mochamah G, et al. Validating physician-certified verbal autopsy and probabilistic modeling (InterVA) approaches to verbal autopsy interpretation using hospital causes of adult deaths. *Popul Health Metr* 2011;9:49
- 33 Tadesse S. Agreement between physicians and the InterVA-4 model in assigning causes of death: the role of recall period and characteristics specific to the deceased and the respondent. *Arch Public Health* 2013;71(01):28
- 34 Fantahun M, Fottrell E, Berhane Y, Wall S, Högberg U, Byass P. Assessing a new approach to verbal autopsy interpretation in a rural Ethiopian community: the InterVA model. *Bull World Health Organ* 2006;84(03):204–210
- 35 James SL, Flaxman AD, Murray CJ; Population Health Metrics Research Consortium (PHMRC) Performance of the Tariff Method: validation of a simple additive algorithm for analysis of verbal autopsies. *Popul Health Metr* 2011;9:31
- 36 Central Statistical Office of Zambia. 2010 Census of Population and Housing. Analytical Report. Lusaka: CSO; 2012
- 37 World Health Organization. Health statistics and information system. Verbal autopsy standards: ascertaining and attributing cause of death. The 2016 WHO verbal autopsy instrument. 2020. Accessed February 19, 2020 at: <https://score.tools.who.int/tools/count-births-deaths-and-causes-of-death/tool/verbal-autopsy-standards-the-2016-who-verbal-autopsy-instrument-23/>
- 38 Open Data Kit. Collect data anywhere. Accessed February 19, 2020 at: <https://getodk.org/#features>
- 39 Byass P, Huong DL, Minh HV. A probabilistic approach to interpreting verbal autopsies: methodology and preliminary validation in Vietnam. *Scand J Public Health Suppl* 2003;62:32–37
- 40 Byass P, Chandramohan D, Clark SJ, et al. Strengthening standardised interpretation of verbal autopsy data: the new InterVA-4 tool. *Glob Health Action* 2012;5:1–8
- 41 Byass P, Fottrell E, Dao LH, et al. Refining a probabilistic model for interpreting verbal autopsy data. *Scand J Public Health* 2006;34 (01):26–31
- 42 Tensou B, Araya T, Telake DS, et al. Evaluating the InterVA model for determining AIDS mortality from verbal autopsies in the adult population of Addis Ababa. *Trop Med Int Health* 2010;15(05): 547–553
- 43 McHugh ML. Interrater reliability: the kappa statistic. *Biochem Med (Zagreb)* 2012;22(03):276–282
- 44 WHO. ICD-10 Version: 2019. February 28, 2021 Available at: <https://icd.who.int/browse10/2019/en>
- 45 McCormick TH, Li ZR, Calvert C, Crampin AC, Kahn K, Clark SJ. Probabilistic cause-of-death assignment using verbal autopsies. *J Am Stat Assoc* 2016;111(515):1036–1049
- 46 Oti SO, Kyobutungi C. Verbal autopsy interpretation: a comparative analysis of the InterVA model versus physician review in determining causes of death in the Nairobi DSS. *Popul Health Metr* 2010;8:21
- 47 Desai N, Aleksandrowicz L, Miasnikof P, et al. Performance of four computer-coded verbal autopsy methods for cause of death assignment compared with physician coding on 24,000 deaths in low- and middle-income countries. *BMC Med* 2014;12:20
- 48 Byass P, Herbst K, Fottrell E, et al. Comparing verbal autopsy cause of death findings as determined by physician coding and probabilistic modelling: a public health analysis of 54 000 deaths in Africa and Asia. *J Glob Health* 2015;5(01):010402
- 49 Weldearegawi B, Melaku YA, Dinant GJ, Spigt M. How much do the physician review and InterVA model agree in determining causes of death? A comparative analysis of deaths in rural Ethiopia. *BMC Public Health* 2015;15(01):669
- 50 Aase S. Autopsy—still the gold standard? *Tidsskr Nor Laegeforen* 2013;133(07):730–731
- 51 Rodriguez L, Reyes H, Tome P, Ridaura C, Flores S, Guiscafne H. Validation of the verbal autopsy method to ascertain acute respiratory infection as cause of death. *Indian J Pediatr* 1998;65 (04):579–584
- 52 Misganaw A, Mariam DH, Araya T, Aneneh A. Validity of verbal autopsy method to determine causes of death among adults in the urban setting of Ethiopia. *BMC Med Res Methodol* 2012;12:130
- 53 Roulson J, Benbow EW, Hasleton PS. Discrepancies between clinical and autopsy diagnosis and the value of post mortem histology; a meta-analysis and review. *Histopathology* 2005;47 (06):551–559