



Partial Shoulder Arthroplasty Guided by Three-dimensional Prototyping

Artroplastia parcial de ombro guiada por prototipagem tridimensional

Lucas Maia¹ Kennedy Tavares Ladeia¹ Bernardo Figueira Althoff² Adriano Marchetto³
Diego Meneghel⁴ Guilherme Valdir Baldo⁵

¹ Division of Shoulder and Elbow Surgery, Pontifícia Universidade Católica de Campinas (PUC), Campinas, São Paulo, Brazil

² Division of Hand Surgery, Pontifícia Universidade Católica de Campinas (PUC), Campinas, São Paulo, Brazil

³ Division of Shoulder and Elbow Surgery, Instituto Wilson Mello, Campinas, São Paulo, Brazil

⁴ Division of Foot and Ankle Surgery, Escola Paulista de Medicina, Universidade Federal de São Paulo, São Paulo, Brazil

Address for correspondence Lucas Maia, Rua Dom Paulo de Tarso Campos, 234-354, Parque dos Jacarandás, Campinas, São Paulo, Brazil (e-mail: maia1505@gmail.com).

⁵ Division of Shoulder and Elbow Surgery, Centro Universitário para o Desenvolvimento do Alto Vale do Itajaí (UNIDAVI), Rio do Sul, Santa Catarina, Brazil

Rev Bras Ortop 2024;59(Suppl S1):e73–e77.

Abstract

Keywords

- ▶ arthroplasty, replacement, shoulder
- ▶ shoulder joint
- ▶ three-dimensional imaging
- ▶ three-dimensional printing

Three-dimensional (3D) printing technology is a reality in medicine. In Orthopedics and Traumatology, 3D printing guides a precise and tailored surgical treatment. Understanding and disseminating its applicability, use, and outcomes can foster academism and improve patient care. This is a report of a rare case of a female young adult patient with osteonecrosis of the humeral head due to avascular necrosis developed in early childhood. The treatment was tailored and optimized with 3D printing, which helped determine the steps for partial humeral arthroplasty.

Resumo

Palavras-chave

- ▶ articulação do ombro
- ▶ artroplastia do ombro
- ▶ imageamento tridimensional
- ▶ impressão tridimensional

A tecnologia de impressão tridimensional (3D) é uma realidade no âmbito da medicina. Na Ortopedia e Traumatologia, a impressão 3D direciona um tratamento cirúrgico preciso e individualizado. Compreender e difundir suas aplicações, utilidades e resultados pode fomentar o academicismo e melhorar o cuidado ao paciente. Trata-se de relato de um caso raro de uma paciente adulta jovem com osteonecrose da cabeça umeral por necrose avascular desenvolvida na primeira infância. O tratamento foi individualizado e otimizado com a impressão 3D, que ajudou a determinar os passos para o procedimento de artroplastia parcial de úmero.

Study carried out at the Pontifícia Universidade Católica de Campinas (PUC), Campinas, SP, Brazil.

received

August 29, 2021

accepted

November 22, 2021

DOI <https://doi.org/10.1055/s-0042-1749682>.

ISSN 0102-3616.

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

Introduction

The emergence of three-dimensional (3D) printing technology is an industrial milestone.¹ Its system of adding layers by automation using some raw materials, based on pre-formulated prototypes, reinvented the means of production.² It spread to medicine quickly, with applications in various fields, including surgery, reconstruction, and rehabilitation, in addition to the manufacture of synthetic tissues, prostheses, implants, and anatomical models.³ In orthopedics, 3D printing is used in a vast number of conditions, traumas, and sequelae with bone deformities. The use of prototypes designed from computed tomography or magnetic resonance imaging provides a new scenario for diagnosis and treatment in Orthopedics.

With 3D printing, it is possible to create anatomical models, facilitating the understanding of anatomical nuances and helping with surgical planning, including access routes, osteotomies, implant positioning, prostheses preparation, orthotics, and personalized implants.⁴ Studies show that preoperative planning of personalized orthopedic implants using 3D prototypes improves the understanding of fracture pattern when compared to two-dimensional or 3D tests visualized on screen.⁵ We present a case of avascular necrosis of the humeral head in a female patient with a history of hemolytic disease of the newborn who underwent surgical treatment with a partial shoulder arthroplasty planned using 3D printing.

Case Report

This study was approved by the institutional ethics committee under the number CAAE-38891420.5.0000.5453.

The patient was a 23-year-old woman with a history of hemolytic disease of the newborn. Her complicated delivery required blood transfusion and hospitalization in an intensive care unit for 40 days. On admission, she presented osteomyelitis of probable hematogenous origin affecting the left femur and humeral head, which was clinically managed.

During osteoarticular development, the patient presented changes at the left humeral head and femur. It is difficult to

determine whether the changes were due to the infection, secondary complications during birth, or both. At 15-years-old, she underwent a left hip arthroplasty due to pain and loss of range of motion (ROM). At 23 years old, she had a 4 cm shortening of the left upper limb, in addition to pain and functional limitation during gait. On the left shoulder, she had adequate muscle trophism, and the following ROM values: 50° for abduction (ABD), 20° for internal rotation (IR), and 0° for external rotation (ER); her Disabilities of the Arm, Shoulder and Hand (DASH) score was 93 of 150, the American Shoulder and Elbow Surgeons (ASES) score was 10 for pain and 18 for function, and pain was rated as 9 to 10 using the visual analog scale (VAS). Radiographs showed humeral head flattening Creuss stage 4⁶ and an intact glenoid, with a medullary canal size of approximately 1.4 cm (►Fig. 1A and 1B).

A computed tomography scan was used for surgical planning and demonstrated the need for a humeral prosthesis with a fracture nail due to its smaller diameter compared to conventional nails (►Fig. 1C).

Despite the scan, we chose a more precise planning method due to the patient's age. The Renato Archer Information Technology Center (CTI) prepared a 3D-printed prototype using selective laser sintering printing, polyamide PA 12, and a 0.1 mm resolution level between layers. Reconstruction was performed with the Invesalious and Magics (CTI, Campinas, SP, Brazil) software (►Fig. 2). The piece was taken to the laboratory for surgical planning and an Equinox Fracture System (Exactech, Inc., Gainesville, FL, USA) prosthesis was chosen. The need for a greater tuberosity (GT) osteotomy was detected during the procedure as an initially unplanned surgical step (►Fig. 3).

Surgery was performed using a deltopectoral approach, with greater and lesser tuberosity osteotomy. The canal was prepared using a humeral reamer, followed by manual insertion of the orthopedic cement, and application of a 6.5 mm x 120 mm fracture nail. A 1.5 mm offset replicator plate and a 38 mm x 16 mm torque with short humeral head were inserted and sutured to the tuberosities using Fiberwire 2.0 (Arthrex Inc., Naples, FL, USA). The surgery took 1 hour and 23 minutes, and the bleeding was




Fig. 1 Humeral head osteonecrosis: (A) progressive Creuss⁶ stages 1) pre-radiographic stage, 2) subchondral sclerosis, 3) subchondral fracture, 4) evident head collapse, 5) degenerative changes of the head; (B) preoperative radiography; (C) computed tomography scan.




Fig. 2 Full-size, 3D-printed prototype: (A) rear aspect; (B) lateral aspect; (C) top aspect; (D) anteromedial aspect of the humerus (post-test).




Fig. 3 (A) Osteotomy of the humeral head and tuberosities; (B) insertion of the intramedullary nail; (C) verification of the greater tuberosity positioning; (D) post-test outcome at the proximal end of the humerus.

minimal. The patient used a sling for 3 weeks and then physical therapy started. ►**Fig. 4** shows radiographic outcomes.

At a follow-up visit, 1 year and 3 months after surgery, the patient presented good trophism and functional shoulder mobility. The following values were detected on examination: 60° ABD, 20° ER, and 30° IR, as shown in ►**Fig. 5**. Scores

improved for the DASH to 30 of 150, ASES to 0 (pain) and 26 (function), and VAS to 0.

Discussion

Glenohumeral conditions, such as avascular osteonecrosis in young patients, require greater therapeutic engagement due




Fig. 4 Immediate postoperative radiograph: (A) true anteroposterior view; (B) anteroposterior view; (C) oblique view.



Fig. 5 Postoperative clinical outcomes: (A) surgical incision; (B) frontal appearance; (C) external rotation; (D) abduction; (E) hand to mouth; (F) hand to head; (G) internal rotation; (H) hand to back.

to the higher life expectancy and demand. Conservative treatment is the first line. Arthroscopic debridement can delay joint survival. Joint replacement surgeries are indicated when symptoms affect daily activities. Although total arthroplasties have been gaining space, hemiarthroplasties in young patients with a functional glenoid cavity remain a good indication.^{6,7}

Orfaly et al.⁸ presented a series of 21 shoulders with a diagnosis of avascular necrosis of the humeral head submitted to partial arthroplasty with significant improvement in pain (average VAS from 88 to 16 in the whole series, $p < 0.01$), function by the average ASES from 36 to 88 ($p < 0.001$), and ABD went from 88 to 123. Smith et al.⁹ evaluated 32 partial arthroplasties for corticosteroid-associated osteonecrosis with 13 excellent outcomes (42%), four satisfactory outcomes (13%), and 14 unsatisfactory outcomes (45%).

According to the current literature, planning with 3D printing improves shoulder surgery by clarifying the following parameters: compliance, retroversion and glenoid diameter, bone stock, joint surface thickness, posterior offset, and medial offset.¹⁰⁻¹² At a recent literature review, Cordona et al. demonstrated that: (1) while planning total and reverse arthroplasties, the glenoid component is the one benefiting the most from the technique, both in terms of implant and bone stock anticipation; (2) in instabilities, printing can design implants to cover specific bone defects; (3) in trauma, it helps the detailed recognition of the fracture pattern, facilitating the surgical procedure. The authors conclude that the technique is beneficial and can reduce surgical time, blood loss, and fluoroscopy use.¹³

The purpose of planning with a 3D-printed prototype in a young female patient with distorted anatomy was to perform all the steps quickly, accurately, and safely, and to increase prosthesis survival. However, an unforeseen extra step, a GT

osteotomy, was required, potentially leading to implant choice issues and iatrogenic injuries, such as periprosthetic fractures and increased safety for the surgery team during the procedure. Such cases highlight the benefit of clarifying anatomical particularities, reducing surgical time and its implications. Although ABD did not show significant improvement (in addition to difficulties in carrying out rehabilitation due to the COVID-19 pandemic), quality of life and functional scores improved.

Three-dimensional printing for orthopedic planning is a reality. Treatment customization, a better understanding of anatomy, and reduced surgical time are well-established benefits, and its use must be considered for cases in which the anatomy raises questions in planning. The increased use and spread of 3D printing will improve its availability as a surgical tool.

Financial Support

There was no financial support from public, commercial, or non-profit sources.

Conflict of Interests

The authors have no conflict of interests to declare.

References

- 1 Bourell DL, Beaman JJ, Wohlers T. History and Evolution of Additive Manufacturing [Internet]. In: Bourell DL, Frazier W, Kuhn H, Seifi M, editors. Additive Manufacturing Processes; 2020:11–8. Available from: <http://dx.doi.org/10.31399/asm.hb.v24.a0006548>
- 2 Awad A, Trenfield SJ, Gaisford S, Basit AW. 3D printed medicines: A new branch of digital healthcare. *Int J Pharm* 2018;548(01): 586–596
- 3 Ventola CL. Medical Applications for 3D Printing: Current and Projected Uses. *P&T* 2014;39(10):704–711

- 4 Eltorai AE, Nguyen E, Daniels AH. Three-Dimensional Printing in Orthopedic Surgery. *Orthopedics* 2015;38(11):684–687
- 5 Bizzotto N, Sandri A, Regis D, Romani D, Tami I, Magnan B. Three-Dimensional Printing of Bone Fractures: A New Tangible Realistic Way for Preoperative Planning and Education. *Surg Innov* 2015;22(05):548–551
- 6 Dialectos A, Namdari S. Glenohumeral osteoarthritis and the young patient: current options for treatment. *Curr Orthop Pract* 2017;28(02):130
- 7 Barlow JD, Abboud J. Surgical options for the young patient with glenohumeral arthritis. *Int J Shoulder Surg* 2016;10(01):28–36
- 8 Orfaly RM, Rockwood CA Jr, Esenyel CZ, Wirth MA. Shoulder arthroplasty in cases with avascular necrosis of the humeral head. *J Shoulder Elbow Surg* 2007;16(3, Suppl):S27–S32
- 9 Smith RG, Sperling JW, Cofield RH, Hattrup SJ, Schleck CD. Shoulder hemiarthroplasty for steroid-associated osteonecrosis. *J Shoulder Elbow Surg* 2008;17(05):685–688
- 10 Hu H, Liu W, Zeng Q, et al. The Personalized Shoulder Reconstruction Assisted by 3D Printing Technology After Resection of the Proximal Humerus Tumours. *Cancer Manag Res* 2019;11:10665–10673
- 11 Cheriachan DM, DiPaola M, Iannotti JP, Ricchetti ET. 3D Printing in Orthopedics—Upper Extremity Arthroplasty. In: DiPaola M, Wodajo FM, editors. *3D Printing in Orthopaedic Surgery* Philadelphia: Elsevier 2019:151–169. Available from: <https://www.sciencedirect.com/science/article/pii/B9780323581189000130?via%3Dihub>
- 12 Jiang M, Chen G, Coles-Black J, Chuen J, Hardidge A. Three-dimensional printing in orthopaedic preoperative planning improves intraoperative metrics: a systematic review. *ANZ J Surg* 2020;90(03):243–250
- 13 Campana V, Cardona V, Vismara V, et al. 3D printing in shoulder surgery. *Orthop Rev (Pavia)* 2020;12(Suppl 1):8681