

Arterial Thrombotic Events in Hospitalized COVID-19 Patients: A Short Review and Meta-Analysis

Matteo Candeloro, MD^{1,2} Sam Schulman, MD, PhD^{2,3}

¹Department of Innovative Technologies in Medicine and Dentistry, "G. D'Annunzio" University, Chieti, Italy

²Department of Medicine, Thrombosis and Atherosclerosis Research Institute, McMaster University, Hamilton, Ontario, Canada

³Department of Obstetrics and Gynecology, I.M. Sechenov First Moscow State Medical University, Moscow, Russia

Address for correspondence Sam Schulman, MD, PhD, Thrombosis Service, Hamilton General Hospital, 237 Barton Street East, Hamilton, ON L8L 2 × 2, Canada (e-mail: schulms@mcmaster.ca).

Semin Thromb Hemost 2023;49:47–54.

Abstract

It is well established that the risk of venous thromboembolism is high in coronavirus disease 19 (COVID-19). The frequency of arterial thromboembolic events (ATEs) in hospitalized patients with COVID-19 is unclear, as is the magnitude of these events in comparison with other infections. We searched MEDLINE from February 2020 to February 2022 for prospective or retrospective cohort studies and randomized clinical trials that reported the number of acute myocardial infarction (AMI), acute ischemic stroke (AIS), acute limb ischemia (ALI), or other ATE as defined by the original authors in hospitalized patients with COVID-19. The pooled frequencies were calculated through meta-analysis using random effects model with logit transformation and presented with relative 95% prediction intervals (95% PI). We retrieved a total of 4,547 studies, 36 of which (28 retrospective cohorts, five prospective cohorts and three randomized trials) were finally included in our analysis. The resulting cohort counted 100,949 patients, 2,641 (2.6%) of whom experienced ATE. The pooled ATE frequency was 2.0% (95% PI, 0.4–9.6%). The pooled ATE frequency for AMI, AIS, ALI, and other ATE was 0.8% (95% PI, 0.1–8.1%), 0.9% (95% PI, 0.3–2.9%), 0.2% (95% PI, 0.0–4.2%), and 0.5% (95% PI, 0.1–3.0%), respectively. In comparison with the ATE incidence reported in three studies on non-COVID viral pneumonia, we did not detect a significant difference from the results in our analysis. In conclusion, we found a non-negligible proportion of ATE in patients hospitalized for COVID-19. Our results are similar to those found in hospitalized patients with influenza or with non-COVID viral pneumonia.

Keywords

- ▶ COVID-19
- ▶ stroke
- ▶ myocardial infarction
- ▶ ischemia
- ▶ thromboembolism

Coronavirus disease 2019 (COVID-19) has affected over 400 million patients worldwide during the period February 2020 to February 2022.¹ The various mechanisms of action of this new Coronavirus (severe acute respiratory syndrome coronavirus 2 [SARS-CoV-2]) are still partially undetermined. The clinical course spans from a completely asymptomatic disease to a critical clinical situation that requires mechanical ventilation and/or sophisticated management in the intensive care unit.

Among other sequelae, COVID-19 has been associated with higher incidence of venous thrombotic events. In a recent meta-analysis of 20 studies that included 1,988 patients with COVID-19, Di Minno et al found a weighted mean prevalence of venous thromboembolism (VTE) of 31.3% (95% confidence interval [CI], 24.3–39.2%).² Increasing age and high body mass index were also associated with higher VTE prevalence.²

article published online
July 6, 2022

Issue Theme Maintaining Hemostasis and Preventing Thrombosis in COVID-19 —Part IV; Guest Editors: Emmanuel J. Favaloro, PhD, FFSc (RCPA), Leonardo Pasalic, FRCPA, FRACP, PhD, and Giuseppe Lippi, MD

© 2022. Thieme. All rights reserved.
Thieme Medical Publishers, Inc.,
333 Seventh Avenue, 18th Floor,
New York, NY 10001, USA

DOI <https://doi.org/10.1055/s-0042-1749661>.
ISSN 0094-6176.

In an additional study that confirmed the high prevalence of VTE in COVID-19, the incidence of arterial thrombotic events (ATEs) in 16 studies with 7,939 patients was 3.9% (95% CI, 2.0–3.0%).³

Another systematic review of 4,466 patients with COVID-19 from the early phase of the pandemic until May 2020, reported a pooled incidence of acute ischemic stroke (AIS) of 1.2%.⁴ Furthermore, the presence of myocardial injury seems to predict more severe COVID-19.⁵ However, a global and updated assessment of ATE frequency in patients hospitalized for COVID-19 is still missing. Hence, the aim of this review and meta-analysis was to estimate the pooled frequency of ATE in these patients.

Methods

We searched MEDLINE from February 2020 to February 2022. The full search history is shown in ►Supplementary Table S1 (available in the online version only). Retrospective and prospective studies involving patients hospitalized for COVID-19 and that reported the number of any ATE during the hospitalization or within 3 months from discharge were included. We excluded case reports and case series and limited our search to the English language and cohorts with ≥200 patients.

Baseline characteristics as age, sex, intensive care unit (ICU) admission, and treatment with antiplatelet drugs were retrieved. Events of interest were acute myocardial infarction (AMI), AIS, acute limb ischemia (ALI), and a composite of all the other ATE, as defined by the original authors. For the assessment of the quality of studies, we used the Jadad score for randomized controlled trials (RCTs), and the methodo-

logical index for non-randomized studies (MINORS) tool for observational studies.^{6,7}

Statistical Analysis

Random-effects model with logit transformation was used to calculate pooled frequency for ATE with relative 95% CI assessed with Clopper-Pearson method.

We then estimated the 95% prediction intervals (95% PI) for the pooled frequencies. We choose to present the results as pooled frequency and relative 95% PI, which estimates where the true effects are to be expected for 95% of similar studies that might be conducted in the future, and in case of heterogeneity. PI covers a wider range than CI.⁸

Subgroup analyses for AMI, AIS, ALI and all the other ATE were computed with the same method. All analyses were performed using R software, version 4.0.4, and Rstudio version 1.1.423 (2009–2018 RStudio, Inc.), meta package.

Results

The search and screening process are shown in ►Fig. 1. A total of 36 articles were included in the pooled analysis (►Table 1).^{9–44} The majority (28; 78%) consisted of retrospective cohort studies whereas three (8%) were RCTs and five (14%) were prospective cohort studies (►Table 1).

The methodological quality of the 33 cohort studies is summarized in ►Supplementary Table S2 and ►Supplementary Fig. S1, available in the online version only. In total, 17/33 studies had three out of eight MINORS' criteria assessed as inadequate whereas the items "loss to follow-up less than 5%" and "unbiased assessment of the study end point" was considered unclear for all the 33 studies.




Fig. 1 Flow diagram of search results.

Table 1 Characteristics of included studies

| Author, year | Study type | Design | N tot | Age | Sex, female | Anti-platelets | Anti-coagulant | ICU | AMI | AIS | ALI | Other ATE | Total ATE | Death | FU days |
|--|------------|--------|--------|-----------------|--------------|----------------|----------------|--------------|----------|-----------|----------|-----------|-------------|--------------|-------------------|
| Bekelis et al, 2020 ¹¹ | C | R | 2,513 | 66 ^a | 1,111 (44.2) | – | – | – | – | 22 (0.9) | – | – | 22 (0.9) | – | – |
| Hanif et al, 2020 ²² | C | R | 921 | 62 ^b | 347 (37.7) | – | 896 (97.3) | 323 (35.1) | – | 11 (1.2) | 2 (0.2) | – | 13 (1.4) | 325 (35.3) | 9 ^b |
| Lodigiani et al, 2020 ²⁸ | C | R | 388 | 66 ^b | 124 (32) | 93 (24) | 307 (79.1) | 61 (15.7) | 4 (1) | 9 (2.3) | – | – | 13 (3.4) | 92 (23.7) | 10 ^b |
| Piazza et al, 2020 ³³ | C | R | 399 | 62 ^a | 170 (42.6) | 149 (37.3) | 346 (86.7) | 170 (42.6) | 14 (3.5) | 1 (0.3) | – | 11 (2.8) | 26 (6.5) | 55 (13.8) | 30 ^c |
| Rothstein et al, 2020 ³⁶ | C | R | 844 | 59 ^a | 439 (52) | – | – | – | 1 (0.1) | 20 (2.4) | – | – | 20 (2.4) | – | – |
| Alabyad et al, 2021 ⁹ | C | R | 276 | 59 ^a | 130 (47.1) | – | – | 158 (57.2) | 4 (1.4) | 5 (1.8) | – | – | 9 (3.3) | 31 (11.2) | 10 ^b |
| Brandão et al, 2021 ¹³ | C | R | 243 | 63 ^b | 97 (39.9) | – | 222 (91.4) | 243 (100) | 6 (2.5) | 3 (1.2) | 3 (1.2) | – | 12 (4.9) | 77 (31.7) | 28 ^c |
| Caro-Codón et al, 2021 ¹⁴ | C | P | 3,042 | 62 ^a | 1372 (45.1) | 434 (14.3) | 307 (10.1) | – | 5 (0.2) | 18 (0.6) | – | 11 (0.4) | 34 (1.1) | 626 (20.6) | 59 ^b |
| Cho et al, 2021 ¹⁵ | C | R | 2,699 | 59 ^b | 936 (34.7) | – | – | 2,699 (100) | – | 19 (0.7) | – | – | 19 (0.7) | 952 (35.3) | – |
| Constans et al, 2021 ¹⁶ | C | P | 211 | 65 ^a | 118 (55.9) | 36 (17.1) | 0 (0) | 6 (2.8) | 3 (1.4) | 2 (0.9) | – | – | 5 (2.4) | 27 (12.8) | 8 ^b |
| De Michieli et al, 2021 ¹⁷ | C | R | 367 | 61 ^a | 147 (40.1) | – | – | 102 (27.8) | 8 (2.2) | – | – | – | 8 (2.2) | 33 (9) | 30 ^c |
| Fournier et al, 2021 ¹⁸ | C | P | 531 | 66 ^b | 180 (33.9) | 137 (25.8) | 267 (50.3) | 0 (0) | 9 (1.7) | 8 (1.5) | 6 (1.1) | 7 (1.3) | 30 (5.6) | 93 (17.5) | 20 ^b |
| Fröhlich et al, 2021 ¹⁹ | C | R | 6,637 | 70 ^b | 3,132 (47.2) | 912 (13.7) | – | 720 (10.8) | 99 (1.5) | 72 (1.1) | – | – | 171 (2.6) | 1,372 (20.7) | – |
| Giannis et al, 2021 ²⁰ | C | P | 4,906 | 61 ^a | 2,273 (46.3) | 1,073 (21.9) | 612 (12.5) | 578 (11.8) | 24 (0.5) | 22 (0.4) | 26 (0.5) | 22 (0.4) | 84 (1.7) | 3,081 (62.8) | 90 ^c |
| Ilyas et al, 2021 ²³ | C | R | 21,528 | 61 ^a | 9,884 (45.9) | 7,066 (32.8) | 16,300 (75.7) | 6,604 (30.7) | 655 (3) | 168 (0.8) | 44 (0.2) | 168 (0.8) | 1,035 (4.8) | 3,326 (15.4) | 10 ^b |
| Kaptein et al, 2021 ²⁴ | C | R | 947 | 66 ^a | 344 (36.3) | – | 947 (100) | 358 (37.8) | 5 (0.5) | 12 (1.3) | 0 (0) | 3 (0.3) | 20 (2.1) | 144 (15.2) | 9 ^b |
| Kolanko et al, 2021 ²⁵ | C | P | 350 | 67 ^b | 139 (39.7) | 163 (46.6) | 347 (99.1) | 124 (35.4) | 8 (2.3) | 4 (1.1) | – | – | 12 (3.4) | 47 (13.4) | 14 ^b |
| Lekoubou et al, 2021 ²⁶ | C | R | 8,815 | 51 ^a | 4,977 (56.5) | – | – | – | – | 77 (0.9) | – | – | 77 (0.9) | – | – |
| Li et al, 2021 ²⁷ | C | R | 2,804 | 70 ^b | 1,300 (46.4) | 949 (33.8) | 1,840 (65.6) | 440 (15.7) | 64 (2.3) | 45 (1.6) | – | – | 109 (3.9) | – | 42 ^b |
| Lopes et al, 2021 ²⁹ | RCT | P | 615 | 57 ^a | 247 (40.2) | 48 (7.8) | 615 (100) | 39 (6.3) | 27 (4.4) | 1 (0.2) | 1 (0.2) | – | 29 (4.7) | 58 (9.4) | 30 ^c |
| Mendes et al, 2021 ³⁰ | C | R | 265 | 86 ^a | 149 (56.2) | – | – | – | – | 11 (4.2) | – | – | 1 (0.4) | 85 (32.1) | 12.6 ^a |
| Muñoz-Rivas et al, 2021 ³¹ | C | R | 1,127 | 65 ^b | 395 (35) | – | 1127 (100) | 13 (1.2) | 6 (0.5) | 13 (1.2) | 1 (0.1) | 8 (0.7) | 28 (2.5) | – | 14 ^b |
| Purroy and Arqué, 2021 ³⁴ | C | R | 1,737 | 66 ^a | 686 (39.5) | – | – | 117 (6.7) | 8 (0.5) | 6 (0.3) | 0 (0) | – | 14 (0.8) | 276 (15.9) | 7 ^b |
| Qureshi et al, 2021 ³⁵ | C | R | 7,709 | 62 ^a | 4,088 (53) | – | – | 272 (3.5) | – | 103 (1.3) | – | – | 103 (1.3) | – | – |
| Sahai et al, 2021 ³⁷ | C | R | 496 | – | – | 258 (52) | 258 (52) | – | 7 (1.4) | 10 (2) | – | – | 17 (3.4) | – | – |
| Siegler et al, 2021 ³⁸ | C | R | 14,483 | – | – | – | – | – | – | 156 (1.1) | – | – | 156 (1.1) | 1,694 (11.7) | – |
| Smadja et al, 2021 ³⁹ | C | R | 1,044 | 63 ^b | 391 (37.5) | 244 (23.4) | 1044 (100) | 157 (15) | 9 (0.9) | 5 (0.5) | 3 (0.3) | – | 17 (1.6) | 93 (8.9) | – |
| Spyropoulos et al, 2021 ⁴⁰ | RCT | P | 253 | 67 ^a | 117 (46.2) | 40 (15.8) | 253 (100) | 83 (32.8) | 3 (1.2) | 2 (0.8) | 2 (0.8) | 1 (0.4) | 8 (3.2) | 56 (22.1) | 30 ^c |
| Tacquard et al, 2021 ⁴¹ | C | R | 538 | 63 ^b | 149 (27.7) | 111 (20.6) | 538 (100) | 538 (100) | 1 (0.2) | 4 (0.7) | – | 1 (0.2) | 6 (1.1) | 1 (0.2) | 14 ^c |
| Topcu et al, 2021 ⁴² | C | R | 681 | 63 ^b | 262 (38.5) | – | – | 90 (13.2) | – | – | 6 (0.9) | – | 6 (0.9) | 94 (13.8) | 30 ^c |
| Vallone et al, 2021 ⁴³ | C | R | 1,621 | 73 ^b | 913 (56.3) | 689 (42.5) | 1,074 (66.3) | 174 (10.7) | 0 (0) | 4 (0.2) | 0 (0) | 1 (0.1) | 5 (0.3) | 348 (21.5) | 90 ^c |
| Violi et al, 2021 ⁴⁴ | C | R | 373 | 67 ^a | 145 (38.9) | 56 (15) | – | 67 (18) | 15 (4) | 10 (2.7) | 9 (2.4) | – | 34 (9.1) | 75 (20.1) | 18 ^b |
| Arachchilage et al, 2022 ¹⁰ | C | R | 5,883 | 74 ^b | 2,636 (44.8) | 1,093 (18.6) | 5,883 (100) | 789 (13.4) | 12 (0.2) | 49 (0.8) | – | 18 (0.3) | 79 (1.3) | 1,719 (29.2) | 90 ^c |
| Bikdeli et al, 2022 ¹² | RCT | P | 562 | 62 ^b | 237 (42.2) | 172 (30.6) | 562 (100) | 562 (100) | 0 (0) | 2 (0.4) | – | – | 2 (0.4) | 250 (44.5) | 90 ^c |

(Continued)

Table 1 (Continued)

| Author, year | Study type | Design | N tot | Age | Sex, female | Anti-platelets | Anti-coagulant | ICU | AMI | AIS | ALI | Other ATE | Total ATE | Death | FU days |
|---|------------|--------|-------|-----------------|-------------|----------------|----------------|-----------|---------|---------|---------|-----------|-----------|------------|---------|
| Gonzalez-Porras et al, 2022 ²¹ | C | R | 690 | 72 ^a | 274 (39.7) | – | 615 (89.1) | 80 (11.6) | 3 (0.4) | 2 (0.3) | 1 (0.1) | – | 6 (0.9) | 263 (38.1) | – |
| Nemetski et al, 2022 ³² | C | R | 4,451 | – | – | – | – | – | – | – | – | – | 411 (9.2) | – | – |

Abbreviations: AIS, acute ischemic stroke; ALI, acute limb ischemia; AMI, acute myocardial infarction; ATE, acute thrombotic event; C, cohort; FU, follow-up; ICU, intensive care unit; P, prospective; R, retrospective.

^aMean.

^bMedian

^cPlanned follow-up duration.

We assessed one of the three RCTs as having a low quality with only one point on the Jadad scale (► **Supplementary Table S3**, available in the online version only).¹²

The resulting population included 100,949 patients. The age of the patients in the studies ranged from a mean of 51 to 86 years (18 studies) or a median of 59 to 74 years (15 studies). Point estimates for age were not available in three cohort studies.

Overall, 37,909 of 81,519 patients (46.5%, 33 studies) were female. Treatment in the ICU was required for 15,567 out of 66,040 patients (23.6%; 28 studies). Furthermore, 13,723 out of 52,181 patients (26.3%, 19 studies) were taking an anti-platelet agent at baseline. Information on anticoagulant treatment was available in 22 studies, and was given to 34,360 out of 49,099 patients (70.0%). Anticoagulant treatment included both prophylactic and therapeutic regimens with low molecular weight heparin, unfractionated heparin, vitamin K antagonists, or direct oral anticoagulants. The follow-up or mean/median hospital stay duration was reported by 17 studies and ranged from 7 to 90 days. Information on mortality was available in 28 studies and 15,293 of 72,190 patients (21.2%) died, with the proportion across the studies ranging from 0.2 to 62.8% (► **Table 1**).

Overall, there were 2,641 ATE (2.6%). The frequency of ATE ranged from 0.3 to 9.2% in the studies included. The pooled frequency was 2.0% with a quite large prediction interval (95% PI, 0.4–9.6%) (► **Fig. 2**).

The pooled frequency was lower for RCTs (1.2%) than for retrospective and prospective cohort studies (1.9 and 2.4%, respectively) (► **Fig. 2**).

The forest plots for the subgroup analyses are displayed in ► **Fig. 3**. In total, 27 studies provided data on AMI for a total of 1,000 of 58,412 (1.7%) patients. After meta-analysis, the pooled frequency turned out to be 0.8% with 95% PI ranging from 0.1 to 8.1%. AIS occurred in 896 of 95,450 patients (0.9%; 33 studies) for a pooled frequency of 0.9% (95% PI, 0.3 to 2.9%). The same figure for ALI was 104 of 37,217 patients (0.3%; 15 studies), pooled frequency 0.2% (95% PI, 0.0–4.2%) (► **Fig. 3**).

Other ATE events occurred in 251 of 40,775 patients (0.6%, 11 studies). These events included 16 systemic embolism, 1 mesenteric infarction, 11 catheter- or device-related arterial thrombosis (all in ICU patients), and the remaining 223 included aortic thrombosis, renal artery thrombosis, splenic infarction, upper-limb ischemia, and acute diffuse microvascular cerebrovascular pathology. The resulting pooled frequency for these events was 0.5% (95% PI, 0.1–3.0%) (► **Fig. 3**).

Discussion

In this systematic review we found a pooled frequency of ATE in COVID-19 hospitalized patients of 2.0% (95% PI, 0.4–9.6). In the subgroup analysis, AIS showed a pooled frequency of 0.9%, followed by AMI (0.8%), other ATE (0.5%), and ALI (0.2%).

Our study covered 2 years from the beginning of the pandemic. Compared with previous reviews and meta-analyses that analyzed only the first part of the pandemic, we found a frequency of AIS that was slightly lower (0.9 vs.




Fig. 2 Forest plot for ATE in overall cohort. ATE, arterial thrombotic events.




Fig. 3 Forest plots for subgroups analysis. (A) AMI, acute myocardial infarction; (B) AIS, acute ischemic stroke; (C) ALI, acute limb ischemia; (D) ATE, arterial thrombotic events.

1.2%).⁵ A previous review reported an extremely high event rate of acute myocardial injury (20%; 95% CI, 17–23%), which is approximately 20 times higher than the pooled frequency we found in our study.⁴⁵ However, the authors included cohorts with at least 100 patients and did not perform a systematic search. Moreover, all the included cohorts but two were from China and the observation covered the early COVID-19 pandemic till May 2020,⁴⁵ and thus before emergence of new viral lineages which may possess lower virulence and pathogenicity.⁴⁶

Accordingly, the pooled frequency of ATE in the included studies tended to decrease from 2020 to 2022, when the Omicron lineages BA.1 and BA.2 become worldwide dominant (data not shown).

If there truly is a decreasing frequency of ATE with time in this pandemic, this could be associated with some other reasons along with emergence of new SARS-CoV-2 variants. First, in the early part of the pandemic there were mainly

case reports and studies based on small cohorts of COVID-19 patients. This could have been led to an overestimation of the true frequency of the ATE. Second, increased attention to antithrombotic prophylaxis during hospitalization in COVID-19 patients began to occur only after the initial months of the pandemic, as attested by the RCTs in this regard.^{12,29,40} This might have reduced the frequency of ATE compared with the early stages of the pandemic. Third, since December 2020 and throughout 2021, a massive vaccine campaign was implemented in the countries where the studies have been performed. This has likely contributed to reduce the frequency of COVID-19 in its most severe form and probably also the frequency of ATE. Moreover, some studies addressed the importance of influenza vaccines in reducing the frequency of ATE in patients with influenza.⁴⁷ As a result, although on a purely conjectural level, this could also have happened in patients with COVID-19. Hence, if this trend is maintained as the vaccination campaign progresses,

Table 2 ATE frequency in our pooled cohort vs viral pneumonia cohort (influenza and non-influenza)

| ATE type | COVID-19 Pooled cohort = 100,949 | Non-COVID-19 viral pneumonia Total cohort = 455,629 ^a |
|----------|----------------------------------|--|
| AMI | 0.8% | 1.3% |
| AIS | 0.9% | 0.7% |
| ALI | 0.2% | 0.07% |

Abbreviations: AIS, acute ischemic stroke; ALI, acute limb ischemia; AMI, acute myocardial infarction; ATE, arterial thromboembolic event.

^aData are from Elgendy et al.⁴⁹

we could expect a further decrease in the frequency of ATE in COVID-19.

Some studies have compared the incidence of ATE between COVID-19 and influenza. In a retrospective cohort, the unadjusted incidence of ATE (AIS + AMI) was more than double in COVID-19 than in influenza (7.1 vs. 3.0%).⁴⁸ However, after propensity score matching there was no difference between the two groups (hazard ratio 1.02, 95% CI, 0.95–1.10).⁴⁸ Nemetski et al found a slightly higher incidence of ATE in patients with COVID-19 ($n=4,451$) when compared with patients hospitalized for influenza ($n=468$) or non-COVID-19 illness ($n=40,359$) and, notably, none of the influenza patients experienced an ATE.³² However, the study was retrospective and the real incidence of ATE in the non-COVID-19 population could have been underestimated.

On the other hand, our data are comparable to those found in a large cohort ($n=455,629$) of patients hospitalized with viral pneumonia (influenza and non-influenza).⁴⁹ Indeed, in this cohort the incidence of ATE was 1.3% for AMI, 0.7% for AIS, and 0.07% for ALI (►Table 2).⁴⁹ This supports the assumption that the incidence of ATE may be similar in patients hospitalized with COVID-19 or with viral influenza or pneumonia. Therefore, whereas it has been proposed that endothelial activation and dysfunction in COVID-19 could be a mechanism that explains the increase in pulmonary thrombosis,⁵⁰ this thromboinflammation might not affect the arteries in the systemic circulation.

This study has limitations that need to be considered. First, because of aggregate data we could not perform more granular subgroup analyses. However, we had enough information to estimate a pooled frequency of ATE as planned. Second, wide variations in duration of follow-up probably contributed to wide prediction intervals.

In conclusion, the frequency of ATE in patients hospitalized for COVID-19 is not negligible although it might have diminished compared with initial estimates from the pandemic onset. When compared with patients hospitalized for viral pneumonia the frequency of ATE is similar.

Funding

M.C. is receiving a salary from “G. d’Annunzio” University for his research fellowship.

Conflict of Interest

M.C. has nothing to disclose. S.S. reports grants and personal fees from Boehringer Ingelheim, grants from Octapharma, personal fees from Bayer, personal fees from Bristol-Myers Squibb, personal fees from Daiichi-Sankyo, personal fees from Pfizer, personal fees from Sanofi, personal fees from Alexion, outside the submitted work.

References

- COVID-19 Dashboard by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University (JHU). Accessed March 1, 2022 at: <https://www.arcgis.com/apps/dashboards/bda7594740fd40299423467b48e9ecf6>
- Di Minno A, Ambrosino P, Calcaterra I, Di Minno MND. COVID-19 and venous thromboembolism: a meta-analysis of literature studies. *Semin Thromb Hemost* 2020;46(07):763–771
- Tan BK, Mainbourg S, Friggeri A, et al. Arterial and venous thromboembolism in COVID-19: a study-level meta-analysis. *Thorax* 2021;76(10):970–979
- Tan YK, Goh C, Leow AST, et al. COVID-19 and ischemic stroke: a systematic review and meta-summary of the literature. *J Thromb Thrombolysis* 2020;50(03):587–595
- Alali AH, Smaiesm MS, Alsheikh AM, et al. Myocardial injuries among patients with COVID-19: a systematic review. *Infez Med* 2021;29(03):345–354
- Jadad AR, Moore RA, Carroll D, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials* 1996;17(01):1–12
- Slim K, Nini E, Forestier D, Kwiatkowski F, Panis Y, Chipponi J. Methodological index for non-randomized studies (minors): development and validation of a new instrument. *ANZ J Surg* 2003;73(09):712–716
- Int’Hout J, Ioannidis JPA, Rovers MM, Goeman JJ. Plea for routinely presenting prediction intervals in meta-analysis. *BMJ Open* 2016; 6(07):e010247
- Alabyad D, Rangaraju S, Liu M, et al. Validation of an admission coagulation panel for risk stratification of COVID-19 patients. *PLoS One* 2021;16(03):e0248230
- Arachchillage DJ, Rajakaruna I, Odho Z, et al. Clinical outcomes and the impact of prior oral anticoagulant use in patients with coronavirus disease 2019 admitted to hospitals in the UK – a multicentre observational study. *Br J Haematol* 2022;196(01):79–94
- Bekelis K, Missios S, Ahmad J, et al. Ischemic stroke occurs less frequently in patients with COVID-19: a multicenter cross-sectional study. *Stroke* 2020;51(12):3570–3576
- Bikdeli B, Talasaz AH, Rashidi F, et al. Intermediate-dose versus standard-dose prophylactic anticoagulation in patients with COVID-19 admitted to the intensive care unit: 90-day results from the INSPIRATION randomized trial. *Thromb Haemost* 2022; 122(01):131–141
- Brandão AAGS, de Oliveira CZ, Rojas SO, et al. Thromboembolic and bleeding events in intensive care unit patients with COVID-19: results from a Brazilian tertiary hospital. *Int J Infect Dis* 2021; 113:236–242
- Caro-Codón J, Lip GYH, Rey JR, et al. Prediction of thromboembolic events and mortality by the CHADS2 and the CHA2DS2-VASc in COVID-19. *Europace* 2021;23(06):937–947
- Cho SM, Premraj L, Fanning J, et al. Ischemic and hemorrhagic stroke among critically ill patients with coronavirus disease 2019: an international multicenter coronavirus disease 2019 critical care consortium study. *Crit Care Med* 2021;49(12):e1223–e1233
- Constans M, Santiago R, Jimenez L, et al. Lupus anticoagulant is an independent risk factor for non-thrombotic in-hospital mortality in COVID-19 patients. *Thromb Res* 2021;208(October):99–105

- 17 De Michieli L, Ola O, Knott JD, et al. High-sensitivity cardiac troponin T for the detection of myocardial injury and risk stratification in COVID-19. *Clin Chem* 2021;67(08):1080–1089
- 18 Fournier M, Faïlle D, Dossier A, et al. Arterial thrombotic events in adult inpatients with COVID-19. *Mayo Clin Proc* 2021;96(02):295–303
- 19 Fröhlich GM, Jeschke E, Eichler U, et al. Impact of oral anti-coagulation on clinical outcomes of COVID-19: a nationwide cohort study of hospitalized patients in Germany. *Clin Res Cardiol* 2021;110(07):1041–1050
- 20 Giannis D, Allen SL, Tsang J, et al. Postdischarge thromboembolic outcomes and mortality of hospitalized patients with COVID-19: the CORE-19 registry. *Blood* 2021;137(20):2838–2847
- 21 Gonzalez-Porras JR, Belhassen-Garcia M, Lopez-Bernus A, et al. Low molecular weight heparin is useful in adult COVID-19 inpatients. Experience during the first Spanish wave: observational study. *Sao Paulo Med J* 2022;140(01):123–133
- 22 Hanif A, Khan S, Mantri N, et al. Thrombotic complications and anticoagulation in COVID-19 pneumonia: a New York City hospital experience. *Ann Hematol* 2020;99(10):2323–2328
- 23 Ilyas S, Henkin S, Martinez-Cambor P, et al. Sex-, race- and ethnicity-based differences in thromboembolic events among adults hospitalized with COVID-19. *J Am Heart Assoc* 2021;10(23):e022829
- 24 Kaptein FHJ, Stals MAM, Grootenboers M, et al; Dutch COVID & Thrombosis Coalition. Incidence of thrombotic complications and overall survival in hospitalized patients with COVID-19 in the second and first wave. *Thromb Res* 2021;199(199):143–148
- 25 Kolanko E, Senderek T, Prokop-Staszecka A, et al. Thromboprophylaxis in hospitalized COVID-19 patients: the efficacy and safety of the approved hospital protocol. *Pol Arch Intern Med* 2021;131(10):16102
- 26 Lekoubou A, Pelton M, Ba DM, Ssentongo P. Racial disparities in ischemic stroke among patients with COVID-19 in the United States. *J Stroke Cerebrovasc Dis* 2021;30(08):105877
- 27 Li A, Kuderer NM, Hsu CY, et al; CCC19 consortium. The CoVID-TE risk assessment model for venous thromboembolism in hospitalized patients with cancer and COVID-19. *J Thromb Haemost* 2021;19(10):2522–2532
- 28 Lodigiani C, Iapichino G, Carenzo L, et al; Humanitas COVID-19 Task Force. Venous and arterial thromboembolic complications in COVID-19 patients admitted to an academic hospital in Milan, Italy. *Thromb Res* 2020;191(April):9–14
- 29 Lopes RD, de Barros E Silva PGM, Furtado RHM, et al; ACTION Coalition COVID-19 Brazil IV Investigators. Therapeutic versus prophylactic anticoagulation for patients admitted to hospital with COVID-19 and elevated D-dimer concentration (ACTION): an open-label, multicentre, randomised, controlled trial. *Lancet* 2021;397(10291):2253–2263
- 30 Mendes A, Herrmann FR, Genton L, et al. Incidence, characteristics and clinical relevance of acute stroke in old patients hospitalized with COVID-19. *BMC Geriatr* 2021;21(01):52
- 31 Muñoz-Rivas N, Abad-Motos A, Mestre-Gómez B, et al; Infanta Leonor Thrombosis Research Group. Systemic thrombosis in a large cohort of COVID-19 patients despite thromboprophylaxis: a retrospective study. *Thromb Res* 2021;199(January):132–142
- 32 Nemetski SM, Ip A, Josephs J, Hellmann M. Clotting events among hospitalized patients infected with COVID-19 in a large multisite cohort in the United States. *PLoS One* 2022;17(01):e0262352
- 33 Piazza G, Campia U, Hurwitz S, et al. Registry of arterial and venous thromboembolic complications in patients with COVID-19. *J Am Coll Cardiol* 2020;76(18):2060–2072
- 34 Purroy F, Arqué G. Influence of thromboembolic events in the prognosis of COVID-19 hospitalized patients. Results from a cross sectional study. *PLoS One* 2021;16(06):e0252351
- 35 Qureshi AI, Baskett WI, Huang W, et al. Acute ischemic stroke and COVID-19: an analysis of 27 676 patients. *Stroke* 2021;52(03):905–912
- 36 Rothstein A, Oldridge O, Schwennesen H, Do D, Cucchiara BL. Acute cerebrovascular events in hospitalized COVID-19 patients. *Stroke* 2020;51(09):e219–e222
- 37 Sahai A, Bhandari R, Godwin M, et al. Effect of aspirin on short-term outcomes in hospitalized patients with COVID-19. *Vasc Med* 2021;26(06):626–632
- 38 Siegler JE, Cardona P, Arenillas JF, et al. Cerebrovascular events and outcomes in hospitalized patients with COVID-19: the SVIN COVID-19 Multinational Registry. *Int J Stroke* 2021;16(04):437–447
- 39 Smadja DM, Bonnet G, Gendron N, et al. Intermediate- vs. standard-dose prophylactic anticoagulation in patients with COVID-19 admitted in medical ward: a propensity score-matched cohort study. *Front Med (Lausanne)* 2021;8(October):747527
- 40 Spyropoulos AC, Goldin M, Giannis D, et al; HEP-COVID Investigators. Efficacy and safety of therapeutic-dose heparin vs standard prophylactic or intermediate-dose heparins for thromboprophylaxis in high-risk hospitalized patients with COVID-19: the HEP-COVID randomized clinical trial. *JAMA Intern Med* 2021;181(12):1612–1620
- 41 Tacquard C, Mansour A, Godon A, et al; French Working Group on Perioperative Hemostasis. Impact of high-dose prophylactic anticoagulation in critically ill patients with COVID-19 pneumonia. *Chest* 2021;159(06):2417–2427
- 42 Topcu AC, Ozturk-Altunyurt G, Akman D, Batirel A, Demirhan R. Acute limb ischemia in hospitalized COVID-19 patients. *Ann Vasc Surg* 2021;74(April):88–94
- 43 Vallone MG, Vazquez C, Chuliber FA, et al. Low incidence of symptomatic thrombotic events in adult patients hospitalized with coronavirus 19: a retrospective cohort study. *Clin Appl Thromb Hemost* 2021;27:10760296211051712
- 44 Violi F, Ceccarelli G, Cangemi R, et al; Intensive Care, Infectious Diseases COVID-19 Study Group of Sapienza University. Arterial and venous thrombosis in coronavirus 2019 disease (Covid-19): relationship with mortality. *Intern Emerg Med* 2021;16(05):1231–1237
- 45 Bavishi C, Bonow RO, Trivedi V, Abbott JD, Messerli FH, Bhatt DL. Special article—acute myocardial injury in patients hospitalized with COVID-19 infection: a review. *Prog Cardiovasc Dis* 2020;63(05):682–689
- 46 Lippi G, Mattiuzzi C, Henry BM. Updated picture of SARS-CoV-2 variants and mutations. *Diagnosis (Berl)* 2021;9(01):11–17
- 47 Yedlapati SH, Khan SU, Talluri S, et al. Effects of influenza vaccine on mortality and cardiovascular outcomes in patients with cardiovascular disease: a systematic review and meta-analysis. *J Am Heart Assoc* 2021;10(06):e019636
- 48 Ward A, Sarraju A, Lee D, et al. COVID-19 is associated with higher risk of venous thrombosis, but not arterial thrombosis, compared with influenza: insights from a large US cohort. *PLoS One* 2022;17(01):e0261786
- 49 Elgendy IY, Kolte D, Mansour MK, et al. Incidence, predictors, and outcomes of thrombotic events in hospitalized patients with viral pneumonia. *Am J Cardiol* 2021;143:164–165
- 50 Schulman S, Hu Y, Konstantinides S. Venous thromboembolism in COVID-19. *Thromb Haemost* 2020;120(12):1642–1653