



Predicting the Position of the Internal Landmarks of Middle Cranial Fossa Using the Zygomatic Root: An Attempt to Simplify Its Complexity

Chandrasekaran Kaliaperumal¹

¹ Department of Clinical Neuroscience, DCN and RHCYP Royal Infirmary of Edinburgh, Edinburgh, United Kingdom

Indian J Neurosurg

Address for correspondence Chandrasekaran Kaliaperumal, DipMedEd, FEBNS, FRCSI, FRCSEd (Neuro.Surg), Department of Clinical Neuroscience, DCN and RHCYP Royal Infirmary of Edinburgh, Edinburgh, EH16 4SA, United Kingdom (e-mail: ckaliaperumal@gmail.com).

Neurosurgery is a rapidly ever evolving, challenging, and young specialty. We are in a race toward technological advancement to achieve excellence in postoperative outcome, avoiding and anticipating complications. Skull base neurosurgery has moved significantly forward with pioneering contributions from eminent neurosurgeons to enrich neuroanatomical knowledge, promoting safe and innovative surgical approaches.

We have evidence on temporal bone embryology and its development.^{1,2} Considering wide variability in the skulls noted in Caucasian and Indian skulls, the current evidence that the mean depths of the lateral projections are smaller and greater distance from anterior root of lateral projections is a useful tool.³ This leaves the corridor open for further anatomical studies.

Patients with trigeminal neuralgia receiving foramen ovale (FO) injection or balloon rhizolysis, radio frequency ablation of trigeminal nerve often pose a challenge in terms of anatomical localization and knowledge about the middle cranial fossa anatomy will potentially avoid failure of procedure and subsequent complications. Zygomatic root (ZR) in this anatomical study has shown that in 89% of specimens analyzed, the FO projected on to the anterior two-third of ZR.³ This is practically a useful tool both for the neurosurgeons and interventional neuroradiologists in terms of localization of FO. ZR identification can help prevention of facial droop by preserving frontalis branch of facial nerve and assist in epilepsy surgery by identifying the temporal horns of lateral ventricle, temporal tip, and the inferior temporal gyrus.^{4,5}

Virtual reality (VR), augmented reality, and haptic simulation are burgeoning in neurosurgery particularly to aid preoperative surgical planning, mental rehearsal, and training residents.^{6–9} They eliminate the task of having to mentally reconstruct and rotate triaxial images and provide additional information on the real operative field that is otherwise invisible and inaccessible.⁶ VR technology might improve neurosurgical skull base teaching quality, which should be promoted in the teaching of clinical subjects.⁷ Case-specific VR-based surgical simulation is shown to allow the users to intuitively understand the positional relationships between organs, blood vessels, and lesions within a patient's body, resulting in assistance for surgical planning and intraoperative management.⁸ “Cost factor” may be a limitation for the utility of these technological adjuncts in the low- and middle-income countries in terms of access but this will hopefully be more accessible and affordable in the near future.

Preoperative computed tomography along with magnetic resonance imaging brain and an understanding of applied neuroanatomy are vital to visualize, plan, and guide the operating neurosurgeon during middle and posterior fossa surgical approach for a variety of neurosurgical pathologies.⁷ Haptic simulation as a tool is emerging and may be a norm in the future albeit it has its limitations.⁹ Neuroanatomical nuance and understanding will help vascular, trauma, functional, oncological neurosurgeons and ENT surgeons (who form a part of the skull base team). As elucidated by the authors in this study, the reliability of ZR in 80% predicting the position of FO, foramen spinosum, and

DOI <https://doi.org/10.1055/s-0042-1750297>.
ISSN 2277-954X.

© 2022. Neurological Surgeons' Society of India. All rights reserved. This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)
Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

trigeminal fossa will remain a useful adjunct to existing technology in delivering a “safe and sustainable” care package to the global neurosurgical patients.³

Conflict of Interest

None declared.

References

- 1 Oettlé AC, Demeter FP, L'abbé EN. Ancestral variations in the shape and size of the zygoma. *Anat Rec (Hoboken)* 2017;300(01):196–208
- 2 Dechow PC, Wang Q. Development, structure, and function of the zygomatic bones: what is new and why do we care? *Anat Rec (Hoboken)* 2016;299(12):1611–1615. Doi: 10.1002/ar.23480
- 3 Current study: Predicting the position of the internal landmarks of middle cranial fossa using the zygomatic root: an attempt to simplify its complexity.
- 4 Dayoub H, Schueler WB, Shakir H, Kimmell KT, Sincoff EH. The relationship between the zygomatic arch and the floor of the middle cranial fossa: a radiographic study. *Neurosurgery* 2010;66(6, Suppl Operative):363–369
- 5 Beckman JM, Vale FL. Using the zygomatic root as a reference point in temporal lobe surgery. *Acta Neurochir (Wien)* 2013;155(12):2287–2291
- 6 Rosahl SK, Gharabaghi A, Hubbe U, Shahidi R, Samii M. Virtual reality augmentation in skull base surgery. *Skull Base* 2006;16(02):59–66
- 7 Shao X, Yuan Q, Qian D, et al. Virtual reality technology for teaching neurosurgery of skull base tumor. *BMC Med Educ* 2020;20(01):3. Doi: 10.1186/s12909-019-1911-5
- 8 Ito T, Kawashima Y, Yamazaki A, Tsutsumi T. Application of a virtual and mixed reality-navigation system using commercially available devices to the lateral temporal bone resection. *Ann Med Surg (Lond)* 2021;72:103063. Doi: 10.1016/j.amsu.2021.103063
- 9 Wong D, Unger B, Kraut J, Pisa J, Rhodes C, Hochman JB. Comparison of cadaveric and isomorphic virtual haptic simulation in temporal bone training. *J Otolaryngol Head Neck Surg* 2014;43(01):31. Doi: 10.1186/s40463-014-0031-9