



# $^{68}\text{Ga}$ -DOTATATE Uptake in COVID-19 Pneumonia: A Non-Neoplastic False Positive

Joshua P. Weissman<sup>1</sup> Twyla Bartel<sup>2</sup>

<sup>1</sup>Northwestern University Feinberg School of Medicine, Chicago, Illinois, United States

<sup>2</sup>Global Advanced Imaging, PLLC, Little Rock, Arkansas, United States

Address for correspondence Twyla Bartel, MD, Global Advanced Imaging, PLLC, Little Rock, 225 E. Chicago Ave., Box 93, Little Rock, AR 72223, United States (e-mail: twylabb@hotmail.com).

World J Nuclear Med 2022;21:163–165.

## Abstract

In the detection and characterization of neuroendocrine tumors (NETs), the NETSPOT  $^{68}\text{Ga}$ -DOTATATE positron emission tomography/computed tomography (PET/CT) utilizes a somatostatin analog to capture whole body imaging of somatostatin receptors. Here, we report a rare case of incidental coronavirus disease 2019 (COVID-19) detection using  $^{68}\text{Ga}$ -DOTATATE PET/CT for a 48-year-old patient with COVID-19 who presented to our nuclear medicine department for oncologic and neuroendocrine indications. Current maximal intensity project image from  $^{68}\text{Ga}$ -DOTATATE PET/CT demonstrated numerous hypermetabolic foci in the upper abdominal lymph nodes, consistent with somatostatin-avid pancreatic islet cell neoplasm. Radiotracer uptake corresponding to the anatomic infiltrates was also seen in the bilateral pneumonia. With the rise of worldwide COVID-19 cases, and the continued use of NETSPOT for somatostatin avid neoplasms, it is imperative that diagnostic measures of this disease are clearly understood. This case highlights the importance of including COVID-19 on the differential diagnosis when radiotracer uptake appears in  $^{68}\text{Ga}$ -DOTATATE PET/CT.

## Keywords

- ▶ Netspot
- ▶ coronavirus
- ▶ PET

## Key Messages

COVID-19 should be included in the differential diagnosis when radiotracer uptake appears in NETSPOT  $^{68}\text{Ga}$ -DOTATATE PET/CT. Current documented false-positive findings for  $^{68}\text{Ga}$ -DOTATATE PET/CT include pancreatic uncinat process activity, prostatitis, splenosis, postradiation therapy change, paraganglioma, pheochromocytoma, neuroblastoma, meningioma, and osteoblastic activity.

## Introduction

The coronavirus disease 2019 (COVID-19) outbreak has infected over 131 million people and caused 2.85 million deaths worldwide, as of early April 2021.<sup>1</sup> In many patients,

COVID-19 presents with nonspecific symptoms such as fever, cough, and chest pain, while in some others, the infection remains asymptomatic.<sup>2</sup> Despite its suboptimal sensitivity, nucleic acid testing (reverse transcription polymerase chain reaction [RT-PCR]) has been regarded as the gold standard to confirm the diagnosis of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). Imaging can be a useful adjunct. In symptomatic patients, chest X-rays are typically obtained. Computed tomography (CT) chest may be added as a key supplemental tool for the diagnosis of SARS-CoV-2 given its reported improved sensitivity relative to RT-PCR (98 vs. 71%).<sup>3</sup> Recently, several groups have reported the incidental [ $^{18}\text{F}$ ]-2-fluoro-2-deoxy-D-glucose (FDG)—positron emission tomography/computed tomography (PET/CT) imaging

DOI <https://doi.org/10.1055/s-0042-1750341>.  
ISSN 1450-1147.


© 2022. World Association of Radiopharmaceutical and Molecular Therapy (WARMTH). All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

findings of COVID-19 (<sup>18</sup>F-FDG PET) in the diagnosis of COVID-19.<sup>4-6</sup> The most prominent features reported on FDG-PET/CT in patients with COVID-19 include FDG uptake in pulmonary parenchymal lesions that are typically peripheral, bilateral, ground glass opacities.<sup>5</sup>

In the detection and characterization of neuroendocrine tumors (NETs), the NETSPOT <sup>68</sup>Ga-DOTATATE PET/CT utilizes a somatostatin analog to capture whole body imaging of somatostatin receptors. Somatostatin receptor imaging is successfully accomplished using <sup>111</sup>In-DTPA-pentetreotide (Octreoscan).<sup>7</sup> However, downsides to Octreoscan (including suboptimal spatial resolution, suboptimal image quality, and multiple day imaging protocols) have led to the development of several chelator-conjugated somatostatin analogs, including DOTATATE. These short amino acid-chelator conjugates have a higher affinity than Octreoscan for somatostatin receptors. Further, they can be labeled with <sup>68</sup>Ga, permitting imaging with PET and improved spatial resolution and image quality.<sup>7</sup> Here, to our knowledge, we report the first case of incidental COVID-19 detection using <sup>68</sup>Ga-DOTATATE PET/CT for a patient with COVID-19 who presented to our nuclear medicine department for oncologic and neuroendocrine indications.




**Fig. 1** (A) Axial computed tomography (CT) with lung windows demonstrates peripheral subpleural ground glass opacities predominantly in the lower lobes, typical of coronavirus disease 2019 pneumonia. These infiltrates correspond to the areas of uptake on scintigraphy. (B) Axial positron emission tomography-computed tomography (PET-CT) fusion with <sup>68</sup>Ga-DOTATATE PET/CT shows radiotracer uptake corresponding to the anatomic infiltrates. Maximum standardized uptake value 2.6.

## Case Report

In July of 2020, a 48-year-old male with known pancreatic neuroendocrine carcinoma was evaluated for status of disease with NETSPOT <sup>68</sup>Ga PET/CT. The scan demonstrated multiple areas of metastatic disease in the liver and abdominal lymph nodes (not shown) and new bilateral ground glass opacities in the lungs (►Fig. 1A and B). The developing infiltrates were associated with low level but clearly abnormal radiotracer activity (►Fig. 2). Upon questioning the patient, it was discovered that he was recovering from COVID-19 infection that was diagnosed 3 weeks prior (►Fig. 3). The findings were not present on a prior scan



**Fig. 2** Maximal intensity projection image from the positron emission tomography-computed tomography demonstrates numerous hypermetabolic foci in the liver and upper abdominal lymph nodes, consistent with somatostatin-avid pancreatic islet cell neoplasm. Maximum standardized uptake value (SUV<sub>max</sub>) liver masses 5.2. SUV<sub>max</sub> abdominal lymph nodes 4.7.



**Fig. 3** Coronal positron emission tomography-computed tomography fusion shows radiotracer accumulation corresponding to the distribution of the bilateral ground glass pneumonia.

3 months earlier (not shown). Subsequent scan several months later for standard treatment follow-up demonstrated resolution of infiltrates and associated radiotracer uptake (not shown).

## Discussion

The increase in reported incidental findings of COVID-19 on FDG-PET/CT imaging underscores the impact nuclear medicine may have outside of the oncologic setting. FDG-PET/CT plays a role in evaluating various inflammatory and infectious diseases, which may guide patient management and treatment.<sup>8</sup> Before the appearance of COVID-19, Das et al reported significant FDG uptake in a patient with Middle East respiratory syndrome coronavirus infection that ultimately progressed to pneumonia.<sup>9</sup>

In our case, an asymptomatic COVID-19 patient presented to our clinic for neuroendocrine cancer screening, and we observed pathologic uptake within the pneumonia using <sup>68</sup>Ga-DOTATATE PET/CT. It is important that the clinician be aware of potential false positives when imaging with this radiotracer.

Current documented false-positive findings for <sup>68</sup>Ga-DOTATATE PET/CT include pancreatic uncinuate process activity, prostatitis, splenosis, postradiation therapy change, paraganglioma, pheochromocytoma, neuroblastoma, meningioma, and osteoblastic activity.<sup>10</sup> This case adds COVID-19 pneumonia to the list. With the growing number of worldwide COVID-19 cases, and the continued use of NETSPOT for somatostatin avid neoplasms, it is imperative

that diagnostic measures of this disease are clearly understood. This case highlights the importance of including COVID-19 on the differential diagnosis when radiotracer uptake appears in <sup>68</sup>Ga-DOTATATE PET/CT.

## Authors' Contributions

JPW was involved in conceptualization, designing, definition of intellectual content, literature search, manuscript preparation, review, and editing. TB contributed substantially in conceptualization, designing, definition of intellectual content, manuscript review and editing. TB has provided guarantee for this manuscript.

## Funding

No financial information to disclose.

## Conflicts of Interest

None declared.

## References

- 1 CDC COVID Data Tracker. <https://covid.cdc.gov/covid-data-tracker/#datatracker-home>. Accessed April 3, 2022
- 2 Cascella M, Rajnik M, Cuomo A, Dulebohn SC, Di Napoli R. Features, Evaluation, and Treatment of Coronavirus (COVID-19). In: StatPearls. Treasure Island (FL): StatPearls Publishing
- 3 Fang Y, Zhang H, Xie J, et al. Sensitivity of chest CT for COVID-19: comparison to RT-PCR. *Radiology* 2020;296(02):E115–E117
- 4 Albano D, Bertagna F, Bertoli M, et al. Incidental findings suggestive of COVID-19 in asymptomatic patients undergoing nuclear medicine procedures in a high-prevalence region. *J Nucl Med* 2020;61(05):632–636
- 5 Qin C, Liu F, Yen T-C, Lan X. <sup>18</sup>F-FDG PET/CT findings of COVID-19: a series of four highly suspected cases. *Eur J Nucl Med Mol Imaging* 2020;47(05):1281–1286
- 6 Lütje S, Marinova M, Kütting D, Attenberger U, Essler M, Bundschuh RA. Nuclear medicine in SARS-CoV-2 pandemic: <sup>18</sup>F-FDG-PET/CT to visualize COVID-19. *Nucl Med (Stuttg)* 2020;59(03):276–280
- 7 Barrio M, Czernin J, Fanti S, et al. The impact of somatostatin receptor-directed PET/CT on the management of patients with neuroendocrine tumor: a systematic review and meta-analysis. *J Nucl Med* 2017;58(05):756–761
- 8 Deng Y, Lei L, Chen Y, Zhang W. The potential added value of FDG PET/CT for COVID-19 pneumonia. *Eur J Nucl Med Mol Imaging* 2020;47(07):1634–1635
- 9 Das KM, Lee EY, Langer RD, Larsson SG. Middle east respiratory syndrome coronavirus: what does a radiologist need to know? *AJR Am J Roentgenol* 2016;206(06):1193–1201
- 10 Hofman MS, Lau WFE, Hicks RJ. Somatostatin receptor imaging with <sup>68</sup>Ga DOTATATE PET/CT: clinical utility, normal patterns, pearls, and pitfalls in interpretation. *Radiographics* 2015;35(02):500–516