







C. W. LAMARTINA, C. A. CHARTIER, S. LEE, N. H. SHAH*, T. ROVIS* (COLUMBIA UNIVERSITY, NEW YORK, USA)

Modular Synthesis of Unnatural Peptides via Rh(III)-Catalyzed Diastereoselective Three-Component Carboamidation Reaction
J. Am. Chem. Soc. **2023**, *145*, 1129–1135, DOI: 10.1021/jacs.2c10793.

Rh(III)-Catalyzed Synthesis of Unnatural Peptides



All yields and dr are isolated, with the crude dr in parentheses.

Significance: Introducing unnatural amino acid residues into the peptide chains is significant in the development of novel drugs. The authors have developed a carboamidation method to insert an unnatural amino acid residue into the peptides from dioxazolones, arylboronic acids and acrylamides.

Comment: A series of peptides containing unnatural amino acid residues are prepared. The yields of these three-component reactions are moderate to good by using Rh(III) as catalyst.

Category

Peptide Chemistry

Key words

rhodium catalysis

unnatural peptides

three-component reaction

carboamidation

dioxazolones

arylboronic acids

acrylamides

Synfact
of the
Month

SYNFACTS Contributors: Hisashi Yamamoto, An Wu
Synfacts 2023, 19(04), 0417 Published online: 17.03.2023
DOI: 10.1055/s-0042-1752646; Reg-No.: H03323SF

© 2023, Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.