


J. YIN, P. H. JACOBSE, D. PYLE, Z. WANG, M. F. CROMMIE*, G. DONG* (UNIVERSITY OF CALIFORNIA AT BERKELEY, LAWRENCE BERKELEY NATIONAL LABORATORY AND THE UNIVERSITY OF CHICAGO, USA)

Programmable Fabrication of Monodisperse Graphene Nanoribbons via Deterministic Iterative Synthesis
J. Am. Chem. Soc. **2022**, *35*, 16012–16019, DOI: 10.1021/jacs.2c05670.

Modular Syntheses of Defined Graphene Nanoribbons



Significance: The accomplishment of bottom-up syntheses of graphene nanoribbons (GNRs) with controlled, well-defined structures will open up a new stage for graphene functionalization, fabrication, and application. Here, by introducing 1,8-diaminonaphthalene (**dan**) as a handy protecting group for the boronic acid to attain iterative site-selective Suzuki–Miyaura cross-coupling (SMC), programmable fabrication of sequence-controlled GNRs is achieved.

Comment: The developed method, so called protecting-group-aided iterative synthesis (PAIS), is demonstrated to be suitable for the construction of tailored, monodisperse GNRs. The masking group **dan** is the key to the precise sequence control, which is readily transformed into reactive boronic acid upon exposure to acid and enables SMC for subsequent chain extension. The final GNRs are eventually obtained upon cyclodehydrogenation, either in solution or on the surface.

SYNFACTS Contributors: Dahui Zhao, Yiming Liu
Synfacts 2022, 18(12), 1299 Published online: 17.11.2022
DOI: 10.1055/s-0042-1753075; Reg-No.: S10922SF

© 2022, Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Synthesis of
Materials and
Unnatural Products

Key words

graphene
nanoribbons
cross-coupling

Synfact
of the
Month