

Use of the SARC-F Score as an Aid in Fragility Fractures Prevention^{*} O uso do escore SARC-F como auxiliar na prevenção de fraturas por fragilidade

Carlos Augusto Nunes Martini^{1®} Carolina Souza Weigert^{1®} Anderson Carlos Bigolin Stiegemaier^{1®} Ana Paula Ribeiro Bonilauri Ferreira^{2®} Ellen Liceras Gonçalves^{2®} Sandro Fortes Valle^{3®}

¹Institute of Orthopedics and Traumatology, Joinville, SC, Brazil

² Scientific Department, Institute of Orthopedics and Traumatology, Joinville, SC, Brazil Address for correspondence Carlos Augusto Nunes Martini, Orthopedic Surgeon IOT/Hospital Municipal São José – Joinville, SC, Brazil (e-mail: carlosnunesmartini@gmail.com).

³Medical School, Universidade da Região de Joinville (UNIVILLE), Joinville, SC, Brazil

Rev Bras Ortop 2023;58(1):157–163.

Abstract	 Objective The present study aimed to relate the strength, assistance with walking, rising from a chair, climbing stairs, and falls (SARCF) score with the presence or absence of fragility fracture in the population over 60 years of age. Methods The risk of sarcopenia was determined through the application of the SARC-F questionnaire, and the patients were divided into 2 groups, according to the
Keywords	occurrence or not of fragility fracture ($n = 100$).
 risk factors 	Results Thirty-two cases of distal radius fractures and eighteen cases of proximal
 osteoporotic fractures osteogenesis imperfecta 	femur fractures were identified. A higher score on the SARC-F is determinant between having or not a fragility fracture, estimating that for each point in the score there is a 70% increase in the chance of a patient having a fracture, regardless of age, gender, and body mass index (BMI).
osteoporosissarcopenia	Conclusion There was a direct correlation between a higher score on the SARC-F and an increase in the chance of fragility fracture.
Resumo	Objetivo O presente estudo teve como objetivo relacionar o escore <i>strength</i> , <i>assistance with walking</i> , <i>rising from a chair</i> , <i>climbing stairs</i> , <i>and falls</i> (SARC-F) com a presença ou não de fratura por fragilidade na população acima de 60 anos.

* Work developed at the Institute of Orthopedics and Traumatology of Hospital Municipal São José, Joinville, SC, Brazil.

received January 26, 2022 accepted after revision July 26, 2022 article published online October 18, 2022 DOI https://doi.org/ 10.1055/s-0042-1756328. ISSN 0102-3616. © 2022. Sociedade Brasileira de Ortopedia e Traumatologia. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (https://creativecommons.org/licenses/by-ncnd/4.0/)

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

Métodos O risco de sarcopenia foi determinado por meio da aplicação do questionão de fratura por fragilidade (n = 100).

- **Palavras-chave**
- fatores de risco
- ► fraturas por osteoporose
- osteogênese imperfeita
- osteoporose
- sarcopenia

- nário SARC-F, sendo os pacientes divididos em 2 grupos, de acordo com a ocorrência ou Resultados Foram levantados 32 casos de fratura de rádio distal e 18 casos de fratura
- de fêmur proximal. Uma maior pontuação no SARC-F determina bem entre ter ou não ter fratura por fragilidade, estimando que a cada ponto a mais no escore há um acréscimo de 70% na chance de o paciente ter fratura, independentemente da idade, sexo e índice de massa corporal (IMC).
- **Conclusão** Houve correlação direta entre uma maior pontuação no SARC-F e aumento na chance de fratura por fragilidade.

Introduction

Fragility fractures are characterized by injuries resulting from low-energy trauma, usually a fall from the person's height, which possibly would not result in fracture in the case of a healthy bone structure.¹ In the Western world, approximately 1 in 3 women and 1 in 5 men over 50 years of age will suffer a fracture in their remaining lifetime.² These fractures occur more commonly in the hip, spine, and wrist, affecting mainly the elderly population, due to the higher incidence of osteoporosis.³ Such fractures can lead to serious complications, decreasing the patient's quality of life, as well as being life-threatening in some cases. In addition, the cost to the health system of treating these fractures is high, since the chance of complications is considerable, and, in some hospitalization and surgical treatment are cases, necessary.4,5

In addition to osteoporosis, more recently, sarcopenia has been correlated with an increased likelihood of an individual suffering from fragility fracture.⁶ Sarcopenia is defined by a decrease in muscle mass, leading to decreased performance capacity of daily activities and increased probability of unfavorable outcomes such as falls, fractures, and death.⁷ The European Working Group on Sarcopenia in Older People (EWGSOP2) describes the strength, assistance with walking, rising from a chair, climbing stairs, and falls (SARC-F) score as a screening for sarcopenia, being a low-cost tool, moderate sensitivity, and high specificity to predict low muscle mass in the studied population.^{6,8–10} This is a useful tool for use in clinical practice to prevent the occurrence of fragility fracture.

The aim of this study was to relate the SARC-F score of patients with the presence or not of fragility fracture.

Materials and Methods

This was an observational, analytical study, approved by the ethics committee, under the opinion of number 4,463,378 -CAAE- 39467220.5.0000.5362.

Patients over 60 years of age, treated at the outpatient clinic of our hospital, from January to June 2021, who presented fragility fracture due to a fall from their own height were included in the present study; or who had never suffered any fragility fracture; and who

agreed to participate in the study and signed the consent form.

Patients whose fractures were due to high-energy trauma were excluded from the study, as well as patients with alteration of cognition or memory that made them unable to answer the questionnaire, patients with deformities or motor limitations in the lower limbs prior to fracture, and fragility fractures that occurred more than 6 months ago.

Patients were allocated to two groups of 50 patients each:

- a) Control group: patients who have never suffered fragility fractures;
- b) Fracture group: patients who suffered fragility fractures.

Intervention

In addition to the demographic variables, information was collected regarding the patients' body mass index, fracture date, whether there were previous treatments for osteoporosis or sarcopenia, and the presence of absence of smoking.

The SARC-F score was the main study data collection tool used to identify patients at high risk of sarcopenia. The SARC-F assesses muscle strength, the need for walking assistance, the ability to get up from a chair, climb stairs, and the frequency of falls. Each item can be scored from 0 to 2, 0 being no difficulty, 1 some difficulty, and 2 very difficult or unable to perform. In the item falls, 0 corresponds to no falls, 1 corresponds from 1 to 3 falls, and 2 corresponds to 4 or more falls. The final score can range from 0 to 10, and the \geq 4 score is considered predictive of sarcopenia (**~Table 1**).

The patients in the control group answered the SARC-F questionnaire when they underwent outpatient consultation due to other orthopedic pathologies (gonarthrosis, coxarthrosis, chronic shoulder injuries, phalanx fracture, etc.). The patients of the *fracture group* answered the SARC-F questionnaire at the time of the first outpatient return consultation, being instructed to give the answers according to their functional status immediately before suffering the fracture. All questionnaires were personally applied by the researchers.

Statistical analysis

The results obtained in the study were described by mean, standard deviation, minimum and maximum (quantitative

COMPONENTS	QUESTIONS	RESULTS
Strength	What is your difficulty in lifting or carrying 4 kg?	None = 0 Some = 1 A lot or incapable = 2
Walking assistance	What is your difficulty in walking through a room?	None = 0 Some = 1 A lot, with help or incapable = 2
Chair lifting	How hard is it to get out of bed or chair?	None = 0 Some = 1 Too much or unable without $help = 2$
Climb stairs	What's your difficulty in climbing 10 steps?	None = 0 Some = 1 A lot or incapable = 2
Falls	How many times have you fallen in the last year?	None = 0 1 to 3 falls = 1 4 or more falls = 2

Table 1 SARC-F score

Abbreviations: SARCF, strength, assistance with walking, rising from a chair, climbing stairs, and falls.

variables) or by frequencies and percentages (categorical variables). For age, the 2 cutoff points, 65 and 73 years, were considered.

For SARC-F, classifications 0 or > 0 and classifications 0, 1 or > 1 were considered.

For the components of the SARC-F classifications, 1 and 2 were grouped, that is, the classifications *none* or *any/very or incapable*). The estimated association measure was the odds ratio (OR) for which a 95% confidence interval was presented. Values of p < 0.05 indicated statistical significance. The data were analyzed with the Stata/SE v.14.1. software (StataCorp LLC, College Station, TX, USA).

Results

Of the 100 patients evaluated, the mean age was 70.1 years, and 58 of these patients were female (58%). Of the 50 patients who suffered fragility fractures, distal radius fracture was the most prevalent (64%), with the other being proximal femur fractures (36%) (\succ Table 2).

Table 2 Descriptive analysis of demographic and localcharacteristics of fractures

Features				
Gender	er Male			
	Female	58		
Age	≤ 65	33		
	> 65	67		
Fragility fracture	Yes	50		
	No	50		
Fracture site	Distal radius	32		
	Proximal femur	18		

Presence of Fragility Fracture versus SARC-F Score

A statistically significant difference was observed between the presence of fragility fracture and the SARC-F score (p < 0.001), that is, patients who suffered fragility fractures had a higher SARC-F score (**\leftarrowTable 3**).

A receiver operating characteristic (ROC) curve was adjusted for the SARC-F score considering the presence or absence of fracture. The area below the curve corresponded to 0.83 presenting statistical significance (p < 0.001), indicating that the SARC-F score discriminates the presence or not of fracture (**-Fig. 1**).

The cut-off point indicated by the ROC curve adjustment was equal to 0. SARC-F equal to 0 is associated with absence of fracture, and SARC-F > 0 was associated with the presence of fracture. The sensitivity of this cutoff point is equal to 90% and specificity is equal to 64%.

Presence of Fragility Fracture versus Patient Age

There was a statistically significant difference between the presence of fragility fracture and the age of the patients (p < 0.001), that is, patients who suffered fragility fractures had the highest mean age (**\succ Table 4**).

A ROC curve was adjusted for the age of the patients considering the presence or not of fracture. The area below the curve corresponded to 0.79 presenting statistical significance (p < 0.001), indicating that the age of the patients discriminates the presence or not of fracture (**-Fig. 2**).

Tab	le 3	Presence o	ft	fragility	fracture	versus	SARC-F	score
-----	------	------------	----	-----------	----------	--------	--------	-------

Presence of fracture	N	SARC-F average score	P* value
No	50	0.66	< 0.001
Yes	50	2.86	

Abbreviations: *Student t-test for independent samples, considering p < 0.05.



Fig. 1 Receiver operating characteristics curve for the SARC-F score considering the presence or absence of fracture.

Table 4 Presence of fragility fracture versus age of patients

Presence	Ν	Age (years)	p *
of fracture		Mean (standard deviation)	
No	50	66.4 (±4.3)	< 0.001
Yes	50	73.8 (±7.5)	

Abbreviation: SARC-F, strength, assistance with walking, rising from a chair, climbing stairs, and falls.

*Student t test for independent samples, considering p < 0.05.



Fig. 2 Receiver operating characteristics curve for the age of patients considering the presence or absence of fracture.

The cutoff point indicated by the ROC curve adjustment was 73 years of age. Age \leq 73 years is associated with absence of fracture and age > 73 years is associated with the presence of fracture. The sensitivity of this cutoff point is equal to 56% and the specificity is 96%. If the cutoff point is equal to 65 years, the sensitivity is equal to 82% and the specificity is equal to 48%.

Association of Demographic Variables and Clinical Variables with the SARC-F Score

According to **- Table 5**, of the variables analyzed, age, gender and treatment for osteoporosis presented statistical significance. That is, the older the patient, the higher the SARC-F

 Table 5
 Association of demographic variables and clinical variables with the SARC-F score

Variables	Category	N	SARC-F score	p *
			Average	
Age (years)	≤ 65	33	1.03	0.020
	> 65	6 7	2.12	
Gender	Female	58	2.29	0.002
	Male	42	1.02	
BMI (kg/m ²)	< 25	28	2.36	0.126
	≥ 25	72	1.53	
Treatment for osteoporosis	No	71	1.20	0.001
	Yes	29	3.14	
Smoking	No	51	1.94	0.105
	Yes / Ex-smoker	49	1.57	

Abbreviation: SARC-F, strength, assistance with walking, rising from a chair, climbing stairs, and falls. *Mann-Whitney nonparametric test, considering p < 0.05.

score (p = 0.020); women had a higher SARC-F score (p = 0.02); and patients undergoing treatment for osteoporosis had a higher SARC-F score (p = 0.001).

Evaluation of Fracture-Related Factors

For categorical variables, the percentages were calculated in relation to the totals in the rows (column n), that is, they add up to 100% in each row (**- Table 6**).

Age was an important factor in the occurrence of fractures, especially in the age group above 73 years, since these patients were about 30 times more likely to suffer fractures than younger patients. Female patients were 8 times more likely to suffer fractures (OR = 8.84). Considering body mass index (BMI) as an isolated variable, individuals with BMI below 25 kg/m² had a nearly three-fold higher chance of suffering fragility fracture. Smokers or former smokers had a lower rate of fractures in relation to the population that never smoked, as well as those who never underwent treatment for osteoporosis.

As for the variables of the SARC-F, patients who scored 0, that is, no difficulty to carry a weight of 4 kg, get up from the chair, climb 10 steps, and no fall in the last year, had a significant reduction in the risk of fragility fractures. Individuals who scored > 0 in the SARC-F were 16 times more likely to suffer fragility fractures than those with SARC-F equal to 0.

Multivariate Analysis

To evaluate the effect of SARC-F on the probability of having a fragility fracture, a multivariate model was

Variable	Category	Ν	Presence of fracture		P* value	OR (95%CI)
			No	Yes		
Age (years)	< 73 (ref)	70	48 (68.6%)	22 (31.4%)	< 0.001	30.6 (6.68–140)
(ROC curve)	≥ 73	30	2 (6.7%)	28 (93.3%)		
Age (years)	< 65 (ref)	33	24 (72.7%)	9 (27.3%)	0.002	4.21 (1.69–10.4)
	≥ 65	67	26 (38.8%)	41 (61.2%)		
Gender	Male (ref)	42	33 (78.6%)	9 (21.4%)	<0.001	8.84 (3.49–22.4)
	Female	58	17 (29.3%)	41 (70.7%)		
BMI (kg/m ²)	≥ 25 (ref)	72	41 (56.9%)	31 (43.1%)	0.029	2.79 (1.11–7.01)
	<25	28	9 (32.1%)	19 (67.9%)		
Treatments for osteoporosis	No (ref)	71	43 (60.6%)	28 (39.4%)	0.002	4.83 (1.82–12.8)
	Yes	29	7 (24.1%)	22 (75.9%)		
Smoking	No	51	20 (39.2%)	31 (60.8%)	0.029	2.45 (1.09-5.47)
	Yes/Ex (ref)	49	30 (61.2%)	19 (38.8%)		
Strength	0 (ref)	84	47 (56%)	37 (44.1%)	0.012	5.51 (1.46-20.8)
	1 or 2	16	3 (18.8%)	13 (81.3%)		
Walking assistance	0 (ref)	85	46 (54.1%)	39 (45.9%)	0.059	3.24 (0.96–11.0)
	1 or 2	15	4 (26.7%)	11 (73.3%)		
Getting up from a chair	0 (ref)	78	46 (59%)	32 (41%)	0.002	6.47 (2.00-20.9)
	1 or 2	22	4 (18.2%)	18 (81.8%)		
Climbing stairs	0 (ref)	74	48 (64.9%)	26 (35.1%)	< 0.001	22.1 (4.85–101)
	1 or 2	26	2 (7.7%)	24 (92.3%)		
Falls	0 (ref)	42	35 (83.3%)	7 (16.7%)	< 0.001	14.3 (5.26–39.0)
	1 or 2	58	15 (25.9%)	43 (74.1%)		
SARC-F	0 (ref)	37	32 (86.5%)	5 (13.5%)	< 0.001	16 (5.38–47.6)
	> 0	63	18 (28.6%)	45 (71.4%)		
SARC-F	0 (ref)	37	32 (86.5%)	5 (13.5%)		
	1	29	13 (44.8%)	16 (55.2%)	0.001	7.88 (2.39–26.0)
	> 1	34	5 (14.7%)	29 (85.3%)	0.011	37.1 (9.74–141)

 Table 6
 Univariate analysis

Abbreviations: 95%CI, 95% confidence interval; Ex, former smoker; OR, odds ratio; Ref, reference; ROC, receiver operating characteristics; SARC-F, strength, assistance with walking, rising from a chair, climbing stairs, and falls. Legend: *Logistic regression model and Wald test considering p < 0.05.

Variable	Category	<i>p</i> *	OR (95%CI)
SARC-F		0.008	1.70 (1.15–2.51)
Age (years)		0.007	1.17 (1.04–1.32)
Gender	Male (ref)	< 0.001	7.91 (2.48–25.2)
	Female		
BMI (kg/m ²)	≥ 25 (ref)	0.183	2.42 (0.66-8.90)
	< 25		

Tab	le	7	Mu	ltiva	iriate	ana	lysis
-----	----	---	----	-------	--------	-----	-------

Abbreviation: BMI, body mass index; 95%CI, 95% confidence interval; OR, odds ratio; Ref, reference; SARCF, strength, assistance with walking, rising from a chair, climbing stairs, and falls.

*Logistic regression model and Wald test considering p < 0.05.

adjusted considering the fracture (yes or no) as a dependent variable and including as explanatory variables those that presented statistical significance in the univariate analysis (**-Table 7**).

The results of the multivariate model adjustment indicate that regardless of age, gender, and BMI, a higher SARC-F score is significantly associated with the probability of having fragility fracture. It is estimated that for each additional point in the SARC-F there is an increase of 70% in the chance of having fracture, regardless of age, gender, and BMI. It can also be said that regardless of SARC-F, gender, and BMI, with each additional year of age there is a 17% increase in the chance of having a fracture. Female patients, regardless of SARC-F, age, and BMI, have 7.91 times more chance of fracture than a male patient. Body mass index, in the presence of the variables SARC-F, age, and gender, is not significantly associated with the probability of having a fracture.

Discussion

Up to our current knowledge, this is the first study that directly comcants the SARC-F with the occurrence of fragility fractures, in view of a possible correlation between sarcopenia and increased risk of falls.

The most important finding of this study was that a higher SARC-F score leads to a significant increase in the risk of fragility fractures. Another finding, no less relevant, however already established in the literature, was that age interferes forcefully in the fracture index.^{5,11}

According to the data presented in this study, age equal to or above 65 years of age presented an important correlation with the fracture index (sensitivity 82%, specificity 48%), and each year older, the individual's chance of suffering a fracture increases by 17%. It is a fact that the older the patient, the higher the incidence of osteoporosis, frailty, and consequently, the higher the risk of pathological fractures.^{5,12,13}

According to Borgström, the female population suffers more fragility fractures, at a ratio of 2:1.² In the present study, it was observed that female patients were 7.91 times more likely to suffer a fracture compared to males. This fact may be related to menopause, low calcium intake, and lack of physical activity.^{11,14}

The SARC-F is considered an important tool in the diagnosis of sarcopenia. The findings in this study showed

that the SARC-F discriminated well between the occurrence or not of fracture, estimating that at each point more in the SARC-F, there was an increase of 70% in the chance of the patient having suffered a fracture, regardless of age, gender, and BMI. However, previous studies have not sought to establish a direct correlation between a higher result in the SARC-F score and the increased chance of the individual suffering fragility fracture, keeping the focus on the application questionnaire test as screening of sarcopenia.^{9,15}

Of the items of the SARC-F score evaluated individually, it was observed that the items *difficulty to climb stairs* and *previous episodes of fall in the last year* were the ones most strongly associated with the occurrence of fractures. The presence of sarcopenia may increase twofold the chance of falls compared to non-sarcopenic individuals.¹⁶

In the present study, only 29 of the 100 patients evaluated had undergone some type of previous treatment for osteoporosis and sarcopenia, and most of them, that is, 22 patients, suffered some fragility fracture, even though they had undergone some treatment for this purpose previously. This finding may be attributed to incomplete or inadequate treatment for osteoporosis, as well as underdiagnosis of the disease. The low rate of patients who underwent some type of treatment for the prevention of fragility fractures is relevant; however, this finding is already reported in the literature, in which, especially in individuals over 75 years of age, osteoporosis is treated inadequately or is simply not treated.^{17,18}

Regarding the location of fractures, no vertebral fracture was identified in the analyzed population, with distal radius fracture being the most prevalent, with 32 cases, followed by proximal femur fracture, with 18 cases. In a previous study, the site most frequently associated with osteoporotic fractures was the proximal femur, followed by fractures of the spine and only then the distal radius.¹⁹

In the present study, it was not possible to establish a correlation between BMI and the probability of fracture occurrence. The vast majority of individuals (n = 72) were overweight (BMI > 25 kg/m²), and only a single patient with BMI < 18.5 kg/m² suffered distal radius fracture. According to Court-Brown et al.,²⁰ there is a positive correlation between underweight patients (BMI < 18.5 kg/m²) and proximal femur fracture due to a higher incidence of sarcopenia in

these individuals, while the authors did not find a correlation between obesity and a higher risk of fractures.

The limitation of this study was the fact that the studied population belonged to the same institution, creating a homogeneous sample of the same socioeconomic profile and even similar life habits. The inclusion of participants from other institutions with different profiles could contribute to a better elucidation of the occurrence of fragility fractures in the general population.

Tools that can assist the physician in the identification of individuals who have a high potential to suffer fragility fracture, are certainly useful in clinical practice.³ Through this research, it was possible to determine that the SARC-F presents a direct correlation with the risk of fractures, demonstrating that sarcopenia is a diagnosis that should be remembered by the physician in the care of elderly patients who have suffered fractures. Scores can guide the physician in requesting tests, prescribing medications, or referral to other specialties, but decision-making should be individualized and based on several factors, not only supported by calculations.

The development of a multicenter study with a more comprehensive age group of patients would be the suggested next step for future research for a solid finding of the relationship between SARC-F and the risk of fragility fractures.^{21,22}

Conclusion

Our findings showed that there is a direct correlation between a higher SARC-F score and an increase in the chance of fragility fracture, especially in individuals over 65 years of age.

Financial Support

There was no financial support from public, commercial, or non-profit sources.

Conflict of Interests

The authors declare that there is no conflict of interests.

References

- 1 Shah GM, Gong HS, Chae YJ, Kim YS, Kim J, Baek GH. Evaluation and Management of Osteoporosis and Sarcopenia in Patients with Distal Radius Fractures. Clin Orthop Surg 2020;12(01):9–21
- 2 Borgström F, Karlsson L, Ortsäter G, et al; International Osteoporosis Foundation. Fragility fractures in Europe: burden, management and opportunities. Arch Osteoporos 2020;15(01):59
- 3 Zerbini CAF, Albergaria BH. The brazilian FRAX model: an introduction. Rev Assoc Med Bras (1992) 2018;64(06):481–483
- 4 Wu Q, Xiao X, Xu Y. Performance of FRAX in Predicting Fractures in US Postmenopausal Women with Varied Race and Genetic Profiles. J Clin Med 2020;9(01):285
- 5 Li G, Thabane L, Papaioannou A, Ioannidis G, Levine MA, Adachi JD. An overview of osteoporosis and frailty in the elderly. BMC Musculoskelet Disord 2017;18(01):46

- 6 Wong RMY, Wong H, Zhang N, et al. The relationship between sarcopenia and fragility fracture-a systematic review. Osteoporos Int 2019;30(03):541–553
- 7 Cruz-Jentoft AJ, Baeyens JP, Bauer JM, et al; European Working Group on Sarcopenia in Older People. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. Age Ageing 2010; 39(04):412–423
- 8 Woo J, Leung J, Morley JE. Validating the SARC-F: a suitable community screening tool for sarcopenia? J Am Med Dir Assoc 2014;15(09):630–634
- 9 Ha YC, Won Won C, Kim M, Chun KJ, Yoo JI. SARC-F as a Useful Tool for Screening Sarcopenia in Elderly Patients with Hip Fractures. J Nutr Health Aging 2020;24(01):78–82
- 10 Wu TY, Liaw CK, Chen FC, Kuo KL, Chie WC, Yang RS. Sarcopenia Screened With SARC-F Questionnaire Is Associated With Quality of Life and 4-Year Mortality. J Am Med Dir Assoc 2016;17(12): 1129–1135
- 11 Blalock SJ, DeVellis RF, Giorgino KB, et al. Osteoporosis prevention in premenopausal women: using a stage model approach to examine the predictors of behavior. Health Psychol 1996;15 (02):84–93
- 12 Kirk B, Phu S, Brennan-Olsen SL, Bani Hassan E, Duque G. Associations between osteoporosis, the severity of sarcopenia and fragility fractures in community-dwelling older adults. Eur Geriatr Med 2020;11(03):443–450
- 13 Balogun S, Winzenberg T, Wills K, et al. Prospective associations of osteosarcopenia and osteodynapenia with incident fracture and mortality over 10 years in community-dwelling older adults. Arch Gerontol Geriatr 2019;82:67–73
- 14 Winzenberg TM, Oldenburg B, Frendin S, Jones G. The design of a valid and reliable questionnaire to measure osteoporosis knowledge in women: the Osteoporosis Knowledge Assessment Tool (OKAT). BMC Musculoskelet Disord 2003;4:17
- 15 Barbosa-Silva TG, Menezes AM, Bielemann RM, Malmstrom TK, Gonzalez MCGrupo de Estudos em Composição Corporal e Nutrição (COCONUT) Enhancing SARC-F: Improving Sarcopenia Screening in the Clinical Practice. J Am Med Dir Assoc 2016;17(12):1136–1141
- 16 Scott D, Seibel M, Cumming R, et al. Does Combined Osteopenia/Osteoporosis and Sarcopenia Confer Greater Risk of Falls and Fracture Than Either Condition Alone in Older Men? The Concord Health and Ageing in Men Project. J Gerontol A Biol Sci Med Sci 2019;74(06):827–834
- 17 Coughlan T, Dockery F. Osteoporosis and fracture risk in older people. Clin Med (Lond) 2014;14(02):187–191
- 18 Vandenbroucke A, Luyten FP, Flamaing J, Gielen E. Pharmacological treatment of osteoporosis in the oldest old. Clin Interv Aging 2017;12:1065–1077
- 19 Warriner AH, Patkar NM, Curtis JR, et al. Which fractures are most attributable to osteoporosis? J Clin Epidemiol 2011;64(01): 46–53
- 20 Court-Brown CM, Duckworth AD, Ralston S, McQueen MM. The relationship between obesity and fractures. Injury 2019;50(08): 1423–1428
- 21 Rolland Y, Dupuy C, Abellan Van Kan G, et al. Sarcopenia Screened by the SARC-F Questionnaire and Physical Performances of Elderly Women: A Cross-Sectional Study. J Am Med Dir Assoc 2017;18 (10):848–852
- 22 Silva Parra BFC, Matos LBN, Ferrer R, Toledo DO. SARCPRO: Proposta de protocolo para sarcopenia em pacientes internados. BRASPEN J 2019;34(01):58–63