

# Recurrent Temporal Meningioma as Exophytic Mass from the External Auditory Canal

Sundarakrishnan Dharanipathy<sup>1</sup> Shweta Kedia<sup>1</sup>


<sup>1</sup>Department of Neurosurgery, All India Institute of Medical Sciences, New Delhi, India

Indian J Neurosurg

Address for correspondence Shweta Kedia, MBBS, MS, MCh, Department of Neurosurgery, All India Institute of Medical Sciences-110029, New Delhi, India (e-mail: drshwetakedia@gmail.com).

A 30-year-old female patient was diagnosed with right medial sphenoid wing meningioma when she was evaluated for her headache and dizziness. She underwent

craniotomy and excision and, biopsy showed grade 1 microcystic meningioma with an MIB-1 labeling index of 1 to 2%. Her follow-up magnetic resonance imaging (MRI) at



**Fig. 1** (A) Pre GKRS CEMRI showing recurrent tumor in Meckel's cave. (B) Two years of follow-up CEMRI showing extension of the tumor and protrusion into the middle ear and external auditory canal. (C) Preop patient image showing a lobular mass with surrounding purulent discharge. (D) Postop image showing planned incision. (E and F) Immediate postoperative CT and 3-month postoperative CEMRI, respectively, showing a near total excision of the lesion.

DOI <https://doi.org/10.1055/s-0042-1757204>.  
ISSN 2277-954X.

© 2022. Neurological Surgeons' Society of India. All rights reserved. This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)  
Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

2 years showed recurrence of the lesion and she was referred to our hospital for gamma knife radiosurgery (GKRS). On examination, she had 30 to 40% sensory loss along the distribution of the right trigeminal nerve. She had a 1.3 × 1.7 cm intensely enhancing globular extra-axial lesion along the right temporal convexity with extension to Meckel's cave and abutting the right trigeminal nerve (►Fig. 1). She underwent secondary GKRS with a dose of 12 Gray. She subsequently noticed a globular mass protruding out from her external auditory canal with associated purulent foul-smelling discharge and had conductive hearing loss (►Fig. 1). Computed tomography (CT) and Contrast enhanced magnetic resonance imaging (CEMRI) scan of the temporal bone showed a lobulated extra axial mass causing erosion of the petrous bone and tegmen tympani with extension to the middle ear, mastoid antrum, and external auditory canal. Intracranial meningiomas extending into the middle ear and subsequently into the external auditory canal are extremely rare. Kusunoki et al noted a case of recurrent sphenoid wing meningioma extending into the middle ear in a 74-year-old female patient.<sup>1</sup> Reitz et al suggested that intracranial meningiomas can enter the middle ear through various routes such as petrous pyramid, tegmen tympani, internal auditory canal, jugular fossa, and canal for greater superficial

petrosal nerve. In our case, the likely path of spread would be the tegmen and petrous pyramid.<sup>2</sup> She underwent preauricular subtemporal transzygomatic approach and near total excision of the lesion by the ENT department and biopsy proved to be meningioma again.

#### Funding

None.

#### Conflict of Interest

None declared.

#### Acknowledgments

We would like to thank Dr. Manmohan Singh, Dr. Shaswat Mishra, and Dr. Arvind Kumar Kairo.

#### References

- 1 Kusunoki T, Ikeda K, Miyashita M. A case report of meningioma extending to the middle ear. *Clin Pract* 2012;2(03):e67. Doi: 10.4081/cp.2012.e67
- 2 Rietz DR, Ford CN, Kurtycz DF, Brandenburg JH, Hafez GR. Significance of apparent intratympanic meningiomas. *Laryngoscope* 1983;93(11 Pt 1):1397-1404