

 $\bigcirc \textcircled{\bullet} \boxdot \textcircled{\bullet} \textcircled{\bullet} \textcircled{\bullet}$

Endoscopic Endonasal Transcavernous Transoculomotor Triangle Approach for the Resection of a Knosp 4 Pituitary Adenoma

Lijun Heng¹ Shuo Zhang¹ Lei Zhao¹ Yan Qu¹

¹Department of Neurosurgery, Tangdu Hospital, Air Force Medical University, Xi'an, China

J Neurol Surg B Skull Base 2022;83:e663-e664.

Address for correspondence Yan Qu, MD, PhD, Department of Neurosurgery, Tangdu Hospital, Air Force Medical University, 569 Xinsi Road, Xi'an, Shaanxi 710038, China (e-mail: yanqu0123@icloud.com).

Abstract	 Objectives The current video presents the nuances of the endoscopic endonasal transcavernous transoculomotor triangle approach for the resection of a Knosp 4 pituitary adenoma protruding through the superior wall of left cavernous sinus. Design The video analyzes the clinical presentation, preoperative workup, and imaging, surgical steps and technical nuances of the surgery, the clinical outcome, and follow-up imaging. Setting The patient was treated by a skull base team with bimanual, three- to four-handed endoscopic technique.
	Participants The patient was a 54-year-old male with a Knosp 4 prolactinoma presented with headaches and blurred vision in both eyes for 3 weeks. He was managed with endoscopic surgery due to his preference of surgery as a rapid primary therapy to relieve mental stress of an invasive brain tumor after awareness of all therapeutic options, including the preoperative medical treatment.
	Main Outcome Measures The main outcome was measured in the following aspects, including the resection extent, the symptoms improvement, the radiological and endo- crinological follow-up results, as well as the postoperative course of complications. Results Nearly gross total resection was achieved with one-stage endoscopic surgery.
Keywords ► endoscopic	recovered in 2 months. Endocrinological remission was kept for 12 months with one pill of bromocrintine daily
 endonasal cavernous sinus oculomotor triangle 	Conclusion The endoscopic endonasal transcavernous transoculomotor triangle approach is an appropriate option for resecting Knosp 4 pituitary tumors with intracranial extension through the roof of cavernous sinus at one stage.

oculomotor triangle
 pituitary adenoma



received March 17, 2020 accepted after revision August 31, 2022 article published online October 7, 2022 DOI https://doi.org/ 10.1055/s-0042-1757619. ISSN 2193-6331.

© 2022. The Author(s).

www.thieme.com/skullbasevideos

www.thieme.com/jnlsbvideos

The link to the video can be found at https://youtu.be/-YJ2PKP9voY.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (https://creativecommons.org/licenses/by-nc-nd/4.0/)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



Fig. 1 Patient's magnetic resonance imaging (MRI). Preoperative (A, D), 2 days postoperative (B, E), and 3 months postoperative (C, F) MRIs with coronal and sagittal views showing the removal of a Knosp 4 pituitary adenoma with left cavernous sinus invasion.



Fig. 2 Intraoperative image showing the superior and lateral exposure through oculomotor triangle and the essential neurovascular structures in this area during tumor resection.

Conflict of Interest None declared.