



Irrigation with Bupivacaine at the Surgical Bed for Postoperative Shoulder Tip and Abdominal Pain Relief after Laparoscopic Cholecystectomy

Affan Iqbal^{1,2} Amir Khodavirdipour³ HR Ravishankar¹

¹Department of General Surgery, Sagar Hospitals, Bangalore, Karnataka, India

²Department of General Surgery, Seventh-Day Adventist Hospital, Bangalore, Karnataka, India

³School of Sport and Health Sciences, University of Brighton, Brighton, United Kingdom.

Address for correspondence Affan Iqbal, MBBS, MS, Department of General Surgery, Seventh-Day Adventist Hospital, Bangalore, Karnataka, India (e-mail: draffaniqbal@gmail.com).

Surg J (NY) 2022;8:e312–e315.

Abstract

Gallstones in western countries are primarily composed of cholesterol. However, mixed or pigment stones, which contain a higher proportion of bilirubin, are more frequently seen in developing nations and Asia than in western countries. Abdominal and shoulder tip pains (STPs) are common complaints following the standard laparoscopic cholecystectomy procedure. To date, all pain management modalities have proven variable outcomes. This prospective randomized study included 82 patients who underwent elective laparoscopic cholecystectomy. The control group received 20 mL of normal saline, whereas the study group received a 20-mL instillation of 0.5% bupivacaine at the gallbladder bed after surgical resection. The Visual Analog Scale (VAS) was used to analyze abdominal pain and STP. The mean age ranged from 20 to 80 years. Abdominal VAS at 6, 12, 18, 24, 30, 36, and 48 hours were statistically insignificant. The majority were discharged on postoperative day 1 (32 studies, 37 control). Follow-up VAS after 1 week for STP VAS and abdominal pain VAS in both groups were statistically insignificant. Even with small numbers of a well-conducted randomized trial, we demonstrated that bupivacaine irrigation at the gallbladder bedpost laparoscopic cholecystectomy does not affect pain relief.

Keywords

- ▶ cholecystectomy
- ▶ gallstone
- ▶ bupivacaine
- ▶ laparoscopy

Introduction

Laparoscopic cholecystectomy (LC) is the gold standard in addressing benign gallbladder disease¹ and is one of the most widely performed surgeries in a general surgery department in India. The most common indications for LC as gallbladder stones are seldomly performed for carcinoma gallbladder.² LC is a minimally invasive surgery with reduced morbidity compared with its open counterpart; however, it has a few disturbing postoperative adverse effects, such as shoulder tip

pain (STP), abdominal pain, port site pain, infection, granuloma, and hernia.

Visceral pain is caused by intra-abdominal cavity content stretching and peritoneal inflammation. Furthermore, phrenic nerve irritation is caused by residual carbon dioxide. Postoperative abdominal pain develops within the first 24 hours, and STP usually manifests only on the second day.² Analgesics and opioids used in pain management have been reported to show variable results.¹ Pain is derived

received
January 30, 2021
accepted after revision
May 23, 2022

DOI <https://doi.org/10.1055/s-0042-1758141>.
ISSN 2378-5128.

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)
Thieme Medical Publishers, Inc., 333 Seventh Avenue, 18th Floor, New York, NY 10001, USA

Table 1 Sex group cross-tabulation

			Group		Total	p-Value
			Control	Study		
Sex	Female	Count	22	22	44	0.82
		% within group	55.0	52.4	53.7	
	Male	Count	18	20	38	
		% within group	45.0	47.6	46.3	
Total		Count	40	42	82	
		% within group	100.0	100.0	100.0	

Note: Chi-square test showing a nonsignificant p-value when correlated.

from various situations, namely, surgical site pain (somatic), deep visceral pain, and STP due to phrenic nerve irritation. Evidence suggests that in 17 to 41% of patients, the most probable cause of prolonged hospital stay is due to pain.¹ Postoperative pain is complex after laparoscopic surgery; hence, various specialties suggest an effective treatment to alleviate pain through multimodal support,¹ which includes desensitizing the sensory afferents (adequately injecting the skin with a local anesthetic [LA] before proceeding with an incision). Opioid administration preoperatively, a LA flushed into the peritoneal cavity, intravenous fluids, and electrolytes were adequately given.¹ Few studies showed that bupivacaine irrigation over the surgical bed proved effective in reducing pain in the first few hours after LC.¹⁻³ Whereas another study did not have such an effect.⁴

Abdominal pain and dyspepsia were the most common presenting symptoms occurring, either alone or together, in patients with gallbladder stones. Many other studies revealed these two symptoms as the most commonly reported symptoms among patients with cholelithiasis.⁵ Ursodeoxycholic acid, used for cholesterol stone diseases, has not been proven successful and is not usually applicable to all patients.⁶ Additionally, gallbladder elevations, called polyps, that protrude onto the lumen most of the time were observed with a higher gallbladder cancer rate in the Indian ethnicity shown in a study including 2,359 patients with gallbladder polyps (multivariate analysis: 5.5% vs. 0.08%).^{7,8} A study shows that intraperitoneal irrigation of the diaphragmatic surface and gallbladder fossa with bupiva-

caine/lignocaine and normal saline may effectively control visceral abdominal pain after LC.⁹ Nonsteroidal anti-inflammatory drugs, wound infiltration with LA, and intermittent intramuscular narcotics are various methods utilized for pain management with variable success after laparoscopic surgeries.¹⁰

Materials and Methods

This prospective randomized study was conducted in the Department of General Surgery, Sagar Hospitals, Bangalore, Karnataka, India over 18 months, which included 82 patients who were admitted and subsequently underwent elective standard four-port LC. The patients had their gallstones imaged by ultrasound before surgery ward admission and were divided randomly by a computer-generated table into two groups (42 studies and 40 control, ▶ **Table 1**). The control group received 20 mL of normal saline and the study group received a 20-mL instillation of 0.5% bupivacaine at the gallbladder bed after surgical resection. All the patients received pain relief as per the World Health Organization (WHO) pain ladder. Surgery was performed by a senior consultant surgeon with a resident as an assistant. Instillation was done via the lateral most port. Patients were postoperatively analyzed every 6 hours for STP and abdominal pain and were followed up until discharge. Pain assessment was done using the Wong-Bakers Visual Analog Scale (VAS) (▶ **Fig. 1**). Inclusion criteria are as follows: patients 18 years and older, both symptomatic and asymptomatic

Wong-Baker FACES™ Pain Rating Scale

**Fig. 1** Wong-Baker pain rating scale.

patients with cholelithiasis, patients with cholecystitis, a virgin abdomen, and single procedure LC (not with double procedures in the same sitting). Exclusion criteria are as follows: patients with open cholecystectomy, LC converted to open, perforated gallbladder (peritonitis), pregnancy, other abdominal procedures in the same sitting with LC, and abdominal drains. Adequate permission was acquired from the medical ethics committee of the hospital, and all research was done with the informed consent of each patient.

Results

The collected data was entered in Microsoft Excel and analyzed using the Statistical Package for the Social Sciences version 17. A total of 82 patients were analyzed and formulated. The demographic data were analyzed as frequencies, central tendency measures, and dispersion. The mean age (years) of groups A and B ranged from 20 to 80 years. With the highest age in the fifties (26%) followed by forties (20.7%). STP was experienced by 15 patients at 6 hours, wherein 10 patients had a VAS of 2 (study group = 8, control group = 2), 2 had a VAS of 4 from the control group, and 2 had a VAS of 6 from the study group, without a significant difference in both groups ($p = 0.128$ and 0.125 , respectively). STP was experienced by 13 patients at 12 hours, wherein 9 had a VAS of 2 (study group = 7, control group = 2), 3 had a VAS of 4 (study group = 1, control group = 2), and 1 had a VAS of 6 from the control group, without a significant difference in both groups ($p = 0.966$ and 0.966 , respectively). At 18 hours, 8 patients had STP, wherein 5 had a VAS of 2 (4 study, 1 control) and 3 had a VAS of 4 (1 study, 2 control), without significant difference in both groups ($p = 0.689$ and 0.688 , respectively). At 24 hours, 4 patients had STP, wherein 2 had a VAS of 2 from the study group, without a statistically significant difference in both groups ($p = 0.400$ and 0.435 , respectively). The majority were discharged on postoperative day 1 (32 studies, 37 control).

The follow-up VAS after 1 week for STP and abdominal pain were statistically insignificant in both groups.

A total of 82 patients, who planned for elective cholecystectomy, were included in this study and were divided into two groups. All are adult males and females. The gender distribution among the two groups showed a nonsignificant difference (chi-square test $p = 0.82$) (► **Table 1**). The Fisher's exact test was used to study the association between gender and study group, and the Student's *t*-test was used to study the association between study group and age, STP (VAS score), and abdominal pain (VAS score). STP VAS at 30, 36, and 48 hours were statistically insignificant. Abdominal pain VAS at 6, 12, 18, 24, 30, 36, and 48 hours was statistically insignificant (► **Table 2**).

Discussion

Nowadays, LC is one of the common procedures in the general surgery and gastroenterology departments. The learning curve for laparoscopy is long; however, surgical

Table 2 STP VAS and abdominal pain VAS were statistically insignificant

Variable	<i>p</i> -Value
Shoulder tip pain 6 h	0.128
	0.125
@12	0.966
	0.966
@18	0.689
	0.688
@24	0.400
	0.435
@30	0.718
	0.709
@36	0.803
	0.795
@48	0.781
	0.772
Abdomen pain vas 6 h	0.162
	0.163
@12	0.685
	0.685
@18	0.589
	0.589
@24	0.327
	0.334
@30	0.271
	0.245
@36	1.000
	1.000
@42	0.776
	0.611
@48	0.754
	0.746
Follow-up	0.777
	0.777

Abbreviations: STP, shoulder tip pain; VAS, Visual Analog Scale.

trainees are remarkably exposed to this procedure, thereby making them more comfortable with performing LC than other procedures.¹¹ Particularly, the postoperative care for these patients must be adequate. This study is conducted to understand the benefits of intraperitoneal bupivacaine irrigation in reducing postoperative abdominal pain and STP. Few study outcomes showed variable results although many studies were of high quality. The present study aimed to obtain high quality evidence for the same. This study showed no particular advantage in the use of bupivacaine.

Radiologic studies demonstrate the presence of pneumoperitoneum for as long as 24 hours after LC, The amount of

free gas remaining in the abdominal cavity is usually small and has not been regarded as a cause of postoperative morbidity. Components of post-LC pain include visceral pain (78.33%), parietal pain (70%), and STP (23.33%). Intraperitoneal irrigation of the diaphragmatic surface and gallbladder fossa using normal saline, bupivacaine, or lignocaine may effectively control visceral abdominal pain after LC.¹² A study observed a reduced postoperative STP in minor gynecological surgery after intraperitoneal instillation of LA.¹³ With the possibility that a similar beneficial effect might be achieved in LC, several studies were conducted with variable results.¹² This study had a small sample size but was conducted in an organized manner. Only the assistant surgeon was aware of the dosage and drug used for irrigating the gallbladder bed. Anesthesia was given following the standard protocol for each case. Postoperative pain management was in union with intravenous analgesia according to the WHO pain ladder. Each data was recorded by the assistant surgeon of the case. A meta-analysis of all studies of intraperitoneal bupivacaine irrigation is required for a better conclusion on the matter. So far, no adverse effects of bupivacaine irrigation have been recorded; hence, it can be used safely by people without allergies.

In conclusion, the present study had a small sample size; thus, drawing a definitive conclusion was difficult. However, this randomized trial was well-conducted and demonstrated that bupivacaine irrigation at the gallbladder bed post LC does not affect pain relief. The background that most patients who underwent LC have minimal pain should also be considered; hence, several patients are required to conclude with a suitable outcome to demonstrate a tangible and significant difference between both groups.

Abbreviations

LC	Laparoscopic Cholecystectomy
OC	Open Cholecystectomy
LA	Local Anesthesia
STP	Shoulder Tip Pain
VAS	Visual Analog Scale

Authors' Contributions

All authors contributed equally.

Funding

This project did not receive any funding from any institution or medical center.

Conflict of Interest

Hereby we confirm that there is no conflict of interests between authors.

Acknowledgment

We would like to thank all the patients, staff, and managerial team of Sagar Hospitals, Bangalore, Karnataka, India.

References

- Castillo-Garza G, Díaz-Elizondo JA, Cuello-García CA, Villegas-Cabello O. Irrigation with bupivacaine at the surgical bed for postoperative pain relief after laparoscopic cholecystectomy. *JLS* 2012;16(01):105–111
- Golubović S, Golubović V, Cindrić-Stancin M, Tokmadžić VS. Intraperitoneal analgesia for laparoscopic cholecystectomy: bupivacaine versus bupivacaine with tramadol. *Coll Antropol* 2009;33(01):299–302
- Wasim M, Shafqatullah SM, Subhan A. Role of intraperitoneal bupivacaine in laparoscopic cholecystectomy. *Pak J Surg* 2014;30(03):222–226
- Zmora O, Stolik-Dollberg O, Bar-Zakai B, et al. Intraperitoneal bupivacaine does not attenuate pain following laparoscopic cholecystectomy. *JLS* 2000;4(04):301–304
- Kumar M, Negi B, Chauhan A, Jhobta RS, Bhandari S. Effect of subdiaphragmatic and gall bladder bed administration of bupivacaine on post operative pain and post operative pulmonary functions in laparoscopic cholecystectomy. *IOSR J Dent Med Sci* 2015;14(01):21–26
- Alkhamisi NA, Peck DH, Lomax D, Darzi AW. Intraperitoneal aerosolization of bupivacaine reduces postoperative pain in laparoscopic surgery: a randomized prospective controlled double-blinded clinical trial. *Surg Endosc* 2007;21(04):602–606
- Latif Kalas G, Khan FH, Khan KH, Malik N, Anwar I. Comparison of intraperitoneal bupivacaine with intravenous tramadol hydrochloride for postoperative pain relief after laparoscopic cholecystectomy. *Journal of University Medical & Dental College* 2019;10(01):1–6
- Golash V. Laparoscopic Primary prosthetic repair of Hiatus hernia with GERD in 88 patients. *British J Surg* 2012;99:57–57
- Ahmed F, Narula H, Khandelwal M, Dutta D. A comparative study of three different doses of nalbuphine as an adjuvant to intrathecal bupivacaine for postoperative analgesia in abdominal hysterectomy. *Indian Journal of Pain* 2016;30(01):23
- Bianchi PP, Rosati R, Bona S, et al. Laparoscopic surgery in rectal cancer: a prospective analysis of patient survival and outcomes. *Dis Colon Rectum* 2007;50(12):2047–2053
- Memon F, Gill RC, Baloch S, et al. Conversion of laparoscopic to open cholecystectomy, is gender a predictor? *Pak J Surg* 2014;30(04):290–295
- Joris JL, Jacob EA, Sessler DI, Deleuse JJ, Kaba A, Lamy ML. Spinal mechanisms contribute to analgesia produced by epidural sufentanil combined with bupivacaine for postoperative analgesia. *Anesth Analg* 2003;97(05):1446–1451
- Boddy AP, Mehta S, Rhodes M. The effect of intraperitoneal local anesthesia in laparoscopic cholecystectomy: a systematic review and meta-analysis. *Anesth Analg* 2006;103(03):682–688