

Pediatrics

Wilms Tumor in India: A Systematic Review

Shyam Srinivasan¹ Subramaniam Ramanathan² Maya Prasad¹¹Department of Pediatric Oncology, Homi Bhabha National Institute, Tata Memorial Hospital, Mumbai, Maharashtra, India²Department of Pediatric Oncology and BMT, Great North Children's Hospital, Royal Victoria Infirmary, Newcastle Upon Tyne, United Kingdom**Address for correspondence** Shyam Srinivasan, DM, Department of Pediatric Oncology, Tata Memorial Hospital, Parel, Mumbai 400 012, Maharashtra, India (e-mail: srinivas.shyam@gmail.com).

South Asian J Cancer 2023;12(2):206–212.

Abstract



Shyam Srinivasan

Background Cure rates of childhood malignancies are inferior in India compared with upper-middle-income countries. There is paucity of quality data addressing outcome of childhood Wilms tumor (WT) from India. This systematic review was conducted to assess the disease trends, treatment strategies, and outcome indicators in WT across India.

Materials and Methods We conducted a systematic search of MEDLINE, Google Scholar, and SCOPUS database, and additionally screened International Society of Pediatric Oncology conference abstracts. Data concerning WT or nephroblastoma published from India were extracted.

Results A total of 17 studies containing 1,170 patients were included in this review. Ninety-four percent of the studies were published after the year 2010. Advanced stage (III and IV) disease was seen in 46% of included patients. In seven studies, patients underwent a pretreatment biopsy before commencement of therapy. A hybrid approach consisting of “surgery first” in a selected subset and “neo-adjuvant chemotherapy” in all others was the most common treatment strategy adopted in half of the studies. The overall survival ranged between 48 and 89%. Key prognostic factors influencing survival across studies included increased tumor volume, metastatic disease, and unfavorable histology. Nonrelapse mortality (2.7–8.5%) was noted to be high.

Conclusion Substantial proportion of children with WT from India present with advanced stages of the disease. Despite several limitations, the current systematic review showed a modest survival among Indian children with WT. Adopting strategies through collaboration to ensure early access to expert care along with involvement of social support team to improve compliance may further improve survival of WT in India.

Keywords

- ▶ Wilms tumor
- ▶ India
- ▶ survival
- ▶ prognostic factors

DOI <https://doi.org/10.1055/s-0042-1758567> ISSN 2278-330X

How to cite this article: Srinivasan S, Ramanathan S, Prasad M. Wilms Tumor in India: A Systematic Review. South Asian J Cancer 2023;12(2): 206–212.

© 2022. MedIntel Services Pvt Ltd. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

Introduction

Wilms tumor (WT) is one of the major success stories in pediatric oncology.¹ Most of the data pertaining to WT has emerged from two major cooperative groups, SIOP (International Society of Pediatric Oncology) and COG (Children's Oncology Group), which have shown improved outcomes over the years with intensive multimodality treatment. The emphasis has been on accurate histological diagnosis and risk stratification, tailored surgical resection, and timely radiotherapy and chemotherapy. While outcomes of localized favorable histology (FH) and that of metastatic WT in developed countries have exceeded 90 and 80%, respectively, the outcomes in lower and middle-income countries (LMICs) and low-income countries (LICs) are relatively inferior.¹⁻⁵ Several factors, including poor access to care, delayed diagnosis and referral, treatment abandonment, treatment related mortality, malnutrition, and lack of expertise in local management are considered to be the contributing factors.^{4,6-8} The purpose of this systematic review is to provide a succinct overview of the epidemiology, clinical characteristics, treatment strategies, outcomes, and prognostic variables in WT across India, based on published literature.

Materials and Methods

We conducted a systematic search of the MEDLINE, Google Scholar, and SCOPUS database for published studies on WT in India. A literature search was performed using text words "Wilms tumor," "nephroblastoma," and "India." Articles published from time of inception of database till 31/08/2021 were included. In addition, studies published as SIOP conference abstracts from the year 2011 to 2020 were screened for data regarding WT published from India. Studies that were published languages other than English and studies containing less than 10 subjects were excluded. Literature search was as per preferred reporting items for systematic reviews and meta-analyses (PRISMA) statement.

Study Selection

The review authors independently screened the titles and abstracts yielded by the search against the inclusion and exclusion criteria. Full reports for all titles and abstracts were obtained if they appeared to meet the inclusion criteria and also in case of any uncertainty. Two reviewers independently extracted the data to reduce the bias and errors in data extraction. In case of disagreements, a third reviewer was consulted for a final decision. The data obtained were integrated into evidence tables and were verified by the two reviewers.

Data Items

The information that was extracted from each study included the surname of first author, year of study, age with range, number of patients, sex ratio, symptomatology, diagnostic approach (biopsy, imaging modality), laterality of disease,

data regarding biopsy, staging of disease, timing of surgery, chemotherapy backbone, event-free survival, overall survival, and prognostic factors.

Results

Literature Search

A total of 826 studies and 14 SIOP conference abstracts were obtained after the initial search. After removing duplicates and screening the title and abstract of the publications, full text of 29 studies were selected for eligibility and 25 could be retrieved. From these, 6 studies included less than 10 patients, 2 studies were duplicates, 1 study did not include WT, 2 studies did not have required data, and 1 study included patients outside India. All these 12 studies were subsequently excluded. Of the 14 SIOP abstracts, 10 abstracts were excluded because of less than 10 patients being enrolled ($n=3$), desired outcomes not being available ($n=2$), studies being subsequently published ($n=3$), and study population overlap ($n=2$). Eventually, 17 studies were included in the systematic review. The PRISMA flowchart is shown in [Fig. 1](#).

Quality of Studies

The Newcastle-Ottawa scale adapted for cross-sectional studies was used to assess the quality of included studies ([Supplemental Table S1](#), available online only).⁹ The quality of the study was unsatisfactory in 3 of the 16 studies included in the systematic review.

Characteristics of the Studies

Of the 16 studies, 2 (13%) were prospective, 1 (6%) was ambispective, and 13 (81%) were retrospective. The salient features of the studies are summarized in [Table 1](#).

Patient Characteristics

A total of 1,170 patients were included from the 17 studies.¹⁰⁻²⁶ Data on presenting symptoms was gathered from eight studies,^{13,14,16,18,19,22,25,26} among which abdominal lump was the most common. Abdominal pain, hypertension, and hematuria were other symptoms seen in 19 to 39%, 8 to 42%, and 4 to 16%, respectively. Systemic symptoms, namely fever, were noted in 26 to 30% of patients.^{14,16,26} Median duration of symptoms prior to presentation at place of care varied between 28 and 35 days.^{19,26} Presence of concomitant congenital anomalies/syndromic associations was noted in three studies that was 3.7, 6.6, and 16%, respectively.^{13,19,22}

All studies employed ultrasonography (US) or computed tomography (CT) to assess locoregional extent and chest radiograph or CT chest to screen for distant metastases. However, the categorical proportion of patients who underwent either US/CT abdomen/CT chest was available in three studies.^{10,16,19} Measurements on tumor volume at baseline were available in two studies, who observed a reduction from mean volume of 523 mL and 481.7 mL to 208 mL, and 109 mL, respectively, following neoadjuvant chemotherapy.^{16,17} Data on baseline staging was available in 16 studies




Fig. 1 Flow diagram of the systematic review according to preferred reporting items for systematic reviews and meta-analysis (PRISMA) guidelines.

of which 1 study each included only metastatic patients and bilateral tumors (Stage V).^{12,20} Of the remaining 14 studies, the proportion of patients with stage III and IV disease ranged between 9–55% and 6–30%, respectively. Overall, the mean incidence of advance stage disease, i.e. stage III and IV combined, was 45% (95%CI: 37–54%). Mean incidence of bilateral WT was 5.8% (95%CI: 4–7.6%).

Information on histological subtypes was available in 13 studies. The pathological subclassification and usage of terminologies were noted to be congruent with the treatment backbone (SIOP vs. NWTG) in eight (62%) of these studies.^{10,16,20–24,26} Overall, high risk/anaplasia/unfavorable histology was reported to be 14% (95%CI: 8–20%) from these studies.

Treatment Strategy

Patients in seven studies had pretreatment histological confirmation prior to initiation of therapy by way of fine-needle aspiration (FNA)^{12,24} or biopsy.^{10,11,14,16,21,26} Data on treatment modality was available from 13 studies and differed across centers. Preoperative chemotherapy and upfront surgery were the standard practice in three (23%) and two (15%) studies, respectively, while remaining seven (54%) studies used a hybrid approach utilizing the benefits of both the approaches: “surgery first” in a selected subset and “neo-adjuvant chemotherapy” in others. One study employed the former strategy for a period of 8 years and switched to the latter after 2008 until present.¹⁵

Among the studies where information on chemotherapy was available ($n = 15$), International Society of Pediatric

Oncology (SIOP)-based protocol was used in four (27%) studies, National Wilms Tumor Study Group/Children’s Oncology Group (NWTG/COG)-based in five (33%) studies, UK WT3 regimen in one (6%) study, and institutional-based protocol in three (20%) studies. Two (13%) studies used different protocols at different time points.^{13,14}

Outcome

Nonrelapse mortality attributable to toxic deaths was 2.7 to 8.5%.^{10,12,16,17,19} Surgical complications were reported in six studies^{15,16,18–20,24} and varied between 11 and 29% of which postoperative ileus and sepsis were most common. Kajal et al noted a high incidence of intestinal perforation (13%), while Verma and Kumar attributed 50% of toxic deaths to postoperative complications.^{18,19} Jain et al noted a higher incidence of intraoperative tumor spillage, mortality, and tumor recurrence among patients who underwent upfront surgery.¹⁵ Survival estimates for the individual studies are summaries in **Table 1**. Regional nodal disease, increased tumor volume, metastatic disease, and unfavorable histology were noted to be adversely influencing survival, in the above studies.

Discussion

WT is the most common childhood renal tumor. While the prevalence of unilateral WT is reported to be marginally higher in girls in Western literature,²⁷ the present review shows a higher male prevalence reflecting the existing gender bias in health-seeking behavior.²⁸ Though most of the studies did not

Table 1 Characteristics of 17 studies included in the systematic review

Study	Type of publication	Study period	n	Mean age (mo)	M:F	Type of study	Baseline biopsy or FNAC	Stage (I/II/III/IV)					Timing of surgery	Histological classification	Chemotherapy backbone	Toxic deaths/NRM	EFS	OS	Poor prognostic factors and other comments
								I	II	III	IV	V							
Qureshi et al 2020 ¹⁰	Research article	2015–19	113	42.5	1.6:1	Prospective	Biopsy	16%	18%	24%	35%	–	Hybrid	LR: 4% IR: 78% HR: 18%	SIOP	5.3%	3-year OS: 81.8 (74.1–85.6)%	Lymph node positivity, stage of disease	
Jeevarathi and Vadivelu 2020 ¹¹	Research article	2008–11	156	NA	1.7:1	Retrospective + prospective	Biopsy	19%	39%	29%	6%	6%	NA	FH: 98.7% UH: 1.3%	NA	NA	NA	LOH 1p, LOH 16q	
Jain et al 2020 ¹²	Research article	2000–12	36	56.2	1.8:1	Retrospective	FNAC	–	–	–	100%	–	Hybrid	NA	AIMS-WT-99	<2.7%	4-year OS: 48% (41.3–75.9)	Liver metastases, poor response to chemotherapy	
Sachdeva et al 2019 ¹³	Conference paper	2005–18	71	30	NA	Retrospective	NA	20%	20%	43%	10%	8%	Hybrid	FH: 82.9% UH: 13.1%	Before 2013: UKCCSG From 2014: SIOP/COG	NA	5-year OS: 87.4%	Upfront chemotherapy, UFH, Stage III/IV	
Wani et al 2019 ¹⁴	Research article	2010–15	23	48	1:1	Retrospective	Biopsy (in 48% patients)	22%	4%	44%	22%	4%	Hybrid	NA	SIOP and COG	NA	NA	NA	
Jain et al 2019 ¹⁵	Conference paper	1998–2018	40	36	2.3:1	Retrospective	No	27%	32%	35%	20%	5%	(see note) ^a	NA	NA	NA	NA	NA	Upfront surgery associated with spillage, mortality and tumor recurrence Abandonment rates: 11–31%
John et al 2018 ¹⁶	Research article	2004–14	59	58	1.4:1	Retrospective	Biopsy	38%	16%	9%	30%	10%	Upfront chemotherapy	LR: 7%; IR: 86% HR: 7%	SIOP	8.5%	3-year OS: 80%	Stage III and IV	
Rahman et al 2018 ¹⁷	Conference paper	2005–16	200	33.5	NA	Retrospective	NA	30%	36%	14%	17%	3%	NA	NA	SIOP	7.2%	3-year OS: 78.5%	Large tumor volume, Stage IV disease Undernutrition Abandonment rates: 20%	
Kajal et al 2017 ¹⁸	Research paper	2008–12	31	40	0.8:1	Retrospective	NA	36%	–	55%	9%	4.5%	Hybrid	Anaplasia: 3.2%	NWTS	NA	NA	NA	
Verma and Kumar 2016 ¹⁹	Research paper	2005–14	108	33	4:1	Retrospective	No	21%	30%	35%	10%	4%	Hybrid	Anaplasia: 11%	NWTS	5.5%	5-year OS: 74%	Stage III and IV	
Agarwala et al 2014 ²⁰	Research paper	1999–2000	11	Range: 6–30	2.7:1	Prospective	No	–	–	–	–	100%	Upfront chemotherapy	FH: 100%	AIMS-WT-99	NA	8 of 11 alive at follow-up	NA	
Bastogi et al 2014 ²¹	Conference paper	1990–2006	147	40	1.4:1	Retrospective	NA	NA	NA	NA	NA	NA	Hybrid	FH: 98.6%	NWTS	NA	10-year OS: 89%	NA	
Guruprasad et al 2013 ²²	Research paper	2003–10	61	40	0.9:1	Retrospective	No	28%	16%	38%	15%	3%	Upfront surgery	FH: 72% Anaplasia: 16.4%	NWTS	NA	5-year OS: 85.2%	UFH	
Yadav et al 2013 ²³	Letter to the editor	2005–10	22	34	0.8:1	Retrospective	NA	7%	33%	44%	4%	11%	NA	FH: 74% UFH: 26%	UK WT3	NA	5-year EFS/OS: 73%/77%	UFH	
Trehan et al 2012 ²⁴	Research paper	1999–2003	20	20	2.3:1	Retrospective	FNAC	20%	65%	15%	–	–	Upfront chemotherapy	FH: 100%	SIOP 6 protocol	NA	75%	NA	
Chang et al 2011 ²⁵	Conference paper	2006–09	52	51	2.2:1	Retrospective	NA	78%	–	–	22%	NA	NA	FH: 77% UFH: 23%	3 drug regimen (VAC regimen)	NA	DFS: 37.8 mo	NA	
Qureshi et al 2007 ²⁶	Research article	1997–2000	20	Range: 9–144	0.8:1	Retrospective	Biopsy (selective)	50%	20%	20%	5%	5%	Upfront surgery	FH: 90% UFH: 20%	NWTS	NA	30% recurrence	NA	

Abbreviations: DFS, disease-free survival; EFS, event-free survival; FH, favorable histology; FNAC, fine-needle aspiration cytology; IR, intermediate risk; HR, high risk; LOH, loss of heterozygosity; LR, low risk; M:F, male: female; NA, not available; NRM, nonrelapse mortality; NWTS, –; OS, overall survival; SIOP, International Society of Pediatric Oncology; UFH, unfavorable histology; VAC, Vincristine, doxorubicin, cyclophosphamide.

^aUpfront surgery was performed between 1999 and 2007 and Upfront chemotherapy was given between 2008 and 2018.

report on the duration of symptoms, with the available literature from two studies, patients seem to be symptomatic for a prolonged duration (>4 weeks) at the time of presentation. Obtaining a specimen for histology prior to commencing treatment was practiced in 64% of the included studies. Data from the UK-W3 trial and SIOP WT 2001 trial have shown needle biopsy to be safe and not associated with increased local relapse.^{29,30} Similar observations have been made on two Indian studies that noted no association between increased tumor rupture, needle tract seeding, or tendency to relapse with a baseline biopsy.^{31,32} Two centers have employed FNA cytology^{12,24} instead of biopsy for establishing diagnosis, which has a comparable sensitivity and specificity.³³

The preferred practice in approximately half of the included studies combined the treatment philosophies of both NWTSG/COG and SIOP. The UK-W3 trial evaluated the role of preoperative chemotherapy in achieving a favorable stage distribution and noted that approximately 20% fewer children received doxorubicin or radiation (because of downstaging by neo-adjuvant chemotherapy).³⁰ Of note, none of the 102 patients who received preoperative chemotherapy had a tumor rupture.³⁰ In this review, the proportion of patients operated upfront varied between 45 and 68%. While the precise reason for selecting certain patients for upfront surgery and deferring surgery in others is unclear from these studies, one study looked at leveraging the benefits of both these approaches in a prospective manner in pediatric renal tumors.³² Based on the presence of image-defined risk factors, the study attempted at objectively classifying tumors at high-risk of tumor rupture and slotting them for delayed resection after neoadjuvant chemotherapy. With this approach the authors had noticed a postoperative complication rate comparable to SIOP and COG reports³² with an excellent survival in the immediate surgery group and a comparable survival in the delayed surgery group. Another factor influencing surgical decision is the baseline tumor size. Several Indian studies have noted the median baseline volume at presentation to be ranging between 480 and 520 mL and the presurgery volume (post neoadjuvant chemotherapy) to be ranging between 200 and 300 mL.^{16,17,32} Strikingly, patients treated on SIOP 2001 also had a similar median baseline tumor volume of 570 mL but showed an impressive decline to 180 mL presurgery.³⁴ However, the proportion of patients presenting in advanced stages was higher when compared with that in the SIOP-2001 trial.

In addition to stage, histological features also drive the treatment strategy in WT. Hence, precise classification is crucial in differentiating WT from non-Wilms pathologies and identifying high-risk versus low-risk features in a WT. Studies have shown that blastemal predominant WT to be misclassified into a lower risk group in approximately 15% of cases.³⁵ Histologic terminologies differ between the two treatment philosophies, that is, low-risk/intermediate risk/high risk is used in the context of SIOP/RTSG-guided treatment, whereas NWTSG/COG-guided approach uses the definitions as favorable and unfavorable histology. In 58% of the studies, where definitions were noted to be congruent, the incidence of high risk/unfavorable WT was found to be similar to larger studies.^{34,35}

Timely radiation therapy forms an important component of adjuvant therapy in at least a third of the patients with WT. A surgery to radiotherapy interval of less than or equal to 14 days correlates with an improved survival.³⁶ While few of the studies provide the proportion of patients needing adjuvant therapy, details about timing and doses have not been specified. Current systematic review also highlighted a high incidence (3–9%) of toxic deaths in a disease that is treated majorly with less than or equal to 3 chemotherapy drugs on an outpatient basis. Malnutrition, sepsis with multi-drug resistant organisms, inadequate supportive care, and excess postoperative complications are probable incriminating factors. There were difficulties in assessing the outcomes in this review mostly due to inconsistencies with definition of survival and data regarding lost to follow-up or treatment abandonment being not available from most of the studies. Despite these setbacks, the current review showed an acceptable survival of 74 to 87% among Indian patients.

According to World Health Organization, the estimated incidence of WT among children between 0 and 14 years in India is 4.4 age-standardized rates per million.⁴ Children of 0 to 14 years constitute 26% of the total 1.38 crore population of India.³⁷ As of date, the expected incidence of WT in India is 1,600 cases per year. Though the incidence of renal tumors is not associated with economic status, mortality is relatively higher in LICs and LMICs. Several barriers exist in childhood cancer treatment and these extend to WT as well. Inappropriate use of chest radiographs instead of CT scans for screening metastases, inaccurate histopathological risk stratification, lack of adherence to nodal sampling alongside nephrectomy, and inability to deliver timely radiation are some barriers specific to WT. In addition, delayed presentation, delayed diagnosis, sociocultural factors such as illiteracy, gender bias, and poor access to care further hamper survival.^{6,38,39} Employing national guidelines to tailor therapy and conducting collaborative trials would improve treatment delivery and build expert clinical capacity.⁴⁰ While the developed countries continue to improve upon the benchmark of overall survival of 90%, with insights into tumor biology and molecular classification being the next step, our immediate steps must be directed at surmounting the aforementioned challenges.

Limitations of the Study

This study was limited by predominance of retrospective and poorly designed studies. There were difficulties in assessing the outcomes mostly due to inconsistencies with definition of survival and data regarding lost to follow-up or treatment abandonment being not available from most of the studies.

Conclusion

The current systematic review highlights the disease trends and treatment strategies of WT in India. Substantial proportion of patients present with advanced stages of the disease and survival seems to be inferior in comparison to developed countries. There is a need to assess for prognostic factors in

Indian patients that will guide in intensifying therapy in selective patients. In addition, long-term follow-up with assessment of quality of life is necessary to establish overall success of therapy. Population-based registries are required to estimate disease burden that would help in health planning and allocation of resources.

Authors' Contributions

Shyam Srinivasan contributed to conceptualization, methodology, formal analysis, writing—original draft. Subramaniam Ramanathan was involved in formal analysis, data curation, and writing—original draft. Maya Prasad contributed to supervision, writing—review and editing, and project administration.

Funding

None.

Conflict of Interest

None declared.

References

- Dome JS, Graf N, Geller JI, et al. Advances in Wilms tumor treatment and biology: progress through international collaboration. *J Clin Oncol* 2015;33(27):2999–3007
- Dix DB, Seibel NL, Chi Y-Y, et al. Treatment of stage IV favorable histology Wilms tumor with lung metastases: a report from the Children's Oncology Group AREN0533 Study. *J Clin Oncol* 2018;36(16):1564–1570
- Ghafoor T, Bashir F, Ahmed S, Khalil S, Farah T. Predictors of treatment outcome of Wilms tumour in low-income country; single centre experience from Pakistan. *J Pediatr Urol* 2020;16(03):375.e1–375.e7
- Cunningham ME, Klug TD, Nuchtern JG, et al. Global disparities in Wilms tumor. *J Surg Res* 2020;247:34–51
- Ekenze SO, Okafor OC, Obasi AA, Okafor DC, Nnabugwu II. Wilms tumor in Africa: a systematic review of management challenges and outcome in two decades (2000–2019). *Pediatr Blood Cancer* 2020;67(11):e28695
- Israels T, Moreira C, Scanlan T, et al. SIOP PODC: clinical guidelines for the management of children with Wilms tumour in a low income setting. *Pediatr Blood Cancer* 2013;60(01):5–11
- Wilde JCH, Lameris W, van Hasselt EH, Molyneux EM, Heij HA, Borgstein EG. Challenges and outcome of Wilms' tumour management in a resource-constrained setting. *Afr J Paediatr Surg* 2010;7(03):159–162
- Libes J, Oruko O, Abdallah F, et al. Risk factors for abandonment of Wilms tumor therapy in Kenya. *Pediatr Blood Cancer* 2015;62(02):252–256
- Ottawa Hospital Research Institute [Internet][cited 2021 Nov 10] Accessed October 28, 2022 at: http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp
- Qureshi SS, Bhagat M, Kazi M, et al. Standardizing lymph nodal sampling for Wilms tumor: a feasibility study with outcomes. *J Pediatr Surg* 2020;55(12):2668–2675
- Jeevarathi T, Vadivelu G. Long term outcome of Wilms' tumor in tertiary care hospital [Internet]. *Int Surg J* 2020;7:3234–3238 [cited 2021 Nov 10]. Accessed October 28, 2022 at: <https://www.ijurgery.com/index.php/ij/article/view/6573>
- Jain V, Dhua A, Agarwala S, et al. Outcome of children with Stage IV Wilms tumor - our experience of 15 years. *J Indian Assoc Pediatr Surg* 2020;25(06):372–377
- Sachdeva P, Danewa A, Thatikonda KB, Paul T, Kalra M, Sachdeva A. Managing Wilms Tumor: A Single Centre Experience From North India. *Pediatric Hematology Oncology Journal* 2019;4(02):S54
- Wani SQ, Khan T, Wani SY, Lone MM, Afroz F. Wilms' tumor-collaborative approach is needed to prevent tumor upstaging and radiotherapy delays: a single institutional Study. [Internet] *Indian J Med Paediatr Oncol* 2019;40:409–412
- Jain N, Kumar S, Sarin Y. V432 SIOP19–1414 Wilms' Tumor Revisited. *Pediatr Blood Cancer* 2019;66(S4):e27989
- John R, Kurian JJ, Sen S, et al. Clinical outcomes of children with Wilms tumor treated on a SIOP WT 2001 protocol in a tertiary care hospital in south India. *J Pediatr Urol* 2018;14(06):547.e1–547.e7
- Rahiman E, Trehan A, Jain R, et al. O-031 tumor volume and under-nutrition at diagnosis adversely affect the outcome of children with Wilms tumor: a study of 200 patients. [Internet] *Pediatr Blood Cancer* 2018;65(S2):e27455
- Kajal P, Rattan KN, Bhutani N, Marwah N. Wilms' tumor: histopathological variants and the outcomes of 31 cases at a tertiary care center in Northern India. *Middle East J Cancer* 2017;8:143–150
- Verma N, Kumar A. Clinicoepidemiological profile and outcome of children with Wilms tumor in a developing country. *J Pediatr Hematol Oncol* 2016;38(07):e213–e216
- Agarwala S, Mittal D, Bhatnagar V, et al. Management and outcomes in massive bilateral Wilms' tumors. *J Indian Assoc Pediatr Surg* 2014;19(04):208–212
- Rastogi S, Qureshi S, Vora T, Chinnaswamy G, Prasad M, Laskar S, Khanna N, Ramadwar M, Medhi, Seema, Kurkure P. P 274. Is three drug chemotherapy protocol for all stages of Wilms Tumor a practical compromise for suboptimal staging in developing countries? Is it worth and safe? *Pediatr Blood Cancer* 2014;61(S2):S105–S433
- Guruprasad B, Rohan B, Kavitha S, Madhumathi DS, Lokanath D, Appaji L. Wilms' tumor: single centre retrospective study from South India. *Indian J Surg Oncol* 2013;4(03):301–304
- Yadav SP, Manchanda H, Sachdeva A. Wilms tumor: a success story in the developing world!. *J Pediatr Hematol Oncol* 2013;35(08):632
- Trehan A, Chowdhary SK, Marwaha RK. Wilms tumor: five-year tumor-free survival on a modified SIOP protocol from an Indian university hospital. *J Pediatr Hematol Oncol* 2012;34(01):57–62
- Chander S, Pathy S, Mohanti B, Agarwal S, Yadav V, Bakshi S. Multidisciplinary approach to treatment of Wilms tumor - AIIMS experience. *Pediatr Blood Cancer* 2011;57(05):705–897
- Qureshi MF, Al Bakrah M, Shenoy UV. Clinical features and treatment strategies of Wilm's tumor: A setup in the last decade of the millennia and possible inclusion of advance researches to improve the clinical management. *J Med Sci* 2007;7:797–803
- Stiller CA, Olshan AF. (2014). Epidemiology of Renal Tumours of Childhood. In: Pritchard-Jones, K, Dome, J. (eds) *Renal Tumours of Childhood*. Pediatric Oncology Springer, Berlin, Heidelberg https://doi.org/10.1007/978-3-662-44003-2_1
- Pant G, Verma N, Kumar A. Gender bias in health seeking behaviour of families of childhood cancer patients from north India: a 6 year retrospective observational study [Internet]. *Pediatric Hematology Oncology Journal* 2019;4(02):S13
- Irtan S, Jitlal M, Bate J, et al;Renal Tumours Committee of the Children's Cancer and Leukaemia Group (CCLG) Risk factors for local recurrence in Wilms tumour and the potential influence of biopsy - the United Kingdom experience. *Eur J Cancer* 2015;51(02):225–232
- Mitchell C, Pritchard-Jones K, Shannon R, et al;United Kingdom Cancer Study Group. Immediate nephrectomy versus preoperative chemotherapy in the management of non-metastatic Wilms' tumour: results of a randomised trial (UKW3) by the UK

- Children's Cancer Study Group. [Internet] *Eur J Cancer* 2006;42(15):2554–2562
- 31 Kurian JJ, Nongpiur KRL, Jehangir S. Use of pretherapy core biopsy in the diagnosis of pediatric renal tumors. [Internet] *J Indian Assoc Pediatr Surg* 2018;23(02):66–69
 - 32 Qureshi SS, Kembhavi SA, Bhagat M, et al. Customized approach for upfront or delayed resection using radiological criteria in unilateral, nonmetastatic pediatric renal tumors: a prospective study. *Pediatr Blood Cancer* 2019;66(suppl 3):e27815
 - 33 Gupta S, Srinivasan R, Trehan A, et al. Diagnostic accuracy and cytomorphological spectrum of Wilms tumour in fine needle aspiration biopsy cytology samples supplemented with cell blocks. *Pediatr Blood Cancer* 2021;68(07):e28996
 - 34 Irtan S, Van Tinteren H, Graf N, et al. Evaluation of needle biopsy as a potential risk factor for local recurrence of Wilms tumour in the SIOP WT 2001 trial. *Eur J Cancer* 2019;116:13–20
 - 35 van den Heuvel-Eibrink MM, van Tinteren H, Bergeron C, et al. Outcome of localised blastemal-type Wilms tumour patients treated according to intensified treatment in the SIOP WT 2001 protocol, a report of the SIOP Renal Tumour Study Group (SIOP-RTSG). *Eur J Cancer* 2015;51(04):498–506
 - 36 Stokes CL, Stokes WA, Kalapurakal JA, et al. Timing of radiation therapy in pediatric Wilms tumor: a report from the national cancer database. [Internet] *Int J Radiat Oncol Biol Phys* 2018;101(02):453–461
 - 37 Population ages 0–14 (% of total population) - India | Data [Internet][cited 2021 Nov 10] Available from: <https://data.worldbank.org/indicator/SP.POP.0014.TO.ZS?locations=IN>
 - 38 Smith MA, Seibel NL, Altekruse SF, et al. Outcomes for children and adolescents with cancer: challenges for the twenty-first century. *J Clin Oncol* 2010;28(15):2625–2634
 - 39 Ribeiro RC, Steliarova-Foucher E, Magrath I, et al. Baseline status of paediatric oncology care in ten low-income or mid-income countries receiving My Child Matters support: a descriptive study. *Lancet Oncol* 2008;9(08):721–729
 - 40 Prasad M, Vora T, Agarwala S, et al. Management of Wilms tumor: ICMR consensus document. *Indian J Pediatr* 2017;84(06):437–445