

New Knowledge About Old Drugs: The Anti-Inflammatory Properties of Cardiac Glycosides*

Authors

Robert Fürst, Ilse Zündorf, Theo Dingermann

Affiliation

Institute of Pharmaceutical Biology, Biocenter,
Goethe University, Frankfurt/Main

Key words

cardiac glycosides, ouabain, inflammation, animal models,
molecular mechanisms

received January 27, 2017

revised February 20, 2017

accepted March 2, 2017

Bibliography

DOI <http://dx.doi.org/10.1055/s-0043-105390>

Published online March 15, 2017 | *Planta Med* 2017; 83: 977–984 © Georg Thieme Verlag KG Stuttgart · New York | ISSN 0032-0943

Correspondence

Robert Fürst, PhD

Institute of Pharmaceutical Biology, Biocenter,
Goethe University

Max-von-Laue-Str. 9, 60438 Frankfurt/Main

Phone: + 49 69 7982 96 55, Fax: + 49 69 7 98 76 32 96 55

fuerst@em.uni-frankfurt.de

ABSTRACT

In the 19th century, cardio-active steroid glycosides, shortly cardiac glycosides, were scientifically established as drugs against heart failure. Their *in vivo*, cellular, and molecular actions as well as their predominant target, Na⁺-K⁺-ATPase, have been comprehensively investigated in the 20th century and the discovery of endogenous cardiac glycosides has fostered this research field. In the last years, however, results from clinical trials and meta-analyses have questioned their therapeutic value due to efficacy and safety issues. This has led to a considerable decline of their usage. Beyond the cardiovascular system, cardiac glycosides have been increasingly recognized as antitumor compounds and Na⁺-K⁺-ATPase has evolved into a promising drug target in oncology. A wealth of review articles exists that intensively discuss these topics. Surprisingly, the anti-inflammatory actions of cardiac glycosides, which were discovered in the 1960s, have so far hardly been perceived and have not yet been summarized. This review provides an overview of the *in vivo* and *in vitro* actions of cardiac glycosides on inflammatory processes and of the signaling mechanisms responsible for these effects: cardiac glycosides have been found to decrease inflammatory symptoms in different animal models of acute and chronic inflammation. Regarding the underlying mechanisms most research has focused on leukocytes. In these cells, cardiac glycosides primarily inhibit cell proliferation and the secretion of proinflammatory cytokines.

Cardio-Active Steroid Glycosides Against Heart Failure: Ups and Downs

Plants that contain cardio-active steroid glycosides (CGs) have been therapeutically used since ancient times. For instance, the usage of squill, *Drimia maritima* (L.) Stearn (Asparagaceae), was reported in the Ebers Papyrus [1]. At the beginning of the early modern history, the German botanist Leonhard Fuchs, who created the scientific name *Digitalis* for foxglove, described the plant in detail and recommended it for diuresis. However, due to the very difficult dosage, which is based on the narrow therapeutic index of CGs [2], digitalis therapy did not succeed at that time.

The successful usage of foxglove, *Digitalis purpurea* L. (Plantaginaceae; ► Fig. 1), started in 1785 with a publication by the English

physician William Withering. Due to his investigations, he could make clear recommendations on the preparation and dosage of the herbal remedy against dropsy (heart failure-associated edema) [1]. Unfortunately, the digitalis therapy became too successful: It was carelessly propagated as a universal remedy, which led to severe reservations among physicians and, consequently, to a shift away from this herbal remedy [3].

The next revival of digitalis started in the middle of the 19th century and was based on the experimental and clinical research by Ludwig Traube in Berlin and on the isolation of CGs by Oswald Schmiedeberg in Strasbourg [1]. With the beginning of the 20th century, digitalis therapy became the gold standard for the treatment of heart failure. A further research milestone was achieved in 1965 when the German pharmacologist Kurt Repke identified the cell membrane-located ion pump Na⁺-K⁺-ATPase as the molecular target of CGs [4]. Besides their action on this pump, they were also found to exert sympathoinhibitory effects and to increase the vagal tone [5].

* Part of the special issue dedicated to Professor Max Wichtl.

ABBREVIATIONS

CFTR	cystic fibrosis conductance regulator
CG	cardio-active steroid glycoside
DC	dendritic cell
IL	interleukin
iNOS	inducible nitric oxide synthase
I κ B	inhibitor of κ B
LPS	lipopolysaccharide
MAPK	mitogen-activated protein kinase
Na ⁺ -K ⁺ -ATPase	sodium potassium adenosine triphosphatase
NF κ B	nuclear factor κ B
PBMC	peripheral blood mononuclear cells
ROR γ t	retinoid acid receptor-related orphan nuclear receptor γ t
TNF α	tumor necrosis factor α
VCAM-1	vascular cell adhesion molecule 1

The discovery of the binding of CGs to the Na⁺-K⁺-ATPase nourished speculations that endogenous CGs, so-called digitalis- or ouabain-like factors, might exist that serve as ligands of this target. In the late 1980s, hints to the existence of endogenous CGs intensified [6–8] and in the early 1990s, these compounds were eventually discovered [9–13]. Their biosynthesis, tissue of origin, and precise physiological as well as pathophysiological function is an ongoing matter of research and debate [14]. Interestingly, in critically ill patients (not treated with CGs) Berendes et al. [15] found that the occurrence of endogenous CGs in the blood was associated with an increased risk of mortality. The authors speculated that the secretion of endogenous CGs depends not only on the impaired myocardial function and the volume overload in these patients, but also on systemic inflammation. However, it is an open question whether the endogenous CGs are a mechanism of protection or part of the detrimental pathology.

Surprisingly, despite their intensive clinical use, randomized controlled trials with CGs have not been performed until the end of the 20th century. The results of these trials and of the respective meta-analyses were sobering: Both the efficacy and the safety of the digitalis-based therapy of heart failure were questioned [16–22]. Since well-tolerated alternative drugs, such as angiotensin-converting enzyme inhibitors and β -blockers, that show clear benefits on the mortality of heart failure patients are available, the use of cardiac glycosides has declined increasingly.

Cardio-Active Steroid Glycosides as Antitumor Agents: A Well Evolved Field

CGs currently experience a fascinating renaissance in a completely different field: They are evaluated as cancer-preventive as well as anticancer drugs (► Fig. 2). Their antitumor action was discovered 50 years ago: In 1967, Osamu Shiratori [23] from the Japanese company Shionogi reported about the antiproliferative action of

several CGs isolated from *Digitalis purpurea* on neoplastic cells *in vitro* and *in vivo*. Moreover, an epidemiologic study published in 1979 indicated that women who were treated with CGs and who developed breast cancer showed a less aggressive tumor growth and lower metastases formation [24]. These discoveries triggered huge research efforts and, most importantly, fueled the idea of the Na⁺-K⁺-ATPase as a promising drug target for the treatment of different types of cancer. The mechanistic basis of these actions has been intensively studied in the last years: CGs inhibit cancer cell growth by various complex and cell-type dependent mechanisms, e.g. by inducing apoptosis or autophagy, by triggering mitochondrial dysfunction due to an accumulation of calcium, and/or by interfering with the mammalian target of rapamycin (mTOR) and NF κ B signaling. Recent comprehensive reviews that summarize this vibrant field of research can be found under the references [25–30].

A New Frontier: The Anti-Inflammatory Potential of Cardio-Active Steroid Glycosides

A schematic overview of the subsequently described actions of CGs is provided in ► Fig. 2.

Cardio-active steroid glycosides exert anti-inflammatory actions *in vivo*

Acute inflammation

The first report on the influence of CGs on inflammatory processes was published in 1967: Lancaster and Vegad [31] found that pretreatment with ouabain (also known as g-strophanthin) (100 μ g/kg bodyweight; ► Fig. 3) reduced the turpentine-triggered early signs of inflammation (edema formation) in the skin and pleural cavity of sheep. De Vasconcelos et al. [32] tested the action of ouabain in different murine models of inflammation and algisia: The authors reported that ouabain (310 or 560 μ g/kg bodyweight) decreased paw edema formation in mice induced by carrageenan, compound 48/80, zymosan, prostaglandine E₂, and bradykinine. Also the concanavalin A-evoked peritoneal inflammation and the associated presence of leukocytes in the peritoneal cavity was reduced by the CG. Moreover, ouabain exhibited nociceptive actions in the acetic acid-induced writhing response and in the hot plate model. These findings were corroborated and expanded in a study by Leite et al. [33], who showed that ouabain (560 μ g/kg bodyweight) decreased zymosan-induced peritonitis in mice. The compound reduced fluid extravasation (edema formation), leukocyte infiltration and the levels of the cytokines IL-1 β and tumor necrosis factor α (TNF α), but did not influence the viability or function (phagocytosis) of leukocytes. The reduced leukocyte extravasation was attributed to a reduced migratory capacity evoked by ouabain. Moreover, ouabain was found to inhibit the zymosan-induced DNA-binding activity of the prototypical proinflammatory transcription factor NF κ B. Taken together, these studies provide clear evidence that ouabain effectively prevents acute inflammation *in vivo*.

Infection-induced inflammation

The activation of inflammatory processes is a prerequisite for the successful elimination of pathogens. Anthony Esposito [34] published his findings on the action of digoxin (4 µg/kg bodyweight per day; ► **Fig. 4**) on the inflammatory response in murine pneumococcal pneumonia. The digoxin-treated animals exhibited a higher mortality rate, since they were unable to clear the infection. Interestingly, these mice showed a decreased recruitment of granulocytes and macrophages into the lung tissue, whereas, at least *in vitro*, digoxin did not affect the bacteria-killing action of alveolar macrophages. Jacob et al. [35] investigated the action of ouabain in murine leishmaniasis. The CG effectively prevented the infiltration of the peritoneal cavity with neutrophils and reduced the levels of TNFα and interferon γ (IFNγ) in the peritoneal exudate.

Chronic inflammatory autoimmune diseases


Regarding autoimmune diseases, digoxin was first evaluated in an animal model resembling multiple sclerosis: Huh and coworkers [36] were seeking for small molecule inhibitors of the retinoid acid receptor-related orphan nuclear receptor RORγt, a transcription factor required for the differentiation of T_H17 cells. In the last years, these cells have been recognized to play a crucial role in autoimmune diseases. In a compound screening approach digoxin was discovered as specific inhibitor of RORγt. In fact, digoxin was found to inhibit murine T_H17 cell differentiation. In murine experimental autoimmune encephalomyelitis, a model of multiple sclerosis, digoxin (40 µg per mouse) delayed the onset and reduced the severity of the disease. Moreover, it strongly decreased the infiltration of the spinal cord with T_H17 cells. To overcome the limitations of CGs regarding their toxicity, the authors searched for chemical modifications of digoxin with a decreased affinity to the Na⁺-K⁺-ATPase and an augmented affinity to RORγt and identified the two compounds 20,22-dihydrodigoxin-21,23-diol and digoxin-21-salicylidene. Inspired by these important findings on the influence of CGs on T_H17 cells, the action of digoxin was tested in two further models of autoimmune diseases: Lee et al. [37] provided evidence for both a preventive and therapeutic action of digoxin (2 and 5 mg/kg bodyweight) on collagen-induced arthritis in mice. Digoxin lowered the joint inflammation and reduced the expression of IL-17, IL-1β, IL-6, TNFα, and IL-21. The digoxin-treated mice showed a decreased amount of T_H17 cells and an increased number of regulatory T cells. *In vitro*, digoxin suppressed the differentiation of T_H17 cells and reduced the production of immunoglobulins from murine B cells. Furthermore, digoxin was tested in experimental autoimmune uveitis in mice. Hinshaw et al. [38] found that digoxin (2 mg/kg bodyweight) reduced uveal inflammation and inhibited the differentiation of T_H17 cells. Unfortunately, this anti-inflammatory action was accompanied by a severe retinal degeneration process that led to the loss of vision. Obviously, digoxin is an improper compound to treat uveitis. These findings strengthen the approach of Huh et al. [36] to separate the actions on RORγt from those on the Na⁺-K⁺-ATPase.

Neuroinflammation

Forshammar et al. [39] reported about the beneficial action of CGs on LPS-activated astrocytes. Both a very low (1 pM) and a very




► **Fig. 1** *Digitalis purpurea* L. (Plantaginaceae), photographed by Dr. Ilse Zündorf.




► **Fig. 2** Major effects of cardiac glycosides.

high concentration (10 µM) of ouabain prevented the downregulation of the Na⁺-K⁺-ATPase, restored the disorganized F-actin cytoskeleton, and reduced the secretion of IL-1β. Forshammar and colleagues [40] also investigated the action of ouabain on rat microglia cells which represent resident brain macrophages. Interestingly, in contrast to astrocytes ouabain did not exert any influence on the LPS-induced secretion of cytokines from microglia. *In vivo* experiments in the context of neuroinflammation were performed by Kinoshita et al. [41] who evaluated the effect of oua-



► Fig. 3 Chemical structure of ouabain (g-strophanthin).



► Fig. 4 Chemical structure of digoxin.

bain (1.8 µg/kg bodyweight) on the hippocampus in a rat model of LPS-induced systemic inflammation/sepsis. Ouabain reduced the LPS-triggered activation of astrocytes, inhibited the raise of the proapoptotic protein Bax, and maintained the expression of the brain-derived neurotrophic factor which indicates a neuroprotective action. Furthermore, ouabain decreased the nuclear translocation of NFκB p65 and the degradation of IκB as well as the expression of IL-1β and of iNOS. Surprisingly, the elevated serum levels of corticosterone and TNFα were not altered by ouabain. Although neither LPS nor ouabain affected the activity of the hippocampal Na⁺-K⁺-ATPase, an alternative target which might be responsible for the observed pharmacological actions of ouabain was not presented.

Cystic fibrosis

Cystic fibrosis is a genetic disease caused by the lack or a malfunction of the chloride channel cystic fibrosis conductance regulator (CFTR). It is characterized by severe lung inflammation and high levels of the proinflammatory cytokine IL-8 in lung tissues. In 2004, Srivastava et al. [42] reported that different CGs (e.g. digitoxin, ouabain, oleandrin) suppress IL-8 hypersecretion *in vitro* in cystic fibrosis lung epithelial cells. With an IC₅₀ of 0.9 nM, digitoxin was the most active CG. Moreover, the authors tested the influence of digitoxin on the global gene expression in these cells and compared the gene expression patterns with those obtained upon restoration of the lacking CFTR ion channel. Interestingly, digitoxin could mimic the effect of this genetic restoration and the overlapping regulated genes were associated with the TNFα/NFκB pathway. This work was followed-up one year later: In a detailed mechanis-

tic analysis, CGs were found to block the TNFα-dependent recruitment of the adaptor protein TNF receptor-associated death domain (TRADD) to the TNF receptor 1 (TNFR1) [43]. This explains the observed inhibitory action of digitoxin on the activation of IKK and on the phosphorylation and degradation of IκB.

Inspired by these two *in vitro* studies, Zeitlin et al. [44] initiated an explorative randomized controlled trial: 24 patients suffering from mild to moderate cystic fibrosis were treated over a period of 28 days with digitoxin (50 µg or 100 µg daily) or with placebo. As primary parameters, the concentration of IL-8 and the count of neutrophils in the sputum were analyzed. The therapy was safe, but steady state blood concentrations of digitoxin could not be achieved until the last few days of the treatment period. Since the duration of the trial was too short, there was no chance to reveal statistically significant outcomes. In fact, digitoxin did not alter IL-8 levels and only slightly reduced the neutrophil count. Larger and much longer studies are needed to answer the question as to whether CGs can beneficially influence inflammatory processes in cystic fibrosis patients.


Ouabain exerts immunomodulatory actions on leukocytes

A schematic overview of the subsequently described effects of CGs on the different types of leukocytes is provided in ► Fig. 5.

Lymphocytes

In 1968, Quastel and Kaplan [45] reported that ouabain (≥ 10 nM) blocks DNA synthesis induced by the mitogen phytohaemagglutinin in isolated human lymphocytes. As expected due to the inhibition of the Na⁺-K⁺-ATPase, the effect could be reversed by increasing potassium concentrations in the culture medium. In a follow-up article [46], the authors corroborated and expanded these findings: ouabain (≥ 10 nM) did not only inhibit the mitogen-triggered DNA, but also the RNA and protein synthesis in lymphocytes. Of note, the inhibitory action of ouabain (1–10 µM) on the *de novo* protein synthesis was confirmed in a recent work by Takada et al. [47], who additionally showed that ouabain decreases the Na⁺-dependent amino acid transport across cell membranes. However, the authors used a human lung carcinoma cell line in this study. After these initial reports, the field was not advanced for a decade. Then, in 1980, Szamel et al. [48] published an article in which they refined the findings of Quastel and Kaplan on the inhibition of ouabain on lymphocytes [45,46]. They analyzed the action of ouabain (20 nM) in dependence of the cell cycle status and showed that lymphocytes were most sensitive to ouabain within the S phase. In a further article, Szamel et al. [49] reported that ouabain (5 µM) is also able to block a very early event in the activation of lymphocytes, i.e. the altered metabolism of plasma membrane phospholipids (e.g. oleate incorporation), by inhibiting the acyl-CoA lysolecithin-acyltransferase due to a functional and spatial association of this enzyme with Na⁺-K⁺-ATPase. Furthermore, ouabain (100 nM) reduced both the phytohaemagglutinin- and the phorbol ester-induced proliferation of peripheral blood lymphocytes [50]. Despite this inhibition of proliferation, the lymphocytes could still be activated.

As a consequence of the discovery of IL-2 as an important regulator of T cells in the late 1970s, Stoeck et al. [51] wondered



whether the actions of ouabain on T lymphocytes could depend on alterations in the production of IL-2 and/or the response to this cytokine. Indeed, ouabain (100 μ M) clearly decreased the mitogenic activity of IL-2 on T cells, but only moderately reduced the concanavalin A-triggered IL-2 production. These findings were confirmed by Lillehoj and Shevach [52] and studied in great detail by Dornand et al. [53]. Of note, ouabain, up to a concentration of 10 mM and a treatment period of 30 h, did not alter the viability of isolated human lymphocytes despite inhibition of the Na⁺-K⁺-ATPase [54].

In thymocytes, which represent hematopoietic progenitor cells in the thymus, a high concentration of ouabain (5 μ M) inhibited the mitogen-induced proliferation [55]. Also a much lower concentration (100 nM) was able to decrease thymocyte activation *via* reducing the phosphorylation of MAPK p38 and the levels of the nuclear factor of activated T cells (NFAT) c1 [56]. In contrast, ouabain (100 nM) was reported to augment intracellular calcium concentrations and to enhance the expression of CD69, an indicator of thymocyte activation [57]. Moreover, ouabain and glucocorticoids synergistically increased the death of thymocytes [58, 59]. The pathophysiological meaning of these results, however, remains elusive.

Besides T lymphocytes, also B cells were found to be susceptible to ouabain: Milthorp et al. [60] demonstrated that ouabain (10 μ M) inhibits the production of antibodies in splenocytes from pigs and rabbits, but not from mice.

Peripheral blood mononuclear cells

In the late 1990s, the influence of CGs on the cytokine secretion from PBMCs (monocytes and lymphocytes) was investigated in detail. These studies provided an interesting and very important insight into the differential actions of CGs: the influence of ouabain on the cytokine production of PBMCs strongly depends on their activation status. Based on the finding that the activity of the Na⁺-K⁺-ATPase is decreased in PBMCs from rheumatoid arthritis patients [61], Foey et al. [62] showed that mimicking this pathological situation in isolated human PBMCs from healthy donors by treating these cells with ouabain (100 nM) strongly augments IL-1 β and TNF α , but suppresses IL-6 secretion. Thus, an impaired

function of the Na⁺-K⁺-ATPase might be associated with the increased levels of proinflammatory cytokines in patients with rheumatoid arthritis. Matsumori et al. [63] also obtained PBMCs from healthy human donors. In their setting, the expression (mRNA and protein) of all three proinflammatory cytokines, IL-1 β , TNF α , and IL-6, was increased by ouabain (100 nM). After activation of PBMCs with LPS, however, ouabain suppressed the production of IL-6 and TNF α . This was in line with the *in vivo* results: ouabain (1 mg/kg bodyweight) protected mice from the LPS-induced lethal toxicity and reduced the levels of IL-6 and TNF α . In a follow-up study, Matsumori et al. [64] could show that the calcium channel blocker amlodipine blocked the ouabain (1 μ M)-triggered increase of IL-1 α , IL-1 β , and IL-6 in PBMCs from healthy human donors.

In 2008, Ihenetu et al. [65] confirmed the earlier reports on the action of CGs on human PBMCs: Both digoxin (< 100 nM) and the endogenous digoxin-like immunoreactive factors (DLIF) inhibited the LPS-triggered activation of NF κ B and the release of TNF α , IL-6, and IL-8.

Monocytes and macrophages

Sowa and Przewlocki [66] analyzed the release of nitric oxide (NO), an important defense mechanism of the innate immunity, from rat peritoneal macrophages. They found that ouabain (125–500 μ M) does not affect basal NO generation, however, it strongly enhanced the LPS-induced NO formation. In monocytes purified from human PBMCs (healthy donors), ouabain treatment (100 nM) resulted in a decreased expression of the membrane-bound form of CD14 [67]. This effect was based on the activation of the epidermal growth factor receptor and p38 MAPK. Furthermore, ouabain (10 nM) augmented the intracellular calcium concentration, the surface levels of the activation markers CD69, HLA-DR, CD86 and CD80, the expression of IL-1 β and TNF α , and the endocytic activity of these cells [68].

Neutrophils, natural killer cells and dendritic cells

Ward and Becker [69] found that ouabain inhibits the chemotactic migration of rabbit neutrophils, but the IC₅₀ value was quite high, ranging between 1–100 μ M. De Moraes et al. [70] wondered whether natural killer cells are also prone to the action of ouabain. However, they could clearly demonstrate that ouabain does not influence these cells. Nascimento et al. [71] were the first to investigate the action of ouabain (100 nM) on human dendritic cells (DCs). Interestingly, the TNF-induced expression of CD83 and the production of IL-12, two features of DC maturation, were reduced by ouabain, whereas the capability of DCs to activate lymphocytes remained unaltered.

Cardiac Glycosides Decrease the Inflammatory Activation of Endothelial Cells

Endothelial cells are not only a crucial part of the vascular system, but also regulate inflammatory processes, such as the extravasation of immune cells from the blood into the tissue. Up to now, only two studies have dealt with the actions of CGs on endothelial cells in the context of inflammation. Bereta et al. [72] showed that

ouabain activates NF κ B and increases the expression of the NF κ B target genes iNOS and VCAM-1 in murine brain endothelial cells. Furthermore, digitoxin was found to strongly inhibit the IL-1 β -induced expression of an important proinflammatory mediator, the monocyte chemoattractant protein-1 (MCP-1), and of VCAM-1 [73]. Most importantly, this study also showed that digitoxin inhibits the adhesion of monocytes to endothelial cells. Regarding the underlying mechanisms, digitoxin blocked the NF κ B cascade as well as the activation of the extracellular signal-regulated kinase (ERK), but not of p38 MAPK and c-Jun N-terminal kinase (JNK). Moreover, digitoxin augmented the activity of the endothelial NO synthase (eNOS) and prevented the TNF α -caused apoptosis of endothelial cells *via* activation of the phosphatidylinositol-3-kinase (PI3K)/Akt pathway.

Summary and Conclusion

The first report on the anti-inflammatory potential of CGs dates back 50 years. Since then, CGs have been tested in several animal models of inflammation ranging from acute and infection-triggered models to chronic inflammatory disease. Despite this knowledge, a translation into clinical applications has not yet occurred. This might be due to the following unresolved issues that call for further research:

CGs have been found to exert anti-inflammatory actions not only by targeting the Na⁺-K⁺-ATPase, but also *via* different systems, such as ROR γ t. Thus, the role of the ATPase for the anti-inflammatory action of CGs is still unclear. This issue is of high relevance, since the toxic activities of CGs are mainly ascribed to this enzyme. The separation of the cardiovascular/Na⁺-K⁺-ATPase-inhibiting effects from the anti-inflammatory activities has been shown to be feasible. Further approaches into this direction are, thus, of great importance and will strongly advance the field.

Most of the studies analyzed the actions of CGs on leukocytes. A major drawback of these analyses is the fact that highly different concentrations of CGs which do not properly reflect the *in vivo* situation were used. Physiological levels of endogenous CGs range from 10 pM to 1 nM. In contrast, both the plasma concentrations of endogenous CGs during stress conditions and the serum levels of administered CGs during pharmacotherapy are much higher, namely in the range of 100 nM [74, 75]. Also species differences have not yet been considered systematically. Thus, from the clinical perspective, some of the reported effects and presented mechanisms might only be of limited relevance.

Up to now, only one very small and preliminary clinical trial has investigated the anti-inflammatory potential of CGs (in the context of cystic fibrosis). Unfortunately, due to the very limited period of drug treatment, the study could not demonstrate significant benefits of CGs. However, the effort to perform clinical trials is highly appreciated and needs to be expanded urgently.

Interestingly, although the pathogenesis of many types of tumors is closely associated with inflammatory processes, no data about the action of CGs on the interface between cancer and inflammation exist. As a future direction, it would be of great interest to investigate both the pathophysiological role of endogenous

CGs and the pharmacological action of exogenously applied CGs in the context of inflammation-associated tumorigenesis.

Taken together, CGs possess a very interesting anti-inflammatory potential. However, the usage of unmodified CGs seems to be very unlikely due to their narrow therapeutic index. The rational design of CG analogs with decreased cardiac and augmented anti-inflammatory actions is a young but promising research area that will greatly advance the field.

Conflict of Interest

The authors declare no conflict of interest.

References

- [1] Greeff K, Schadewaldt H. Introduction and Remarks on the History of cardiac Glycosides. In: Greeff K, ed. Cardiac Glycosides. Berlin, Heidelberg: Springer; 1981: 1–12
- [2] Lampe D, Lampe H, Berwing K, Parsi RA, Mai I. On the problem of optimum digitalization in man. *Int J Clin Pharmacol Biopharm* 1978; 16: 380–383
- [3] Regal W, Nanut M. Von der Geheimrezeptur zum Herzglykosid. Available at <http://www.springermedizin.at/artikel/2184-von-der-geheimrezeptur-zum-herzglykosid>. Accessed January 5, 2017
- [4] Repke K, Portius HJ. Comparison of the effect of steroid lactones on the transport-ATPase and ion transport of erythrocytes. *Folia Haematol Int Mag Klin Morphol Blutforsch* 1965; 83: 108–118
- [5] Ferguson DW, Berg WJ, Sanders JS, Roach PJ, Kempf JS, Kienzle MG. Sympathoinhibitory responses to digitalis glycosides in heart failure patients. Direct evidence from sympathetic neural recordings. *Circulation* 1989; 80: 65–77
- [6] Haddy FJ. Endogenous digitalis-like factor or factors. *N Engl J Med* 1987; 316: 621–623
- [7] Poston L. Endogenous sodium pump inhibitors: a role in essential hypertension? *Clin Sci (Lond)* 1987; 72: 647–655
- [8] Haber E, Hauptert GT jr. The search for a hypothalamic Na⁺,K⁺-ATPase inhibitor. *Hypertension* 1987; 9: 315–324
- [9] Goto A, Yamada K, Ishii M, Yoshioka M, Ishiguro T, Eguchi C, Sugimoto T. Purification and characterization of human urine-derived digitalis-like factor. *Biochem Biophys Res Commun* 1988; 154: 847–853
- [10] Goto A, Ishiguro T, Yamada K, Ishii M, Yoshioka M, Eguchi C, Shimura M, Sugimoto T. Isolation of a urinary digitalis-like factor indistinguishable from digoxin. *Biochem Biophys Res Commun* 1990; 173: 1093–1101
- [11] Hamlyn JM, Blaustein MP, Bova S, DuCharme DW, Harris DW, Mandel F, Mathews WR, Ludens JH. Identification and characterization of a ouabain-like compound from human plasma. *Proc Natl Acad Sci U S A* 1991; 88: 6259–6263
- [12] Bova S, Blaustein MP, Ludens JH, Harris DW, DuCharme DW, Hamlyn JM. Effects of an endogenous ouabainlike compound on heart and aorta. *Hypertension* 1991; 17: 944–950
- [13] Ferrandi M, Manunta P, Balzan S, Hamlyn JM, Bianchi G, Ferrari P. Ouabain-like factor quantification in mammalian tissues and plasma: comparison of two independent assays. *Hypertension* 1997; 30: 886–896
- [14] Buckalew VM. Endogenous digitalis-like factors: an overview of the history. *Front Endocrinol (Lausanne)* 2015; 6: 49
- [15] Berendes E, Cullen P, Van Aken H, Zidek W, Erren M, Hübschen M, Weber T, Wirtz S, Tepel M, Walter M. Endogenous glycosides in critically ill patients. *Crit Care Med* 2003; 31: 1331–1337
- [16] Rathore SS, Curtis JP, Wang Y, Bristow MR, Krumholz HM. Association of serum digoxin concentration and outcomes in patients with heart failure. *JAMA* 2003; 289: 871–878

- [17] Georgiopolou VV, Kalogeropoulos AP, Giamouzis G, Agha SA, Rashad MA, Waheed S, Laskar S, Smith AL, Butler J. Digoxin therapy does not improve outcomes in patients with advanced heart failure on contemporary medical therapy. *Circ Heart Fail* 2009; 2: 90–97
- [18] Freeman JV, Yang J, Sung SH, Hlatky MA, Go AS. Effectiveness and safety of digoxin among contemporary adults with incident systolic heart failure. *Circ Cardiovasc Qual Outcomes* 2013; 6: 525–533
- [19] Opie LH. Digitalis, yesterday and today, but not forever. *Circ Cardiovasc Qual Outcomes* 2013; 6: 511–513
- [20] Vamos M, Erath JW, Hohnloser SH. Digoxin-associated mortality: a systematic review and meta-analysis of the literature. *Eur Heart J* 2015; 36: 1831–1838
- [21] Chaggar PS, Shaw SM, Williams SG. Is foxglove effective in heart failure? *Cardiovasc Ther* 2015; 33: 236–241
- [22] Ziff OJ, Lane DA, Samra M, Griffith M, Kirchhof P, Lip GY, Steeds RP, Townend J, Kotecha D. Safety and efficacy of digoxin: systematic review and meta-analysis of observational and controlled trial data. *BMJ* 2015; 351: h4451
- [23] Shiratori O. Growth inhibitory effect of cardiac glycosides and aglycones on neoplastic cells: *in vitro* and *in vivo* studies. *Gan* 1967; 58: 521–528
- [24] Stenkvist B, Bengtsson E, Eriksson O, Holmquist J, Nordin B, Westman-Naeser S. Cardiac glycosides and breast cancer. *Lancet* 1979; 1: 563
- [25] Slingerland M, Cerella C, Guchelaar HJ, Diederich M, Gelderblom H. Cardiac glycosides in cancer therapy: from preclinical investigations towards clinical trials. *Invest New Drugs* 2013; 31: 1087–1094
- [26] Alevizopoulos K, Calogeropoulou T, Lang F, Stourmaras C. Na⁺/K⁺ ATPase inhibitors in cancer. *Curr Drug Targets* 2014; 15: 988–1000
- [27] Calderón-Montaño JM, Burgos-Morón E, Orta ML, Maldonado-Navas D, García-Domínguez I, López-Lázaro M. Evaluating the cancer therapeutic potential of cardiac glycosides. *Biomed Res Int* 2014; 2014: 794930
- [28] Durlacher CT, Chow K, Chen XW, He ZX, Zhang X, Yang T, Zhou SF. Targeting Na(+)/K(+) -translocating adenosine triphosphatase in cancer treatment. *Clin Exp Pharmacol Physiol* 2015; 42: 427–443
- [29] Patel S. Plant-derived cardiac glycosides: Role in heart ailments and cancer management. *Biomed Pharmacother* 2016; 84: 1036–1041
- [30] Diederich M, Muller F, Cerella C. Cardiac glycosides: From molecular targets to immunogenic cell death. *Biochem Pharmacol* 2017; 125: 1–11
- [31] Lancaster MC, Vegad JL. Suppression of the early inflammatory response in the sheep by strophanthrin G. *Nature* 1967; 213: 840–841
- [32] De Vasconcelos DI, Leite JA, Carneiro LT, Piuvezam MR, de Lima MR, de Moraes LC, Rumjanek VM, Rodrigues-Mascarenhas S. Anti-inflammatory and antinociceptive activity of ouabain in mice. *Mediators Inflamm* 2011; 2011: 912925
- [33] Leite JA, Alves AK, Galvão JG, Teixeira MP, Cavalcante-Silva LH, Scavone C, Morrot A, Rumjanek VM, Rodrigues-Mascarenhas S. Ouabain modulates zymosan-induced peritonitis in mice. *Mediators Inflamm* 2015; 2015: 265798
- [34] Esposito AL. Digoxin disrupts the inflammatory response in experimental pneumococcal pneumonia. *J Infect Dis* 1985; 152: 14–23
- [35] Jacob PL, Leite JA, Alves AK, Rodrigues YK, Amorim FM, Neris PL, Oliveira MR, Rodrigues-Mascarenhas S. Immunomodulatory activity of ouabain in *Leishmania leishmania amazonensis*-infected Swiss mice. *Parasitol Res* 2013; 112: 1313–1321
- [36] Huh JR, Leung MW, Huang P, Ryan DA, Krout MR, Malapaka RR, Chow J, Manel N, Ciofani M, Kim SV, Cuesta A, Santori FR, Lafaille JJ, Xu HE, Gin DY, Rastinejad F, Littman DR. Digoxin and its derivatives suppress TH17 cell differentiation by antagonizing ROR γ activity. *Nature* 2011; 472: 486–490
- [37] Lee J, Baek S, Lee J, Lee J, Lee DG, Park MK, Cho ML, Park SH, Kwok SK. Digoxin ameliorates autoimmune arthritis via suppression of Th17 differentiation. *Int Immunopharmacol* 2015; 26: 103–111
- [38] Hinshaw SJ, Ogbeifun O, Wandu WS, Lyu C, Shi G, Li Y, Qian H, Gery I. Digoxin inhibits induction of experimental autoimmune uveitis in mice, but causes severe retinal degeneration. *Invest Ophthalmol Vis Sci* 2016; 57: 1441–1447
- [39] Forshammar J, Block L, Lundborg C, Biber B, Hansson E. Naloxone and ouabain in ultralow concentrations restore Na⁺/K⁺-ATPase and cytoskeleton in lipopolysaccharide-treated astrocytes. *J Biol Chem* 2011; 286: 31586–31597
- [40] Forshammar J, Jörneberg P, Björklund U, Westerlund A, Lundborg C, Biber B, Hansson E. Anti-inflammatory substances can influence some glial cell types but not others. *Brain Res* 2013; 1539: 34–40
- [41] Kinoshita PF, Yshii LM, Vasconcelos AR, Orellana AM, Lima Lde S, Davel AP, Rossoni LV, Kawamoto EM, Scavone C. Signaling function of Na,K-ATPase induced by ouabain against LPS as an inflammation model in hippocampus. *J Neuroinflammation* 2014; 11: 218
- [42] Srivastava M, Eidelman O, Zhang J, Paweletz C, Caohuy H, Yang Q, Jacobson KA, Heldman E, Huang W, Jozwick C, Pollard BS, Pollard HB. Digitoxin mimics gene therapy with CFTR and suppresses hypersecretion of IL-8 from cystic fibrosis lung epithelial cells. *Proc Natl Acad Sci U S A* 2004; 101: 7693–7698
- [43] Yang Q, Huang W, Jozwick C, Lin Y, Glasman M, Caohuy H, Srivastava M, Esposito D, Gillette W, Hartley J, Pollard HB. Cardiac glycosides inhibit TNF- α /NF- κ B signaling by blocking recruitment of TNF receptor-associated death domain to the TNF receptor. *Proc Natl Acad Sci U S A* 2005; 102: 9631–9636
- [44] Zeitlin PL, Diener-West M, Callahan KA, Lee S, Talbot CC jr., Pollard B, Boyle MP, Lechtzin N. Digitoxin for airway inflammation in cystic fibrosis: preliminary assessment of safety, pharmacokinetics, and dose finding. *Ann Am Thorac Soc* 2017; 14: 220–229
- [45] Quastel MR, Kaplan JG. Inhibition by ouabain of human lymphocyte transformation induced by phytohaemagglutinin *in vitro*. *Nature* 1968; 219: 198–200
- [46] Quastel MR, Kaplan JG. Lymphocyte stimulation: the effect of ouabain on nucleic acid and protein synthesis. *Exp Cell Res* 1970; 62: 407–420
- [47] Takada Y, Matsuo K, Ogura H, Bai L, Toki A, Wang L, Ando M, Kataoka T. Odoricide A and ouabain inhibit Na⁺/K⁺-ATPase and prevent NF- κ B-inducible protein expression by blocking Na⁺-dependent amino acid transport. *Biochem Pharmacol* 2009; 78: 1157–1166
- [48] Szamel M, Somogyi J, Csukás I, Solymosy F. Effect of ouabain on macromolecular synthesis during the cell cycle in mitogen-stimulated human lymphocytes. *Biochim Biophys Acta* 1980; 633: 347–360
- [49] Szamel M, Resch K. Inhibition of lymphocyte activation by ouabain. Interference with the early activation of membrane phospholipid metabolism. *Biochim Biophys Acta* 1981; 647: 297–301
- [50] Pires V, Harab RC, Olej B, Rumjanek VM. Ouabain effects on activated lymphocytes: augmentation of CD25 expression on TPA-stimulated cells and of CD69 on PHA-and TPA-stimulated cells. *Int J Immunopharmacol* 1997; 19: 143–148
- [51] Stoeck M, Northoff H, Resch K. Inhibition of mitogen-induced lymphocyte proliferation by ouabain: interference with interleukin 2 production and interleukin 2 action. *J Immunol* 1983; 131: 1433–1437
- [52] Lillehoj H, Shevach EM. A comparison of the effects of cyclosporin A, dexamethasone, and ouabain on the interleukin-2 cascade. *J Immunopharmacol* 1985; 7: 267–284
- [53] Dornand J, Favero J, Bonnafous JC, Mani JC. Mechanism whereby ouabain inhibits human T lymphocyte activation: effect on the interleukin 2 pathway. *Immunobiology* 1986; 171: 436–450
- [54] Falciola J, Volet B, Anner RM, Moosmayer M, Lacotte D, Anner BM. Role of cell membrane Na,K-ATPase for survival of human lymphocytes *in vitro*. *Biosci Rep* 1994; 14: 189–204
- [55] Szamel M, Schneider S, Resch K. Functional interrelationship between (Na⁺ + K⁺)-ATPase and lysolecithin acyltransferase in plasma membranes

- of mitogen-stimulated rabbit thymocytes. *J Biol Chem* 1981; 256: 9198–9204
- [56] Rodrigues-Mascarenhas S, Bloise FF, Moscat J, Rumjanek VM. Ouabain inhibits p38 activation in thymocytes. *Cell Biol Int* 2008; 32: 1323–1328
- [57] Rodrigues-Mascarenhas S, Echevarria-Lima J, Fernandes dos Santos N, Rumjanek VM. CD69 expression induced by thapsigargin, phorbol ester and ouabain on thymocytes is dependent on external Ca^{2+} entry. *Life Sci* 2003; 73: 1037–1051
- [58] Mann CL, Bortner CD, Jewell CM, Cidlowski JA. Glucocorticoid-induced plasma membrane depolarization during thymocyte apoptosis: association with cell shrinkage and degradation of the Na^{+}/K^{+} -adenosine triphosphatase. *Endocrinology* 2001; 142: 5059–5068
- [59] Rodrigues-Mascarenhas S, dos Santos NF, Rumjanek VM. Synergistic effect between ouabain and glucocorticoids for the induction of thymic atrophy. *Biosci Rep* 2006; 26: 159–169
- [60] Milthorpe P, Quastel MR, Kaplan JG, Vogelfanger IJ. Effect of the inhibition of cation transport by ouabain on plaque formation by splenic lymphoid cells: time course and dose response. *Cell Immunol* 1974; 14: 128–133
- [61] Maubach K, Foey AD, Hall ND. Impaired activity of thiol-dependent ATPases in rheumatoid mononuclear cell membranes. *Agents Actions* 1993; 39 (Spec No): C107–C109
- [62] Foey AD, Crawford A, Hall ND. Modulation of cytokine production by human mononuclear cells following impairment of Na^{+} , K^{+} -ATPase activity. *Biochim Biophys Acta* 1997; 1355: 43–49
- [63] Matsumori A, Ono K, Nishio R, Igata H, Shioi T, Matsui S, Furukawa Y, Iwasaki A, Nose Y, Sasayama S. Modulation of cytokine production and protection against lethal endotoxemia by the cardiac glycoside ouabain. *Circulation* 1997; 96: 1501–1506
- [64] Matsumori A, Ono K, Nishio R, Nose Y, Sasayama S. Amlodipine inhibits the production of cytokines induced by ouabain. *Cytokine* 2000; 12: 294–297
- [65] Ihenetu K, Espinosa R, de Leon R, Planas G, Perez-Pinero A, Waldbeser L. Digoxin and digoxin-like immunoreactive factors (DLIF) modulate the release of pro-inflammatory cytokines. *Inflamm Res* 2008; 57: 519–523
- [66] Sowa G, Przewlocki R. Ouabain enhances the lipopolysaccharide-induced nitric oxide production by rat peritoneal macrophages. *Immunopharmacology* 1997; 36: 95–100
- [67] Valente RC, Nascimento CR, Araujo EG, Rumjanek VM. mCD14 expression in human monocytes is downregulated by ouabain via transactivation of epithelial growth factor receptor and activation of p38 mitogen-activated protein kinase. *Neuroimmunomodulation* 2009; 16: 228–236
- [68] Teixeira MP, Rumjanek VM. Ouabain affects the expression of activation markers, cytokine production, and endocytosis of human monocytes. *Mediators Inflamm* 2014; 2014: 760368
- [69] Ward PA, Becker EL. Potassium reversible inhibition of leukotaxis by ouabain. *Life Sci* 1970; 9: 355–360
- [70] De Moraes VL, Olej B, de la Rocque L, Rumjanek VM. Lack of sensitivity to ouabain in natural killer activity. *FASEB J* 1989; 3: 2425–2429
- [71] Nascimento CR, Valente RC, Echevarria-Lima J, Fontes CF, de Araujo-Martins L, Araujo EG, Rumjanek VM. The influence of ouabain on human dendritic cells maturation. *Mediators Inflamm* 2014; 2014: 494956
- [72] Bereta J, Cohen MC, Bereta M. Stimulatory effect of ouabain on VCAM-1 and iNOS expression in murine endothelial cells: involvement of NF- κ B. *FEBS Lett* 1995; 377: 21–25
- [73] Jagielska J, Salguero G, Schieffer B, Bavendiek U. Digitoxin elicits anti-inflammatory and vasoprotective properties in endothelial cells: Therapeutic implications for the treatment of atherosclerosis? *Atherosclerosis* 2009; 206: 390–396
- [74] Goto A, Yamada K, Nagoshi H, Terano Y, Omata M. Stress-induced elevation of ouabainlike compound in rat plasma and adrenal. *Hypertension* 1995; 26: 1173–1176
- [75] Gottlieb SS, Rogowski AC, Weinberg M, Krichthen CM, Hamilton BP, Hamlyn JM. Elevated concentrations of endogenous ouabain in patients with congestive heart failure. *Circulation* 1992; 86: 420–425