


## Endoscopic submucosal dissection of a squamous cell carcinoma of the esophagus developing in the area of a previous Heller's myotomy for achalasia



► **Fig. 1** Endoscopic appearance of the esophageal lesion showing: **a, b** very mild vessel irregularities on narrow-band imaging; **c, d** an unstained lesion with the pink-color sign after staining with Lugol dye.


Achalasia is considered a risk factor for esophageal cancer; nevertheless, there is no consensus for any regular screening [1, 2]. A few reports have described early esophageal cancer associated with achalasia that has been resected by endoscopy [3, 4]. Here we report a case of endoscopic submucosal dissection (ESD) for a squamous cell carcinoma that developed in the area of a previous Heller's myotomy.

A 77-year-old woman with achalasia underwent Heller's myotomy in 2015. In

March 2017 she underwent follow-up endoscopy and a squamous cell carcinoma was detected in the lower third of the esophagus. Examination using magnified narrow-band imaging (NBI) revealed very mild vessel irregularities (► **Fig. 1a, b**), but the use of Lugol dye demonstrated an unstained lesion with the pink-color sign (► **Fig. 1c, d**). ESD was successfully performed but the lesion was located on the area of the previous myotomy and there was no circular muscle layer left under the submucosa (► **Video 1**; ► **Fig. 2**).

During the procedure, the scarred tissue under the submucosa maintained the structure of the esophageal wall and no perforation occurred. At the end of the procedure, a 24-mm covered metal stent (Taewoong Medical, Seoul, South Korea) was placed and fixed with two clips to avoid delayed perforation.

A subsequent computed tomography (CT) scan showed neither a pneumomediastinum nor signs of esophageal perforation (► **Fig. 3**). Pathology confirmed an intramucosal squamous cell carcinoma.



► **Fig. 2** Views during the procedure showing: **a** submucosal fibrosis in the area of the previous Heller's myotomy; **b** the tunnel with lower enlargement in the previous myotomy area; **c** muscle fibrosis in the area of the previous myotomy; **d** a stent positioned to cover the area.




► **Video 1** Endoscopic submucosal dissection of an esophageal squamous cell carcinoma found in the area of a previous Heller's myotomy, a covered metal stent is placed at the end of the procedure and is subsequently removed 2 weeks later.

ma that had been resected with free margins. The patient recovered completely, being able to eat on day 1, and was discharged on day 3 following the procedure with no delayed morbidity. The stent was removed after 2 weeks because of pain, which disappeared after stent removal.

To our knowledge, this is the first case of ESD for an early esophageal cancer in the area of a previous Heller's myotomy. ESD has been shown to be a safe and effective procedure for early esophageal cancer [5], even in a previously treated achalasia patient [3]. Previous myotomy should not be considered a contraindication to an ESD procedure in such patients, but a prophylactic stent can be placed to reduce the risk of complications.

Endoscopy\_UCTN\_Code\_TTT\_1AO\_2AI



► **Fig. 3** Appearances following the procedure: **a, b** on chest computed tomography, showing a 24-mm covered metal stent (white arrows) fixed with two clips (black arrows) and no evidence of either a pneumomediastinum or esophageal perforation; **c** on endoscopy after stent removal.

### Competing interests

None

### The authors

**Dario Biasutto**<sup>1,2</sup>, **Jérôme Rivory**<sup>3</sup>, **Sabine Roman**<sup>1,2,4</sup>, **François Mion**<sup>1,2,4</sup>, **Jean-Christophe Saurin**<sup>3</sup>, **Thierry Ponchon**<sup>3,4</sup>, **Mathieu Pioche**<sup>3,4</sup>

- 1 Digestive Physiology, Hospices Civils de Lyon, Hôpital Edouard Herriot, Lyon, France
- 2 Digestive Physiology, Université de Lyon 1, Lyon, France
- 3 Department of Endoscopy and Gastroenterology, Pavillon L, Hôpital Edouard Herriot, Lyon, France
- 4 Inserm U1032, LabTAU, Lyon, France

### Corresponding author

**Mathieu Pioche, MD**

Gastroenterology and Endoscopy, Hôpital Edouard Herriot, 5, place d'Arsonval, 69437 Lyon cedex 03, France  
 Fax: +33-4-72110147  
 mathieu.pioche@chu-lyon.fr

## References

- [1] Ravi K, Geno DM, Katzka DA. Esophageal cancer screening in achalasia: is there a consensus? *Dis Esophagus* 2015; 28: 299–304
- [2] Vaezi MF1, Pandolfino JE, Vela MF. ACG clinical guideline: diagnosis and management of achalasia. *Am J Gastroenterol* 2013; 108: 1238–1249
- [3] Ota M, Narumiya K, Kudo K et al. Incidence of esophageal carcinomas after surgery for achalasia: usefulness of long-term and periodic follow-up. *Am J Case Rep* 2016; 17: 845–849
- [4] Ohkura Y, Iizuka T, Kikuchi D et al. Endoscopic submucosal dissection for early esophageal cancer associated with achalasia. *Turk J Gastroenterol* 2013; 24: 161–166
- [5] Kitagawa Y, Suzuki T, Hara T et al. Safety and efficacy of endoscopic submucosal dissection using IT knife nano with clip traction method for early esophageal squamous cell carcinoma. *Surg Endosc*. doi:10.1007/s00464-017-5703-x

### Bibliography

DOI <https://doi.org/10.1055/s-0043-121565>  
Published online: 14.11.2017  
*Endoscopy* 2018; 50: E38–E41  
© Georg Thieme Verlag KG  
Stuttgart · New York  
ISSN 0013-726X

### ENDOSCOPY E-VIDEOS <https://eref.thieme.de/e-videos>



*Endoscopy E-Videos* is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at <https://mc.manuscriptcentral.com/e-videos>