


## Endoscopic mucosal resection of rectal squamous cell papilloma



► **Fig. 1** Endoscopic view of squamous cell papilloma of the rectum using white light (a,c,d) and narrow-band imaging (b,e).


A 22-year-old male patient presented with a 3-month history of persistent, colicky, lower abdominal pain. He was referred for colonoscopy. Total colonoscopy showed multiple small-to-medium sized pale pink, nonulcerated, polypoid lesions in the distal part of the rectum, occupying most of the circumference (► **Fig. 1**).

The patient's medical history was not significant. After discussing his sexual history, he mentioned having a male partner for 2 years. Basic investigations were carried out, including human immunodeficiency virus serology; they were all nega-




► **Fig. 2** Complete endoscopic mucosal resection of squamous cell papilloma.





► **Fig. 3** Histopathological examination. **a, b** Hyperplastic stratified squamous epithelium with keratosis, vascular cores (**a**, arrow), and epithelioid (**b**, arrow). **c** Squamous cell papilloma (SCP) with mild atypia (arrow). **d** SCP with focal moderate atypia (arrow).



► **Video 1** Endoscopic mucosal resection of rectal squamous cell papilloma.



tive. After discussing the treatment options with the patient, endoscopic mucosal resection of the lesions was performed (► **Fig. 2**, ► **Video 1**). Histopathological examination showed squamous cell papillomatosis of the rectum, with mild-to-moderate atypia (► **Fig. 3**). Isolation of the human papilloma virus (HPV) from the specimen was not possible. Squamous cell papilloma (SCP) is a benign lesion that is a result of infection with HPV [1]. HPV-associated diseases are sexually transmitted and mainly affect the genitals, causing wart formation [2]. SCP arises from the stratified squamous epithelium of the skin, lip, oral cavity, tongue, pharynx, larynx, esophagus, cervix, vagina or anal canal [3]. SCP of the gastrointestinal tract is

mainly localized to the oropharynx, esophagus, and anal canal [4]. The current case was an extremely rare case of rectal SCP.

Endoscopy\_UCTN\_Code\_CCL\_1AD\_2AC

### Competing interests

None

### The authors

**Mohamed Nabil Alkady<sup>1</sup>, Shaimaa Elkholy<sup>2</sup>**

- 1 Tropical Department, Faculty of Medicine, Cairo University, Cairo, Egypt
- 2 Internal Medicine Department, Faculty of Medicine, Cairo University, Cairo, Egypt

### Corresponding author

**Shaimaa Elkholy, MD**

Internal Medicine Department, Cairo University, 531, 17th street, 5th district 6 of October City, Cairo 00202, Egypt  
Fax: +20-2-37493563  
shuma50082@gmail.com

### References

- [1] New Zealand Dermatological Society. Squamous cell papilloma. New Zealand Dermatological Society. Available from: <https://www.dermnetnz.org/topics/squamous-cell-papilloma/> Accessed November 2007
- [2] Ljubojevic S, Skerlev M. HPV-associated diseases. *Clin Dermatol* 2014; 32: 227 – 234
- [3] Gerada J, Savic A, Vassallo M. Squamous papilloma of the anal canal. *Endoscopy* 2013; 45: E42 – E43
- [4] Karantanis D, Allen-Auerbach M, Czernin J. Squamous papilloma of the oral cavity and oropharynx on 18F-FDG PET/CT imaging. *Clin Nucl Med* 2012; 37: e98 – e99

### Bibliography

DOI <https://doi.org/10.1055/s-0043-123819>  
Published online: 19.1.2018  
*Endoscopy* 2018; 50: E83–E85  
© Georg Thieme Verlag KG  
Stuttgart · New York  
ISSN 0013-726X

### ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



*Endoscopy E-Videos* is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at <https://mc.manuscriptcentral.com/e-videos>