

# The Combination of Fosfomycin, Metronidazole, and Recombinant Human Granulocyte-Macrophage Colony-Stimulating Factor is Stable *in vitro* and Has Maintained Antibacterial Activity

## Authors

Siv Fonnes<sup>1</sup>, Barbara Juliane Holzknacht<sup>2</sup>, Lærke Smidt Gasbjerg<sup>3,4</sup>, Johan Juhl Weisser<sup>5</sup>, Hanne Wiese Hallberg<sup>2</sup>, Magnus Arpi<sup>2</sup>, Jacob Rosenberg<sup>1</sup>

## Affiliations

- 1 Centre for Perioperative Optimisation, Department of Surgery, Herlev and Gentofte Hospital, University of Copenhagen, Herlev, Denmark
- 2 Department of Clinical Microbiology, Herlev and Gentofte Hospital, University of Copenhagen, Herlev, Denmark
- 3 Novo Nordisk Foundation Center for Basic Metabolic Research and Department of Biomedical Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen N, Denmark
- 4 Center for Diabetes Research, Herlev and Gentofte Hospital, University of Copenhagen, Hellerup, Denmark
- 5 Department of Pharmacy, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen Ø, Denmark

## Key words

Antibacterial drugs, inflammation, chemistry

received 13.09.2017

accepted 28.11.2017

## Bibliography

DOI <https://doi.org/10.1055/s-0043-123933>

Published online: 19.12.2017

Drug Res 2018; 68: 349–354

© Georg Thieme Verlag KG Stuttgart · New York

ISSN 2194-9379

## Correspondence

Siv Fonnes

Centre for Perioperative Optimisation, Department of Surgery

Herlev and Gentofte Hospital, University of Copenhagen

Herlev Ringvej 75

DK-2730 Herlev

Denmark

Tel.: +45/23/21 65 34, Fax: +45/38/68 37 72

siv.fonnes@gmail.com

## ABSTRACT

**Background** Treatment of secondary peritonitis includes surgery and antimicrobial agents. Antimicrobial agents are often administered intravenously, however, the alternative route intraperitoneal administration could be considered. Investigations must be conducted prior to clinical application. Therefore, we aimed to investigate the combination of fosfomycin, metronidazole, and recombinant human granulocyte-macrophage colony-stimulating factor with regard to its chemical properties and the solution's stability. In addition, the antibacterial effect of the mixed drug solution was compared with the effect of the individual antibacterial agents.

**Methods** The drugs were mixed to an aqueous solution. Basic chemical investigations of pH, precipitation, and calculated osmolarity of the drug combination were conducted. Fosfomycin and metronidazole's chemical stability was investigated using High Pressure Liquid Chromatography-Mass Spectrometry. Microbiological investigations using the agar cup method were carried out to measure the antibacterial effect of fosfomycin and metronidazole.

**Results** The aqueous solution of the combination of the three drugs had a pH of 7.46–7.62, which was stable during 24 h, was without precipitation, and had a calculated osmolarity of 293 mOsm/l. High Pressure Liquid Chromatography-Mass Spectrometry found stable concentrations of fosfomycin and metronidazole both alone and in combination during 24 h. The antibacterial effect of the drug combination solution was similar to the antibacterial effects of fosfomycin and metronidazole alone.

**Conclusion** The drug combination had neutral and stable pH, was iso-osmotic, and had stable concentrations during 24 h of storage. The antibacterial effect of fosfomycin and metronidazole were not altered when the drugs were mixed.

## Introduction

Antimicrobial therapy for intraabdominal infections, either as prophylaxis or treatment, can be administered by different routes. Intravenous administration is the most frequently recommended route of administration [1, 2], however, local administration could be superior to systemic administration. This has been employed in e. g. intraperitoneal administration in peritoneal dialysis-associated peritonitis [3].

In peritonitis due to perforation of a hollow organ, which is termed secondary peritonitis, the infection is usually polymicrobial [2]. The recommended treatment is source control [4] together with combination regimens of antimicrobial agents [2]. We propose that combination therapy with fosfomycin, metronidazole, and recombinant human granulocyte-macrophage colony-stimulating factor (rhGM-CSF) administered intraoperatively and intraperitoneally could effectively treat secondary, infectious peritonitis. The drugs have not previously been administered in combination. However, individual administration of each drug intraperitoneally has been reported [5–10]. Fosfomycin (cis-(1R,2S)-Epoxypropylphosphonic acid [11]) is a phosphonic antibiotic agent [12] with low molecular weight of 138 Da [13]. It enters the bacteria and inhibits the bacterial cell wall synthesis through an irreversible inhibition of the enzyme enolpyruvyl transferase [14]. Therefore, it exhibits bactericidal activity on both Gram-positive and Gram-negative aerobic bacteria in the growth phase. Metronidazole (1H-Imidazole-1-ethanol, 2-methyl-5-nitro) is a nitroimidazole antimicrobial agent, which has a bactericidal action on anaerobic bacteria and certain protozoa [15]. It has a molecular weight of 171 Da [16]. Its mechanism of action is not fully understood. However, intermediate metabolites of metronidazole are considered cytotoxic and interact with bacterial DNA, which results in destabilisation [15]. The third compound, rhGM-CSF, is a glycoprotein and cytokine that is produced by cells of the immune system such as T lymphocytes, monocytes, fibroblasts, and endothelial cells [17]. The main structure of rhGM-CSF is a four  $\alpha$ -helix bundle. It works by promoting growth, differentiation, and maturation of macrophages and neutrophil granulocytes and stimulate their activities [17–19]. Thereby, it complements antibiotic treatment by enhancement of the patient's immune response. Previous studies found rhGM-CSF to improve the clinical outcomes of both patients with abdominal sepsis [20] and patients with sepsis admitted to the intensive care unit [21].

Fluids or drugs administered intraperitoneally should preferably be physiological in their constitution prior to clinical application [22]. Therefore, the aim of this study was to investigate the combination of fosfomycin, metronidazole and rhGM-CSF regarding to their chemical properties as an aqueous solution and the antibacterial activity.

## Materials and Methods

### Drugs

The drugs used in this investigation were the following: fosfomycin 2 g (Infectofos, Infectopharm, Germany) dissolved in sterile water for injections (Sterilt Vand "SAD", Denmark), metronidazole 5 mg/ml (Metronidazole "B. Braun", Denmark), and rhGM-CSF

0.25  $\mu\text{g}/\mu\text{l}$  (Repomol, Reponex Pharmaceuticals, Denmark). We used a total of 4 g fosfomycin, 1 g metronidazole, and 50  $\mu\text{g}$  rhGM-CSF. To optimise the parameters for High Pressure Liquid Chromatography-Mass Spectrometry (HPLC-MS) pure analytical standards were used. These were fosfomycin sodium (Sigma-Aldrich, Glostrup, Denmark) and metronidazole (Sigma-Aldrich, Glostrup, Denmark). We used the internal standards (IS) ethylphosphonic acid (Sigma-Aldrich, Glostrup, Denmark) and metronidazole- $d_4$  (Toronto Research Chemicals, ON, Canada).

### Basic chemical investigations

The aims were to determine the pH of the drugs individually and in combination, and to evaluate if precipitations were present.

A pH-meter (PHM210 Standard pH Meter, Radiometer analytical SAS, Denmark) was calibrated with standard solutions of pH 4 and 10 before use. We dissolved 2 g fosfomycin in 50 ml water for injections. This step was repeated. We measured pH and collected photographic documentation. The two fosfomycin solutions were then mixed and 200 ml water for injection was added. Hereafter, we measured pH and collected photographic documentation. 0.2 ml of rhGM-CSF (50  $\mu\text{g}$ ) was added to the 300 ml of fosfomycin solution and gently mixed. We measured pH and collected photographic documentation. Lastly, a total of 200 ml metronidazole 5 mg/ml (1 g) was gathered and pH was measured and photographic documentation was collected. This was mixed with the 300.2 ml solution consisting of fosfomycin and rhGM-CSF and gently mixed resulting in 500.2 ml solution. We measured pH and collected photographic documentation. The abovementioned protocol was conducted in duplicate. The investigations described in the following section were conducted once. We kept 25 ml of the solution at room temperature (20–25 °C). We measured pH and collected photographic documentation after 30 min, 1, 2, 4, 8, and 24 h. The remaining portion of the solution was kept at 37 °C (simulating the body temperature). We measured pH and collected photographic documentation after 30 min, 1, 2, 4, 8, and 24 h.

The theoretical osmolarity was calculated based on the available information of the drugs used.

### Chemical stability

The aim was to investigate whether the concentration of fosfomycin and metronidazole alone and in combination was stable during a time period of 24 h at 37 °C.

Analytical standards dissolved in methanol were used to optimise and develop the methods for HPLC-MS (AB Sciex Qtrap 4500, CA, USA). For both fosfomycin and metronidazole a Kinetex 2.6  $\mu\text{m}$  Biphenyl 100 A, 50  $\times$  2.1 mm column (Phenomenex, USA) was used. In both methodologies mobile phase A consisted of MilliQ water with 0.1 % formic acid (v/v) and mobile phase B consisted of acetonitrile with 0.1 % formic acid (v/v). For metronidazole the injection volume was 0.5  $\mu\text{l}$ , autosampler tray temperature was set to 10 °C, and the column oven was set to 35 °C. A binary 1290 Agilent Infinity System performed a gradient flow at 350  $\mu\text{l}/\text{min}$ . The gradient elution was maintained at 2 % B from 0–1.3 min, and 2–98 % B from 1.3–1.8 min where it was held for 2 min before equilibrating the column again. The retention time was 2.15 min and the total method was 7.5 min long. Metronidazole was analysed in positive mode where source parameters were set to curtain gas (40 l/min),

collision gas (medium), ionspray voltage (5000 V), temperature (400 °C), ion source gas 1 (30 l/min), and 2 (30 l/min). The Q1 parent ion and Q2 daughter ions were 172.0, 128.0, and 110.9 m/z, respectively. For both 128.0 m/z and 110.9 m/z the compound parameters were set to declustering potential (60 V), entrance potential (10 V), collision energy (40 V), cell exit potential (11 V), and the dwell time was 50 msec. For fosfomycin an isocratic method was applied with 10 % B with a flow of 300 µl. The injection volume was 0.1 µl, tray temperature 8 °C and the column oven was 30 °C. The retention time was 0.49 min and the total method was 1.5 min long. Fosfomycin was analysed in negative mode where source parameters were set to curtain gas (40 l/min), collision gas (medium), ionspray voltage (-4500 V), temperature (450 °C), ion source gas 1 (30 l/min), and 2 (30 l/min). The Q1 parent ion and Q2 daughter ions were 136.5, 78.7, and 62.8 m/z, respectively. For both 78.7 m/z and 62.8 m/z the compound parameters were set to declustering potential (-60 V), entrance potential (-6 V), and the dwell time was 150 ms. Furthermore, the collision energy (-33 V and -18 V) and cell exit potential (-7 V and -6 V) were 78.7 m/z and 62.8 m/z, respectively. Linearity and injection variation was tested within the working range: fosfomycin 2–10 mg/ml and metronidazole 0.95–2.25 mg/ml. The chemical stability experiment was conducted by four different prepared solutions at 20 °C: fosfomycin 8 mg/ml alone, metronidazole 2 mg/ml alone, rhGM-CSF  $1 \times 10^{-4}$  mg/ml alone, and combination; fosfomycin 8 mg/ml, metronidazole 2 mg/ml, and rhGM-CSF  $1 \times 10^{-4}$  mg/ml. After preparation, all solutions were placed in a dark room at 37 °C. Hereafter, samples of 100 µl were collected at 0, ½, 1, 2, 4, 8, and 24 h and immediately frozen to -80 °C where they were kept until analysis. When samples were thawed for analysis 50 µl IS (10 ng/µl) was added, then diluted with 850 µl MilliQ water and transferred to 1.5 ml LC vials. All samples were collected in triplicates and data were processed in MultiQuant 3.0 (AB Sciex, CA, USA). Peak areas at time zero was defined as 2 mg/ml and 8 mg/ml for metronidazole and fosfomycin, respectively. The concentration in the following time series was determined by comparing the peak area with time zero. Thus, we compared if fosfomycin or metronidazole in either of the setups (alone or in combination) differed significantly during the 24 h.

## Microbiological in vitro investigations

We investigated the antibacterial effect with the agar cup method, which is a diffusion based method to investigate antimicrobial activity [23, 24]. The agar cup is a cylindrical hole that is punched into an agar plate, which has been pre-inoculated with a microorganism. A fluid containing the tested agent is pipetted into the hole and the agar plate incubated. The growth inhibition zone correlates with the antimicrobial activity of the substance tested.

The antibacterial effect of fosfomycin was investigated with the bacteria strains *Escherichia coli* ATCC 25922 and *Enterococcus faecalis* ATCC 29212 on Mueller-Hinton agar plates (Oxoid, Thermo Fisher Scientific Inc., Denmark) supplemented with 25 µg/ml glucose-6-phosphate (Sigma Aldrich, Brøndby, Denmark) under aerobic conditions. The antibacterial effect of metronidazole was investigated with *Bacteroides fragilis* ATCC 25285 and *Bacteroides thetaio-taomicron* ATCC 29741 on Brucella Agar plates (Becton Dickinson, Heidelberg, Germany) under anaerobic conditions. Plates were in-

oculated with the test strain using an automatic plate rotator. For *E. coli* and *E. faecalis*, an inoculum of 0.5 McFarland was used and for *B. fragilis* and *B. thetaio-taomicron* an inoculum of 1 McFarland was used. Then, the plates were dried and two cylinders were punched into each agar plate. One of the holes was filled with a solution of the investigated drug alone in sterile water ("control solution" containing either 200 µg fosfomycin or 5 µg metronidazole) and the other hole was filled with the combination of the drugs ("combination solution"). For fosfomycin, 200 µg in a volume of 40 µl was pipetted in a 4 mm diameter agar cup and for metronidazole, 5 µg in a volume of 18 µl were pipetted in a 3 mm agar cup. In the control solution, the same amount of the investigated drug was used and the other two drugs were added. All plates were prepared as quintuples, which resulted in 20 plates in total. The plates with *E. coli* and *E. faecalis* were incubated overnight in ambient air at 35 °C (19 h and 19 min). The plates with *B. fragilis* and *B. thetaio-taomicron* were incubated overnight in an anaerobic chamber at 35 °C (18 h and 54 min).

Inhibition zone diameters were read at the point of complete inhibition as judged by the naked eye by an experienced biomedical laboratory scientist with the plate held around 30 cm from the eye. Isolated colonies (seen with *E. coli*) within the inhibition zone were ignored. A negative control without the active agent had been performed in a pilot trial and showed that there was no inhibition of growth.

## Statistical methods

For the investigations of chemical stability with HPLC-MS, it was tested if peak area in time zero was significantly different from time 24 h using unpaired t-test. For the microbiological investigations the inhibition zone diameters of the two solutions ("combination solution" and "control solution") were compared. As data was not normally distributed, the median of the two solutions' diameter was calculated due to the distribution of data. Inter-test variability was evaluated with a Mann-Whitney-U-test of the diameters due to the distribution of data. A p-value < 0.05 was considered statistically significant.

## Results


The basic investigations showed that the combination of fosfomycin, metronidazole, and rhGM-CSF resulted in a neutral pH of 7.46–7.62 (► **Table 1**). The pH remained stable at both room temperature and 37 °C during 24 h (► **Fig. 1**). Furthermore, no precipitation was present at any stage of the combination of drugs (► **Table 1**). The osmolarity was theoretically calculated to 293 mOsm/l.

The HPLC-MS investigations found a stable concentration of both fosfomycin and metronidazole over time (► **Fig. 2**). There was no difference at 24 h between the measured concentration of the drugs alone or in combination for neither fosfomycin (p = 0.5) nor metronidazole (p = 0.9).

The results of the microbiological in vitro investigations are presented in ► **Fig. 3**. A low intra-test variability was found with a range of ± 1 mm from median for each series. The medians of the inhibition zones were similar for the "combination solution" and the "control solution" in all the bacteria strains (► **Fig. 3**).

► **Table 1** An overview of the drug content in the different solutions including the concentrations, the pH, and an evaluation of precipitations at room temperature. rhGM-CSF: recombinant human granulocyte-macrophage colony-stimulating factor.

| Drug content and concentration, mg/ml |                     |               | Precipitation | pH        |
|---------------------------------------|---------------------|---------------|---------------|-----------|
| Fosfomycin                            | rhGM-CSF            | Metronidazole |               |           |
| 40                                    |                     |               | No            | 7.95-7.99 |
| 13                                    |                     |               | No            | 7.84-7.90 |
|                                       | 0.25                |               | No            | 7.32-7.39 |
|                                       |                     | 5.0           | No            | 5.04-5.17 |
| 13                                    | $1.7 \cdot 10^{-4}$ |               | No            | 7.84-8.01 |
| 8.0                                   | $1.0 \cdot 10^{-4}$ | 2.0           | No            | 7.46-7.62 |




► **Fig. 1** The pH of the drug combination of fosfomycin, metronidazole and recombinant human granulocyte-macrophage colony-stimulating factor drugs kept at 20–25°C (black squares), and 37°C (grey diamonds) over a time period of 24 h.

## Discussion

This investigation found that the combination of fosfomycin, rhGM-CSF, and metronidazole had a neutral pH and was without precipitation. The concentrations of both fosfomycin and metronidazole measured by HPLC-MS were stable during 24 h. Finally, the investigation of the *in vitro* antibacterial activity by the agar cup method showed that the antibacterial effect of fosfomycin and metronidazole was maintained in the drug combination.

Precipitation has previously been reported for the combination of GM-CSF and other drugs [25]. Sagramostim, which is GM-CSF produced in yeast, was found visually incompatible with antibiotic agents such as ampicillin and ceftazidime. However, visual compatibility was found for sagramostim and metronidazole. We also found rhGM-CSF to be visually compatible when mixed with fosfomycin and metronidazole, thus the drugs can be mixed.

We found that the drug combination had a neutral pH. A neutral pH is of importance, when administrating drugs intraperitoneally. Low pH of 5 caused discomfort at administration in peritoneal dialysis [26], and discomfort diminished when the pH was neutralised [22, 27, 28]. Low pH of the early dialysis fluids has been suspected to cause chronic alterations of the mesothelium of the peritoneal membrane [29]. These alterations were not present when using biocompatible fluids with neutral pH [30, 31]. There-



► **Fig. 2** Concentrations of fosfomycin (dots) and metronidazole (triangles) alone (grey) and mixed (black) at 37°C during a time period of 24 h measured with High Liquid Chromatography-Mass Spectrometry. The dotted line represents the baseline concentration of each drug. p: p-value.

fore, the pH of 7.46–7.62 found in this investigation is not expected to cause neither discomfort for the patient nor alterations of the peritoneal membrane when the drug combination is administered intraperitoneally.


The *in vitro* inhibitory effect of fosfomycin on aerobic bacteria and metronidazole on anaerobic bacteria was maintained after the drugs were mixed. This is the first investigation on the antibacterial activity of the drug combination. However, previous studies confirmed the preserved antibacterial activity and stability of fosfomycin in different peritoneal dialysis fluids with bioassays [32, 33]. Furthermore, our HPLC-MS analysis confirmed that degradation was neither present for the drugs alone nor in the drug combination solutions during 24 h. Previously, only an unchanged concentration of fosfomycin after mixture with four different peritoneal fluids has been confirmed with HPLC-MS analysis [32].

The strength of these investigations was that all outcomes were predefined in a study protocol, and intra-test variability of the agar cup method was assessed by quintuples to validate interpretation of the test. However, there are some limitations. We do not know the exact limit of detection of the comparative analysis of the antibacterial effect. Since, the results were unambiguous and antagonism of these two drug classes is not expected, we believe that this investigation is adequate to ensure preserved antibacterial activity when the drugs are mixed. Furthermore, *in vitro* studies cannot predict the exact response *in vivo*.

In conclusion, we found that the drug combination of fosfomycin, metronidazole and rhGM-CSF in a specific solution could be mixed without precipitation, had neutral pH, and was iso-osmotic. The concentrations of fosfomycin and metronidazole did not change during 24 h, and the antibacterial activity of the two antibacterial agents was not impaired by their combination.

## Acknowledgements

None.



► **Fig. 3** Antibacterial effect of fosfomycin (panel A) and metronidazole (panel B). Inhibition zone diameters in millimeters (mm) are shown for “combination solution” (black dots) and the “control solution” (grey dots) of each of the five plates for the four different strains: *Escherichia coli*, *Enterococcus faecalis*, *Bacteroides fragilis*, and *Bacteroides thetaiotaomicron*. The punctuated line represents the medians, which were identical for the “combination solution” and the “control solution”. p: p-value.

## Conflict of Interest

This study was funded by a grant from Reponex Pharmaceuticals ApS to the Department of Surgery, Herlev Hospital. The company did not have any influence on the study design, study conduct, or writing of the manuscript.

## References

- [1] O'Neal PB, Itani KMF. Antimicrobial formulation and delivery in the prevention of surgical site infection. *Surg Infect (Larchmt)* 2016; 17: 275–285
- [2] Hasper D, Schefold JC, Baumgart DC. Management of severe abdominal infections. *Recent Pat Antiinfect Drug Discov* 2009; 4: 57–65
- [3] Ballinger AE, Palmer SC, Wiggins KJ et al. Treatment for peritoneal dialysis-associated peritonitis. *Cochrane Database Syst Rev* 2014; 4: CD005284
- [4] Malangoni MA, Inui T. Peritonitis – the Western experience. *World J Emerg Surg* 2006; 1: 25
- [5] Tobudic S, Matzneller P, Stoiser B et al. Pharmacokinetics of intraperitoneal and intravenous fosfomycin in automated peritoneal dialysis patients without peritonitis. *Antimicrob Agents Chemother* 2012; 56: 3992–3995
- [6] Saha SK. Efficacy of metronidazole lavage in treatment of intraperitoneal sepsis. A prospective study. *Dig Dis Sci* 1996; 41: 1313–1318
- [7] Saha SK. Peritoneal lavage with metronidazole. *Surg Gynecol Obstet* 1985; 160: 335–338
- [8] El-Sefi TA, El-Awady HM, Shehata MI et al. Systemic plus local metronidazole and cephazolin in complicated appendicitis: A prospective controlled trial. *J R Coll Surg Edinb* 1989; 34: 13–16
- [9] Selgas R, Fernández de Castro M, Jiménez C et al. Immunomodulation of peritoneal macrophages by granulocyte-macrophage colony-stimulating factor in humans. *Kidney Int* 1996; 50: 2070–2078
- [10] Toner GC, Gabrilove JL, Gordon M et al. Phase I trial of intravenous and intraperitoneal administration of granulocyte-macrophage colony-stimulating factor. *J Immunother Emphasis Tumor Immunol* 1994; 15: 59–66
- [11] Drugbank. <http://www.drugbank.ca/drugs/DB00828> last accessed 31-08-2017)
- [12] Raz R. Fosfomycin: an old-new antibiotic. *Clin Microbiol Infect* 2012; 18: 4–7
- [13] PubChem. <https://pubchem.ncbi.nlm.nih.gov/compound/446987#section=2D-Structure> (last accessed 31-08-2017)
- [14] Kahan FM, Kahan JS, Cassidy PJ et al. The mechanism of action of fosfomycin (phosphonomycin). *Ann N Y Acad Sci* 1974; 235: 364–386
- [15] Lamp KC, Freeman CD, Klutman NE et al. Pharmacokinetics and pharmacodynamics of the nitroimidazole antimicrobials. *Clin Pharmacokinet* 1999; 36: 353–373
- [16] Drugbank. <http://www.drugbank.ca/drugs/DB00916> (last accessed 31-08-2017)
- [17] Liles WC, Van Voorhis WC. Review: Nomenclature and biologic significance of cytokines involved in inflammation and the host immune response. *J Infect Dis* 1995; 172: 1573–1580

- [18] Lieschke GJ, Burgess AW. Granulocyte colony-stimulating factor and granulocyte-macrophage colony-stimulating factor (1). *N Engl J Med* 1992; 327: 28–35
- [19] Weisbart RH, Gasson JC, Golde DW. Physiology of granulocyte and macrophage colony-stimulating factors in host defense. *Hematol Oncol Clin North Am* 1989; 3: 401–409
- [20] Orozco H, Arch J, Medina-Franco H et al. Molgramostim (GM-CSF) associated with antibiotic treatment in nontraumatic abdominal sepsis: A randomized, double-blind, placebo-controlled clinical trial. *Arch Surg* 2006; 141: 150–154
- [21] Presneill JJ, Harris T, Stewart AG et al. A randomized phase II trial of granulocyte-macrophage colony-stimulating factor therapy in severe sepsis with respiratory dysfunction. *Am J Respir Crit Care Med* 2002; 166: 138–143
- [22] Yamamoto T, Sakakura T, Yamakawa M et al. Clinical effects of long-term use of neutralized dialysate for continuous ambulatory peritoneal dialysis. *Nephron* 1992; 60: 324–329
- [23] Rose SB, Miller RE. Studies with the agar cup-plate method: I. A standardized agar cup-plate technique. *J Bacteriol* 1939; 38: 525–537
- [24] Bang J. Zone formation in the agar-cup method for determining resistance to antibiotics. *Acta Pathol Microbiol Scand Suppl* 1956; 39: 192–193
- [25] Trissel LA, Bready BB, Kwan JW et al. Visual compatibility of sargramostim with selected antineoplastic agents, anti-infectives, or other drugs during simulated Y-site injection. *Am J Hosp Pharm* 1992; 49: 402–406
- [26] Prakash J, Tripathi K, Singh M. Complications of intermittent peritoneal dialysis. *J Assoc Physicians India* 1990; 38: 906–908
- [27] Rippe B, Simonsen O, Wieslander A et al. Clinical and physiological effects of a new, less toxic and less acidic fluid for peritoneal dialysis. *Perit Dial Int* 1997; 17: 27–34
- [28] Mactier RA, Sprosen TS, Gokal R et al. Bicarbonate and bicarbonate/lactate peritoneal dialysis solutions for the treatment of infusion pain. *Kidney Int* 1998; 53: 1061–1067
- [29] Williams JD, Craig KJ, Topley N et al. Morphologic changes in the peritoneal membrane of patients with renal disease. *J Am Soc Nephrol* 2002; 13: 470–479
- [30] Williams JD, Topley N, Craig KJ et al. The Euro-Balance Trial: the effect of a new biocompatible peritoneal dialysis fluid (balance) on the peritoneal membrane. *Kidney Int* 2004; 66: 408–418
- [31] Ayuzawa N, Ishibashi Y, Takazawa Y et al. Peritoneal morphology after long-term peritoneal dialysis with biocompatible fluid: Recent clinical practice in Japan. *Perit Dial Int* 2012; 32: 159–167
- Kussmann M, Baumann A, Hauer S et al. Compatibility of fosfomycin with different commercial peritoneal dialysis solutions. *Eur J Clin Microbiol Infect Dis*. 2017; 36: 2237–2242
- [32] Quentin C, Bouchet JJ, Gaillard I et al. Stability of fosfomycin and quinolones in peritoneal dialysis solution. *J Antimicrob Chemother* 1990; 25: 878–880