



What Factors Affect Pain Tolerance during Hysteroscopy?

Quais fatores afetam a tolerância à dor durante a histeroscopia?

Cássia Fernanda dos Santos Rosa¹ Marianna Marques Rodrigues Dourado¹
Rebecca Schuster Dorea Leite¹ Laís Viana Aragão Almeida¹ Brenda Lima Meireles Martins¹
Johnnatas Mikael Lopes²

¹Universidade Tiradentes, Aracaju, SE, Brazil

²Universidade Federal do Vale São Francisco, Paulo Afonso, BA, Brazil

Address for correspondence Johnnatas Mikael Lopes, Universidade Federal do Vale São Francisco, Avenida da Amizade, s/n°, 48605-780, Paulo Afonso, BA, Brazil (e-mail: johnnataslopes@univasf.edu.br).

Rev Bras Ginecol Obstet 2023;45(8):e499–e500.

Hysteroscopy is considered the gold standard for the evaluation and management of intrauterine pathologies because it is capable of simultaneously offering diagnosis and treatment for many of them. Genital tract infections, pregnancy, pelvic inflammatory disease, active herpetic infections, or human papilloma virus infections are contraindications for its performance. The indications are diverse, including suspected intracavitary mass, abnormal endometrial thickening, infertility, congenital anomalies, intrauterine adherence, in addition to post-treatment follow-up, and biopsy may be performed when necessary.¹

Initially, outpatient hysteroscopy was restricted to diagnostic procedures. With advances in technology, there has been an increase in their practice in the office, rather than in the operating room with anesthesia. Despite this, the outpatient method is associated with higher levels of pre- and intraoperative anxiety, which impairs patient satisfaction with the intervention and is associated with a greater perception of pain.² Furthermore, higher levels of pain during the intervention were associated with high rates of refusal to perform the procedure in the future, as well as higher rates of unsuccessful procedures.³ Thus, the impact of pain on the continuity of health care by patients is notorious, as well as its interference in the adequate control of their pathologies.

The article by Coimbra et al.,⁴ entitled Predictive Factors of Tolerance in Office Hysteroscopy – a 3-Year Analysis from a

Tertiary Center, addresses a topic of great relevance in the management of women's health. However, we would like to point out suggestions for improvement in the production of results that are of practical use.

We understand that in longitudinal and cross-sectional research designs, where the investigated outcome has a high prevalence or incidence (> 10%), the effect measure used to estimate the relationship/prediction between the independent variable and the outcome cannot be the odds ratio estimated by logistic regression.⁵ Applying this analytical approach produces overestimated point and interval estimates. The best strategy is the application of Poisson or Cox regression, depending on the type of outcome, generating measures such as relative risk or hazard ratio, respectively.⁵

In the study by Coimbra et al.,⁴ it is possible that the oversizing interfered with the significance of the age variable and the risk values of the other variables presented in Table 2, since the considered outcome is low tolerance (terrible and poor), which has an incidence of 14.9%. In addition, the reason for not including the variable intracavitary pathology, which revealed a difference in the outcome, and others such as parity number, was not identified in the model.

It is extremely important to create a raw model with the estimates of effects for the independent variables as a whole and the selection criteria for them to appear in the final

received
May 5, 2023
accepted
May 13, 2023

DOI <https://doi.org/10.1055/s-0043-1772469>.
ISSN 0100-7203.

© 2023. Federação Brasileira de Ginecologia e Obstetrícia. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)
Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

model. Commonly, factors may have a marginal relationship with the outcome in the crude analysis and in the adjusted model if it reveals a significant predictor, such as age, for example. Furthermore, overestimated estimates may limit findings in systematic reviews with meta-analysis.

Conflict of Interests

The authors have no conflict of interests to declare.

References

- 1 Yen CF, Chou HH, Wu HM, Lee CL, Chang TC. Effectiveness and appropriateness in the application of office hysteroscopy. *J Formos Med Assoc.* 2019;118(11):1480–1487
- 2 Giovanni B, Alonso PL, et al. Pain Management during Office Hysteroscopy: An Evidence-Based Approach. *Medicina MDPI.* 2022;x:1132–1132
- 3 McIlwaine K, Readman E, Cameron M, Maher P. Outpatient hysteroscopy: factors influencing post-procedure acceptability in patients attending a tertiary referral centre. *Aust N Z J Obstet Gynaecol.* 2009;49(06):650–652. Doi: 10.1111/j.1479-828X.2009.01096.x
- 4 Coimbra AC, Falcão V, Pinto P, Cavaco-Gomes J, Fernandes AS, Martinho M. Predictive Factors of Tolerance in Office Hysteroscopy - a 3-Year Analysis from a Tertiary Center. *Rev Bras Ginecol Obstet.* 2023;45(01):38–42. Doi: 10.1055/s-0043-1764361
- 5 Knol MJ, Le Cessie S, Algra A, Vandenbroucke JP, Groenwold RH. Overestimation of risk ratios by odds ratios in trials and cohort studies: alternatives to logistic regression. *CMAJ.* 2012;184(08): 895–899. Doi: 10.1503/cmaj.101715