



# Clinical assessment of upper limb impairments and functional capacity in Parkinson's disease: a systematic review

## *Avaliação clínica de comprometimentos de membros superiores e capacidade funcional na doença de Parkinson: uma revisão sistemática*

Tamine T. C. Capato<sup>1,2</sup>  Rúbia Rodrigues<sup>1</sup>  Rubens G. Cury<sup>1</sup>  Manoel Jacobsen Teixeira<sup>3</sup>   
Egberto R. Barbosa<sup>1</sup> 

<sup>1</sup> Universidade de São Paulo, Faculdade de Medicina, Departamento de Neurologia, Centro de Distúrbios do Movimento, São Paulo SP, Brazil.

<sup>2</sup> Radboud University Medical Centre, Donders Institute for Brain, Cognition and Behavior, Department of Neurology, Nijmegen, The Netherlands.

<sup>3</sup> Universidade de São Paulo, Departamento de Neurocirurgia, São Paulo SP, Brazil.

Address for correspondence Tamine T. C. Capato (email: taminec@usp.br)

Arq. Neuropsiquiatr. 2023;81(11):1008–1015.

### Abstract

**Background** Parkinson's disease (PD) may progressively reduce the upper limb's functionality. Currently, there is no standardized upper limb functional capacity assessment in PD in the rehabilitation field.

**Objective** To identify specific outcome measurements to assess upper limbs in PD and access functional capacity.

**Methods** We systematically reviewed and analyzed the literature in English published from August/2012 to August/2022 according to PRISMA. The following keywords were used in our search: "upper limbs" OR "upper extremity" and "Parkinson's disease." Two researchers searched independently, including studies accordingly to our inclusion and exclusion criteria. Registered at PROSPERO CRD42021254486.

**Results** We found 797 studies, and 50 were included in this review ( $n=2.239$  participants in H&Y stage 1–4). The most common upper limbs outcome measures found in the studies were: (i) UPDRS-III and MDS-UPDRS to assess the severity and progression of PD motor symptoms (tremor, bradykinesia, and rigidity) (ii) Nine Hole Peg Test and Purdue Pegboard Test to assess manual dexterity; (iii) Spiral test and Funnel test to provoke and assess freezing of upper limbs; (iv) Technology assessment such as wearables sensors, apps, and other device were also found.

**Conclusion** We found evidence to support upper limb impairments assessments in PD. However, there is still a large shortage of specific tests to assess the functional capacity of the upper limbs. The upper limbs' functional capacity is insufficiently

### Keywords

- ▶ Parkinson Disease
- ▶ Upper Extremity
- ▶ Physical Therapy Modalities
- ▶ Treatment Outcome
- ▶ Rehabilitation
- ▶ Freezing

received  
May 9, 2023  
received in its final form  
July 16, 2023  
accepted  
July 19, 2023

DOI <https://doi.org/10.1055/s-0043-1772769>.  
ISSN 0004-282X.

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (<https://creativecommons.org/licenses/by/4.0/>).

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

investigated during the clinical and rehabilitation examination due to a lack of specific outcome measures to assess functionality.

## Resumo

**Antecedentes** A doença de Parkinson (DP) reduz progressivamente a funcionalidade do membro superior. Não existe uma avaliação padronizada da capacidade funcional do membro superior na DP na área da reabilitação.

**Objetivo** Identificar medidas de resultados específicos para avaliar membros superiores na DP e avaliar capacidade funcional.

**Métodos** Revisamos e analisamos sistematicamente a literatura publicada de agosto/2012 a agosto/2022 de acordo com PRISMA. Usamos as seguintes palavras-chave “membros superiores” OU “extremidade superior” e “doença de Parkinson.” Dois pesquisadores fizeram a busca de forma independente, incluindo estudos de acordo com os critérios de inclusão e exclusão. Registro PROSPERO CRD42021254486.

**Resultados** Encontramos 797 estudos, 50 foram incluídos no estudo ( $n=2.239$  participantes no estágio 1–4 de H&Y). As medidas de resultados de membros superiores mais comuns encontradas foram: (i) UPDRS-III e MDS-UPDRS, para avaliar a gravidade e a progressão dos sintomas motores da DP (tremor, bradicinesia, e rigidez); (ii) Nine Hole Peg Test e Purdue Pegboard Test para avaliar a destreza manual; (iii) Teste da Espiral e Teste do Funil para provocar e avaliar o congelamento de membros superiores; (iv) Avaliação de tecnologia, como sensores vestíveis, aplicativos e outros dispositivos também foram encontrados.

**Conclusão** Encontramos evidências para dar suporte para as avaliações de deficiências de membros superiores na DP. No entanto, ainda há grande escassez de testes específicos para avaliar a capacidade funcional dos membros superiores. A capacidade funcional dos membros superior é insuficientemente investigada durante o exame clínico e de reabilitação devido à falta de medidas de resultados específicos para avaliar a funcionalidade.

## Palavras-chave

- ▶ Doença de Parkinson
- ▶ Extremidade Superior
- ▶ Modalidades de Fisioterapia
- ▶ Resultado do Tratamento
- ▶ Reabilitação
- ▶ Congelamento

## INTRODUCTION

Parkinson's disease (PD) is a neurodegenerative disease<sup>1,2</sup> diagnosed using clinical criteria, including bradykinesia, tremor, rigidity, and postural instability.<sup>3,4</sup> The clinical presentation can be multifaceted, including other motor and non-motor symptoms, differing among patients and subtypes.<sup>5</sup> The onset of symptoms is asymmetric, and since the early stages of PD, people experience a decrease in arm swing,<sup>6</sup> progressive speed reduction, and a decrease in the amplitude of the upper limb's repetitive movements.<sup>7</sup> Progressively, reduction of the upper limb's functional capacity generated by bradykinesia, tremor and rigidity may impact daily life activities, and freezing of upper limb (FOUL) episodes can be very disabling.<sup>8</sup>

Various measurement instruments used to assess gait, freezing of gait (FOG), and balance in Parkinson's disease are reported.<sup>9,10</sup> However, only a few instruments are available for clinical assessing upper limb impairments in PD.<sup>11</sup> Usually, the test and scales do not provide sufficient information about the quality of task performance quality or the test target according to the intervention proposed.<sup>12,13</sup>

Most of the tests and scales used in clinical practice and research measure the severity of PD motor symptoms including tremor, rigidity, bradykinesia,<sup>14</sup> manual dexterity,<sup>15,16</sup> and FOUL.<sup>8,11</sup> There is a gap of this instrument in clinical and

rehabilitation fields that can reliably measure the upper limb functional capacity of a person with PD. Currently, there are no recommendations to assess upper limb functionality<sup>12,13,17–19</sup> and standardized upper limb impairments assessment in PD.<sup>1,3,12,13,18</sup> Here, in this systematic review, we aimed:

- to identify the available outcome measures to assess upper limb impairments in people with PD; and
- to identify specific outcome measures to assess functional capacity in PD.

## METHODS

### Registration

This study was registered at PROSPERO CRD42021254486.

### Search strategy and selection criteria

We reviewed systematically the literature published from August 2012 to August 2022 according to PRISMA<sup>20</sup> (checklist - supplementary material). We analyzed published studies from a systematic review in the PubMed, using the following search: (“upper extremity”[MeSH Terms] OR (“upper”[All Fields] AND “extremity”[All Fields]) OR “upper extremity”[All Fields] OR (“upper”[All Fields] AND “limb”[All Fields]) OR “upper limb”[All Fields]) AND (“Parkinson disease”[MeSH Terms] OR (“Parkinson”[All Fields] AND “disease”[All Fields])

OR "Parkinson disease"[All Fields] OR "Parkinson s disease"[All Fields])) AND ("2012/08/22"[PDAT]: "2022/08/22"[PDAT]) AND.

## Eligibility criteria

### Inclusion criteria

- We included all peer-reviewed studies that reported an upper limb or upper extremity assessment and rehabilitation interventions in PD; only studies published in English were included in this review;
- observational studies, experimental studies, and quantitative study designs, including clinical trials, meta-analyses, systematic reviews, and case reports published from August 2012 to August 2022 were also included in this review.

### Exclusion criteria

- All studies that do not mention Parkinson's disease, and that do not present upper limbs assessments and rehabilitation interventions in methodology were excluded. Studies that used outcome measures and interventions that were not tested in PD patients were excluded.

### Study selection, study quality and risk of bias appraisal

Two researchers performed the search independently (RR and TC). A consensus meeting was held when needed

to include or excluded accordingly our criteria. The researchers also investigated the trial's effect size, and an outcome measure cut-off that could be used as an effect size of the interventions in future trials or at least indicate motor disease severity or level of upper limb disability. The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)<sup>20</sup> recommends the use of checklists to appraise study quality in systematic reviews. So, to evaluate the methodological quality of the included studies to determine whether the study was eligible for this review, and to reduce selection bias in the review, we used the Physiotherapy Evidence Database PEDro.<sup>21</sup> This database provides a good of information to evaluate the methodologic quality of the studies and risk of bias.

## RESULTS

Initially, we found 797 studies, 785 in PubMed, and 12 in other sources (►Figure 1). According to the inclusion and exclusion criteria, 50 studies were included in this systematic review (2.239 participants in H&Y stage 1–4). ►Supplementary Material Table 1 (<https://www.arquivos-deneuropsiquiatria.org/wp-content/uploads/2023/10/ANP-230102-Supplementary-Material.pdf>) shows in detail the characteristics of studies assessing upper limb impairments




Figure 1 Flow diagram.

in PD. In summary, we found many tests and scales which are used to assess upper limbs in PD:

- PD motor symptoms (severity and progression) can be assessed by Unified Parkinson's Disease Rating Scale<sup>22</sup> (UPDRS III) ( $n = 28$ ),<sup>7,15,23-47</sup> and MDS-Unified Parkinson's Disease Rating Scale<sup>14</sup> (MDS-UPDRS) ( $n = 18$ )<sup>6,8,11,16,48-61</sup>; To assess tremors we found Fahn-Tolosa-Marin Tremor Rating Scale<sup>62</sup> ( $n = 5$ ),<sup>32,44-46,60</sup> Tremor Rating Assessment Scale ( $n = 1$ )<sup>45</sup> and Spiral test ( $n = 2$ ).<sup>45,46</sup> Other tests and scales were found to assess bradykinesia such as the Finger tapping test ( $n = 2$ )<sup>38,63</sup> and Patient Global Impression of Change ( $n = 1$ ),<sup>46</sup> Motor assessment scale ( $n = 1$ ),<sup>61</sup> Performance measure ( $n = 1$ )<sup>61</sup> and Altering tapping performance ( $n = 1$ ).<sup>36</sup>
- Manual dexterity can be assessed using the Nine Hole Peg Test ( $n = 11$ )<sup>15,16,23,31,47-49,51,61,64,65</sup> and Purdue Pegboard Test ( $n = 9$ )<sup>16,24,33,44,50,61,63,64,66</sup>; Goal attainment scaling ( $n = 2$ )<sup>61,64</sup>; DextQ-24 ( $n = 1$ )<sup>48</sup>; Coin Rotation task ( $n = 03$ )<sup>16,48,49</sup>; Spiral (test  $n = 3$ )<sup>8,38,45</sup>; Manual Ability Measure-36 questionnaire ( $n = 1$ )<sup>49</sup>; Functional Reach test ( $n = 1$ )<sup>27</sup>; Occupational Therapy Neurologic Assessment battery dexterity task ( $n = 1$ )<sup>50</sup>; Bimanual dexterity hardware and experimental setup ( $n = 1$ )<sup>52</sup>; Box & Blocks test ( $n = 3$ )<sup>16,30,47</sup>; Hand temporal and spatial parameters ( $n = 1$ )<sup>30</sup>; Edinburgh handedness Inventory ( $n = 2$ )<sup>11,53</sup>; Patient-Specific Index-Parkinson's Disease and Self-assessment Parkinson's Disease Disability Scale ( $n = 1$ )<sup>15</sup>; Functional motor task ( $n = 1$ )<sup>51</sup>; Disabilities of the Arm, Shoulder and Hand (DASH) ( $n = 2$ )<sup>16,29</sup>;
- FOUL have been assessed by using the Spiral test ( $n = 3$ )<sup>8,38,44</sup> Funnel test ( $n = 2$ ).<sup>8,53</sup> We also found PD studies using technology to assess FOUL during alternating bimanual movements ( $n = 2$ )<sup>54,57</sup>; Finger tapping test ( $n = 2$ )<sup>44,55</sup>; handwriting and drawing patterns ( $n = 1$ )<sup>45</sup>;
- Technology has been used to assess the upper limb by using wearables sensors and apps or digital platforms.<sup>44,67</sup> Clinical-based, kinematic-based or kinematic have been assessed by using EMG ( $n = 3$ ).<sup>6,68,69</sup> Sensor units attached to the arms, hand and fingers can assess strength, movement power ( $n = 3$ ),<sup>6,37,56</sup> and arm swing ( $n = 1$ ).<sup>40</sup> Other outcomes were related to access Hand grip strength and finger measured by Dynamometer ( $n = 05$ ).<sup>25,47,48,50,64</sup> Power have been also assessed by using one repetition maximum (1RM) ( $n = 1$ )<sup>26</sup>; and movement resistance in the wrist and finger muscles ( $n = 1$ ).<sup>39</sup>

Only few studies included quality of life scales to verify the impact of upper limb impairments in daily life activities<sup>35,46,48,61,70</sup>; We did not find specific outcome measures to assess the functional capacity of upper limbs in PD. We did not find in the studies a specific intervention effect size, or an outcome measure cut-off that can be used to indicate the level of upper limb disability.

## DISCUSSION

In this study, we aimed to identify the available outcome measures to assess upper limb impairments in people with

PD; and specific outcome measures to assess functional capacity. Although we have found some evidence and useful outcome measures to assess of upper limb impairment assessments in PD, there is still a large shortage of specific tests to assess the functional capacity of the upper limbs in rehabilitation filed.

### Motor severity and disease progression

To assess the level of PD progression and motor severity of the disease, the UPDRS III and the MDS-UPDRS scales are widely used in many studies.<sup>6,7,23-27,33,48-50,53,63</sup> The MDS-UPDRS is the "gold standard" scale to access upper limbs tremors, rigidity and bradykinesia in clinical practice. By using specifically MDS-UPDRS domains (subscales), makes it possible to objectively evaluate upper limb resting tremor, and action tremor. It has also been reported that tremors can be assessed by using the Fahn-Tolosa-Marin Tremor Rating Scale<sup>30,42-44,58</sup> and Spiral test ( $n = 2$ ).<sup>43,44</sup> Our findings did not find any advantages in using the Fahn-Tolosa-Marin scale instead of MDS-UPDRS to access tremors. Future trials should investigate if one scale is superior to another to access tremors in the rehabilitation field. Other tests and scales can access bradykinesia, like the Finger tapping test.<sup>36,61</sup> Clinic-based experience and kinematic-based (EMG) assessments are also used to treat upper limb tremors in PD during BONT-A injection.<sup>32,68</sup>

### Specific impairments which directly impact functional movements

The current guidelines for PD provide no strong recommendations to assess upper limb functionality,<sup>12,13,17-19</sup> and there is no consensus about outcome measures to assess specific impairments that directly impact functional movements. However, studies have shown that it is possible to clinical assess other upper limb impairments, such as manual dexterity.<sup>16</sup> It is well known that upper limb impairments can be very disabling during daily life activities.<sup>16,55</sup> On the other hand, the current measures can be potentially difficult to assess specific motor symptoms, such as rigidity can impact functional movements.<sup>38,54</sup> The scarcity of specific tests to evaluate upper limbs functional capacity leads to insufficiently investigation during a clinical examination and in research protocols.

The most common tests found in our search to assess manual dexterity were Nine Hole Peg Test<sup>15,16,23,31,42,48,51,61,64,65</sup> and Purdue Pegboard Test.<sup>16,24,33,35,50,51</sup> Both tests are effective and are the most assertive in evaluating the manual dexterity in PD. The Coin rotation test has been used in few studies on dexterity in PD.<sup>29,48</sup> The DextQ-24 is an interesting questionnaire to access dexterity in daily live activities and everyday tasks such as washing/grooming, dressing and others.<sup>48</sup> Both, Coin Rotation task and DextQ-24 are easy and low cost to apply.

Our findings showed some outcome measures used for PD assessment in the studies searched were originally designed to access other diseases, such as "Test devaluation des Membres Supérieurs des Personnes Agées" (TEMPA).<sup>71</sup> Other examples are the Fugl-Meyer scale,<sup>31</sup> (it is a stroke-specific, performance-based impairment

index) and the Jebsen Hand Function Test, originally developed to assess gross and hand function in patients with cervical spinal cord injury.<sup>41</sup>

### Assess and provoke freezing of upper limbs (FOUL)

FOUL episodes can be very disabling during daily life activities. For this reason, FOUL should be objectively verified its presence, e.g., by evaluating the spiral-drawing task or the funnel task.<sup>8,9</sup> Interestingly, studies have shown that it is possible to provoke and assess FOUL in a clinical setting.<sup>11,55,72</sup> It is important to highlight, the tests most used to assess FOUL: flexion and extension of the index finger, finger tapping (index finger on thumb collected by MDS-UPDRS)<sup>23,55</sup> and Funnel Task.<sup>53</sup> During these tasks, it is possible to identify that movements of very small amplitude, high frequency, and execution in dual tasks lead to more FOUL episodes.<sup>11,53,57</sup> Only one of these studies found a correlation between their intervention and the effects of transcranial direct current stimulation on FOUL during the funnel test.<sup>53</sup> Previous studies affirm the importance of a therapy based on specific FOUL goals to achieve significant improvements.<sup>8,48,64</sup> Moreover, lower limb motor control can be associated to upper limb control, being more easily incorporated into regular daily tasks.<sup>73</sup> The spiral and funnel test are an assessment tool for FOUL during a task, which can be an important marker of the development of the pathology during the test time, as well as providing feedback for both clinicians and patient.<sup>38,74</sup> These studies align with the findings of,<sup>49</sup> emphasizing the need to consider the specificity of the proposed task, optimizing gains by including dual-task tasks exercises.<sup>6,34,56,73</sup> Therefore, FOUL deserves tailored treatment, and patients must be educated about compensation strategies by a physiotherapist with expertise in PD management.<sup>8</sup>

It is important to emphasize the importance of dual-task exercises<sup>23</sup> and the use of rhythmic cues to ameliorate FOUL.<sup>8,9</sup> Prior studies warned about the possibility of patients affected by FOUL and FOG, becoming dependent on rhythmic cues and highlighting how technology can be allied in assessing the delivering cues, optimizing their effectiveness.<sup>47</sup> It is still unclear whether the FOUL and FOG share the same physiological mechanism and the scarcity of specific tests to evaluate FOUL may limit the therapeutic approach.<sup>61</sup>

In this context, technology, through applications, can be an important tool for evaluating the commitment of the upper limbs, as well as an ally in the therapeutic approach and, a promising strategy in the area.<sup>6,32</sup>

### Technology in upper limb assessment

Our finds showed the use of technology to measure arm swing, and arm swing magnitude. Perhaps it can be useful for differentiating PD patients in early stage from healthy individuals.<sup>58</sup> In addition, some devices are portable and have been used to assess the impairments of the upper limbs in the clinical setting, and perhaps at home. Bradykinesia have been assessed on telemedicine with touch-pad to evaluation of alternating tapping performance using a touch-pad handheld, however the implementation of this method requires

more research.<sup>36</sup> An appropriate assessment will prevent false-negative results and allow the phenomenon to be identified and treated. So far, there is no strong evidence that these methods have any advantage over assessment traditional methods. Therefore, despite they are promising, future studies should evaluate the ability of these technologies to complement traditional upper limb clinical exams to optimize pharmacological, surgical, and non-pharmacological interventions.

### Limitations

Limitations of this review included possible bias due to the lack of measures currently in the development or studies that include motor fluctuations such as dyskinesias and individual participation in daily life activities. In our inclusion, there was heterogeneity across many studies regarding the outcome measures used to assess the same variable, which made it challenging to compare all the studies' results. In addition, the small samples and variability of methods were often difficult to assess. Therefore, it was not possible to perform a meta-analysis, a specific intervention effect size, and indicate an outcome measure cut-off that can be used to indicate the level of upper limb disability. Finally, we cannot make any statement about the technology to assess functional capacity and its use in the late stages of PD. Using technology during clinical measurements to monitor upper limb impairments in a clinical setting and home-based can be a promising strategy. However, large clinical trials should confirm these findings.

### Clinical implications

In our opinion, the upper limbs' functional capacity is under investigation during the clinical and rehabilitation examination due to a lack of specific outcome measures in the movement disorders field. Therefore, a proper outcome measure may be important not only as a marker of the progression of the pathology but also as a basis for therapeutic interventions that improve the quality of life of individuals.<sup>71</sup> For various reasons, we strongly believe that standard evaluation of upper limb functional capacity can be an essential element in PD clinical management. First, we can use this kind of assessment to develop and follow a pharmacological, non-pharmacological, and surgical treatment program in all disease stages of PD, to be consistently referred and to select the interventions according to the upper limb deficits. Second, to quantify motor deficits before and to demonstrate the results after clinical and surgical interventions including neuromodulation and rehabilitation program to optimize these treatments since the early stages of PD up to advanced stages. Finally, a functional capacity assessment could detect, monitor and support the decision of whether a person with PD can continue independently, work, and determine when tasks work-related should be adapted or discontinued. More studies are needed to verify this concept.

In conclusion, we found evidence to support upper limb impairment assessments in PD. However, there is still a large shortage of specific tests to assess the functional capacity of the upper limbs in movement disorders rehabilitation field.

Further studies should investigate technological advances to refine and support outcomes of assessing upper limb impairments.

#### Authors' Contributions

TTCC: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, resources, supervision, writing – review & editing; RR: data curation, formal analysis, investigation, methodology, writing – original draft, writing – review & editing; RG, MJT, ERB: writing – review & editing; TTCC, RR: These authors contributed equally. All authors contributed to the article and approved the submitted version.

#### Conflict of Interest

There is no conflict of interest to declare.

#### Acknowledgments

We also thank the support of University of São Paulo and PHYSICAL Parkinson's Disease and Movement Disorders Rehabilitation Center, São Paulo, Brazil.

#### References

- 1 Fox SH, Katzenschlager R, Lim SY, et al; Movement Disorder Society Evidence-Based Medicine Committee. International Parkinson and movement disorder society evidence-based medicine review: Update on treatments for the motor symptoms of Parkinson's disease. *Mov Disord* 2018;33(08):1248–1266
- 2 Espay AJ, Aybek S, Carson A, et al. Current Concepts in Diagnosis and Treatment of Functional Neurological Disorders. *JAMA Neurol* 2018;75(09):1132–1141
- 3 Armstrong MJ, Okun MS. Diagnosis and Treatment of Parkinson Disease: A Review. *JAMA* 2020;323(06):548–560
- 4 Postuma RB, Berg D, Stern M, et al. MDS clinical diagnostic criteria for Parkinson's disease. *Mov Disord* 2015;30(12):1591–1601
- 5 Bloem BR, Okun MS, Klein C. Parkinson's disease. *Lancet* 2021;397(10291):2284–2303
- 6 van den Noort JC, Verhagen R, van Dijk KJ, et al. Quantification of Hand Motor Symptoms in Parkinson's Disease: A Proof-of-Principle Study Using Inertial and Force Sensors. *Ann Biomed Eng* 2017;45(10):2423–2436
- 7 Jitkriksadikul O, Thanawattano C, Anan C, Bhidayasiri R. Tremor's glove—an innovative electrical muscle stimulation therapy for intractable tremor in Parkinson's disease: A randomized sham-controlled trial. *J Neurol Sci* 2017;381:331–340
- 8 Capato TT. C NJ, Barbosa E.R, Bloem R.B. Internal and external compensation strategies to alleviate upper limb freezing in Parkinson's disease. *Parkinsonism Relat Disord* 2019
- 9 Capato TTdC. Clinical Assessment and Management of Balance Impairments in Parkinson's disease. Donders Institute. Nijmegen, the Netherlands: Radboud University; 2022:168
- 10 Bloem BR, Marinus J, Almeida Q, et al; Movement Disorders Society Rating Scales Committee. Measurement instruments to assess posture, gait, and balance in Parkinson's disease: Critique and recommendations. *Mov Disord* 2016;31(09):1342–1355
- 11 Heremans E, Nackaerts E, Vervoort G, et al. Amplitude Manipulation Evokes Upper Limb Freezing during Handwriting in Patients with Parkinson's Disease with Freezing of Gait. *PLoS One* 2015;10(11):e0142874
- 12 Keus S, Munneke M, Graziano M, et al. European physiotherapy guideline for Parkinson's disease. 2014
- 13 Osborne JA, Botkin R, Colon-Semenza C, et al. Physical Therapist Management of Parkinson Disease: A Clinical Practice Guideline From the American Physical Therapy Association. *Phys Ther* 2022;102(04):pzab302
- 14 Goetz CG, Tilley BC, Shaftman SR, et al; Movement Disorder Society UPDRS Revision Task Force. Movement Disorder Society-sponsored revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): scale presentation and clinimetric testing results. *Mov Disord* 2008;23(15):2129–2170
- 15 Proud E, Morris ME, Bilney B, et al. Hand dexterity assessment in Parkinson's disease: construct validity of the 9-Hole peg test for the more affected hand. *Disabil Rehabil* 2021;43(26):3834–3838
- 16 Proud EL, Miller KJ, Bilney B, Balachandran S, McGinley JL, Morris ME. Evaluation of measures of upper limb functioning and disability in people with Parkinson disease: a systematic review. *Arch Phys Med Rehabil* 2015;96(03):540–551.e1
- 17 Capato TTdC, Domingos JMM, de Almeida LRS. Versão em Português da Diretriz Europeia de Fisioterapia para a Doença de Parkinson. Omini Farma. 2015:1
- 18 Saba RA, Maia DP, Cardoso FEC, et al. Guidelines for Parkinson's disease treatment: consensus from the Movement Disorders Scientific Department of the Brazilian Academy of Neurology – motor symptoms. *Arq Neuropsiquiatr* 2022;80(03):316–329
- 19 Sturkenboom IH, Graff MJ, Borm GF, et al. Effectiveness of occupational therapy in Parkinson's disease: study protocol for a randomized controlled trial. *Trials* 2013;14:34
- 20 Page MJ, Moher D, Bossuyt PM, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ* 2021;372(160):n160
- 21 Sherrington C, Herbert RD, Maher CG, Moseley AM. PEDro. A database of randomized trials and systematic reviews in physiotherapy. *Man Ther* 2000;5(04):223–226
- 22 Fahn SER. UPDRS Development Committee Unified Parkinson's Disease Rating Scale. In: Fahn S, Marsden CD, Calne DB, Goldstein M, editors. Recent developments in Parkinson's disease. Florham Park, NJ: Macmillan; 1987:153–163
- 23 Braun Janzen T, Haase M, Thaut MH. Rhythmic priming across effector systems: A randomized controlled trial with Parkinson's disease patients. *Hum Mov Sci* 2019;64:355–365
- 24 Momin S, Mahlknecht P, Georgiev D, et al. Impact of Subthalamic Deep Brain Stimulation Frequency on Upper Limb Motor Function in Parkinson's Disease. *J Parkinsons Dis* 2018;8(02):267–271
- 25 David FJ, Robichaud JA, Vaillancourt DE, et al. Progressive resistance exercise restores some properties of the triphasic EMG pattern and improves bradykinesia: the PRET-PD randomized clinical trial. *J Neurophysiol* 2016;116(05):2298–2311
- 26 Ni M, Mooney K, Signorile JF. Controlled pilot study of the effects of power yoga in Parkinson's disease. *Complement Ther Med* 2016;25:126–131
- 27 Ni M, Signorile JF, Mooney K, et al. Comparative Effect of Power Training and High-Speed Yoga on Motor Function in Older Patients With Parkinson Disease. *Arch Phys Med Rehabil* 2016;97(03):345–354.e15
- 28 Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ* 2016;354:i4086
- 29 Pazzaglia C, Imbimbo I, Tranchita E, et al. Comparison of virtual reality rehabilitation and conventional rehabilitation in Parkinson's disease: a randomised controlled trial. *Physiotherapy* 2020;106:36–42
- 30 Cikajlo I, Peterlin Potisk K. Advantages of using 3D virtual reality based training in persons with Parkinson's disease: a parallel study. *J Neuroeng Rehabil* 2019;16(01):119
- 31 Picelli A, Tamburin S, Passuello M, Waldner A, Smania N. Robot-assisted arm training in patients with Parkinson's disease: a pilot study. *J Neuroeng Rehabil* 2014;11:28

- 32 Samotus O, Lee J, Jog M. Long-term tremor therapy for Parkinson and essential tremor with sensor-guided botulinum toxin type A injections. *PLoS One* 2017;12(06):e0178670
- 33 Eggers C, Günther M, Rothwell J, Timmermann L, Ruge D. Theta burst stimulation over the supplementary motor area in Parkinson's disease. *J Neurol* 2015;262(02):357–364
- 34 Messa LV, Ginanneschi F, Momi D, et al. Functional and Brain Activation Changes Following Specialized Upper-Limb Exercise in Parkinson's Disease. *Front Hum Neurosci* 2019;13:350
- 35 Nowinski CJ, Siderowf A, Simuni T, Wortman C, Moy C, Cella D. Neuro-QoL health-related quality of life measurement system: Validation in Parkinson's disease. *Mov Disord* 2016;31(05):725–733
- 36 Memedi M, Khan T, Grenholm P, Nyholm D, Westin J. Automatic and objective assessment of alternating tapping performance in Parkinson's disease. *Sensors (Basel)* 2013;13(12):16965–16984
- 37 Jo HJ, Ambike S, Lewis MM, Huang X, Latash ML. Finger force changes in the absence of visual feedback in patients with Parkinson's disease. *Clin Neurophysiol* 2016;127(01):684–692
- 38 Saunders-Pullman R, Derby C, Stanley K, et al. Validity of spiral analysis in early Parkinson's disease. *Mov Disord* 2008;23(04):531–537
- 39 Zetterberg H, Frykberg GE, Gäverth J, Lindberg PG. Neural and nonneural contributions to wrist rigidity in Parkinson's disease: an explorative study using the NeuroFlexor. *BioMed Res Int* 2015;2015:276182
- 40 Sterling NW, Cusumano JP, Shaham N, et al. Dopaminergic modulation of arm swing during gait among Parkinson's disease patients. *J Parkinsons Dis* 2015;5(01):141–150
- 41 Ryan D, Fullen B, Rio E, Segurado R, Stokes D, O'Sullivan C. Effect of Action Observation Therapy in the Rehabilitation of Neurologic and Musculoskeletal Conditions: A Systematic Review. *Arch Rehabil Res Clin Transl* 2021;3(01):100106
- 42 França C, de Andrade DC, Teixeira MJ, et al. Effects of cerebellar neuromodulation in movement disorders: A systematic review. *Brain Stimul* 2018;11(02):249–260
- 43 GBD 2016 Parkinson's Disease Collaborators. Global, regional, and national burden of Parkinson's disease, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol* 2018;17(11):939–953
- 44 Simpson MW, Mak M. The effect of transcranial direct current stimulation on upper limb motor performance in Parkinson's disease: a systematic review. *J Neurol* 2020;267(12):3479–3488
- 45 Lora-Millan JS, Delgado-Oleas G, Benito-León J, Rocon E. A Review on Wearable Technologies for Tremor Suppression. *Front Neurol* 2021;12:700600
- 46 Mittal SO, Jog M, Lee J, Jabbari B. Novel Botulinum Toxin Injection Protocols for Parkinson Tremor and Essential Tremor - the Yale Technique and Sensor-Based Kinematics Procedure for Safe and Effective Treatment. *Tremor Other Hyperkinet Mov (N Y)* 2020;10:61
- 47 Moudjian L, Sarkamo T, Leone C, Leman M, Feys P. Effectiveness of music-based interventions on motricity or cognitive functioning in neurological populations: a systematic review. *Eur J Phys Rehabil Med* 2017;53(03):466–482
- 48 Vanbellinghen T, Nyffeler T, Nigg J, et al. Home based training for dexterity in Parkinson's disease: A randomized controlled trial. *Parkinsonism Relat Disord* 2017;41:92–98
- 49 Allen NE, Song J, Paul SS, et al. An interactive videogame for arm and hand exercise in people with Parkinson's disease: A randomized controlled trial. *Parkinsonism Relat Disord* 2017;41:66–72
- 50 Mateos-Toset S, Cabrera-Martos I, Torres-Sánchez I, Ortiz-Rubio A, González-Jiménez E, Valenza MC. Effects of a Single Hand-Exercise Session on Manual Dexterity and Strength in Persons with Parkinson Disease: A Randomized Controlled Trial. *PM R* 2016;8(02):115–122
- 51 Paul SS, Dibble LE, Olivier GN, Walter C, Duff K, Schaefer SY. Dopamine replacement improves motor learning of an upper extremity task in people with Parkinson disease. *Behav Brain Res* 2020;377:112213
- 52 Jansen AE, Koop MM, Rosenfeldt AB, Alberts JL. Aerobic exercise does improve bimanual coordination in Parkinson's disease: Response to Samuel and colleagues. *Parkinsonism Relat Disord* 2021;93:103–104
- 53 Broeder S, Heremans E, Pinto Pereira M, et al. Does transcranial direct current stimulation during writing alleviate upper limb freezing in people with Parkinson's disease? A pilot study. *Hum Mov Sci* 2019;65:65
- 54 Nieuwboer A, Vercruyse S, Feys P, Levin O, Spildooren J, Swinnen S. Upper limb movement interruptions are correlated to freezing of gait in Parkinson's disease. *Eur J Neurosci* 2009;29(07):1422–1430
- 55 Delval A, Defebvre L, Tard C. Freezing during tapping tasks in patients with advanced Parkinson's disease and freezing of gait. *PLoS One* 2017;12(09):e0181973
- 56 Pradhan S, Scherer R, Matsuoka Y, Kelly VE. Grip force modulation characteristics as a marker for clinical disease progression in individuals with Parkinson disease: case-control study. *Phys Ther* 2015;95(03):369–379
- 57 Williams AJ, Peterson DS, Ionno M, Pickett KA, Earhart GM. Upper extremity freezing and dyscoordination in Parkinson's disease: effects of amplitude and cadence manipulations. *Parkinsons Dis* 2013;2013:595378
- 58 Ospina BM, Chaparro JAV, Paredes JDA, Pino YJC, Navarro A, Orozco JL. Objective Arm Swing Analysis in Early-Stage Parkinson's Disease Using an RGB-D Camera (Kinect®). *J Parkinsons Dis* 2018;8(04):563–570
- 59 Cavallo F, Moschetti A, Esposito D, Maremmanni C, Rovini E. Upper limb motor pre-clinical assessment in Parkinson's disease using machine learning. *Parkinsonism Relat Disord* 2019;63:111–116
- 60 Chan PY, Ripin ZM, Halim SA, et al. An In-Laboratory Validity and Reliability Tested System for Quantifying Hand-Arm Tremor in Motions. *IEEE transactions on neural systems and rehabilitation engineering: a publication of the IEEE Engineering in Medicine and Biology Society*. 2018;26(02):460–467
- 61 Proud EL, Miller KJ, Martin CL, Morris ME. Upper-limb assessment in people with Parkinson disease: is it a priority for therapists, and which assessment tools are used? *Physiother Can* 2013;65(04):309–316
- 62 Fahn S, Tolosa E, Marín C. Clinical rating scale for tremor. *Parkinson's disease and movement disorders*. 1993;2:271–280
- 63 Kueper JK, Speechley M, Lingum NR, Montero-Odasso M. Motor function and incident dementia: a systematic review and meta-analysis. *Age Ageing* 2017;46(05):729–738
- 64 Cabrera-Martos I, Ortiz-Rubio A, Torres-Sánchez I, Rodríguez-Torres J, López-López L, Valenza MC. A randomized controlled study of whether setting specific goals improves the effectiveness of therapy in people with Parkinson's disease. *Clin Rehabil* 2019;33(03):465–472
- 65 Park JK, Kim SJ. Dual-Task-Based Drum Playing with Rhythmic Cueing on Motor and Attention Control in Patients with Parkinson's Disease: A Preliminary Randomized Study. *Int J Environ Res Public Health* 2021;18(19):10095
- 66 Vingerhoets FJ, Schulzer M, Calne DB, Snow BJ. Which clinical sign of Parkinson's disease best reflects the nigrostriatal lesion? *Ann Neurol* 1997;41(01):58–64
- 67 Knippenberg E, Verbrugghe J, Lamers I, Palmaers S, Timmermans A, Spooren A. Markerless motion capture systems as training device in neurological rehabilitation: a systematic review of their

- use, application, target population and efficacy. *J Neuroeng Rehabil* 2017;14(01):61
- 68 Samotus O, Mahdi Y, Jog M. Real-World Longitudinal Experience of Botulinum Toxin Therapy for Parkinson and Essential Tremor. *Toxins (Basel)* 2022;14(08):557
- 69 Abbruzzese G, Trompetto C, Mori L, Pelosin E. Proprioceptive rehabilitation of upper limb dysfunction in movement disorders: a clinical perspective. *Front Hum Neurosci* 2014;8: 961
- 70 Samotus O, Lee J, Jog M. Transitioning from Unilateral to Bilateral Upper Limb Tremor Therapy for Parkinson's Disease and Essential Tremor Using Botulinum Toxin: Case Series. *Toxins (Basel)* 2018; 10(10):394
- 71 de Freitas PR, Lemos AE, Santos MP, Michaelsen SM, Corrêa CL, Swarowsky A. "Test D'évaluation Des Membres Supérieurs Des Personnes Âgées" (TEMPA) to assess upper limb activity in Parkinson's disease. *J Hand Ther* 2017;30(03):320-327
- 72 Vercruyse S, Spildooren J, Heremans E, et al. Freezing in Parkinson's disease: a spatiotemporal motor disorder beyond gait. *Mov Disord* 2012;27(02):254-263
- 73 Heideman SG, te Woerd ES, Praamstra P. Rhythmic entrainment of slow brain activity preceding leg movements. *Clin Neurophysiol* 2015;126(02):348-355
- 74 Snijders AH, Haaxma CA, Hagen YJ, Munneke M, Bloem BR. Freezer or non-freezer: clinical assessment of freezing of gait. *Parkinsonism Relat Disord* 2012;18(02):149-154