

# Endoscopic Band Ligation for Rectal Dieulafoy's Lesion: Serial Endoscopic Images

Y. Mizukami<sup>1</sup>, K. Akahoshi<sup>1</sup>, N. Kondoh<sup>1</sup>,  
N. Harada<sup>2</sup>, H. Nawata<sup>1</sup>

<sup>1</sup> Department of Gastroenterology,  
Aso Iizuka Hospital, Iizuka, Japan

<sup>2</sup> Third Department of Internal Medicine,  
Faculty of Medicine, Kyushu University,  
Fukuoka, Japan

## Corresponding Author

K. Akahoshi, M.D., Ph.D.

Department of Gastroenterology  
Aso Iizuka Hospital


3-83 Yoshio

Iizuka 820-8505


Japan

Fax: +81-948-298747

E-mail: kakahoshih1@aih-net.com



**Figure 1** An 85-year-old Japanese woman with rectal Dieulafoy's lesion presented with fresh blood hematochezia. Colonoscopy revealed a 5 mm shallow mucosal defect with an exposed vessel in the anterior wall of the lower rectum.



**Figure 2** A gastroduodenoscope with an attached single-band ligator was employed, and 4 ml of hypertonic saline epinephrine solution was injected beneath the lesion. After endoscopic band placement, the hemorrhage stopped completely. Note that the Dieulafoy's lesion had been wholly encased by the band.



**Figure 3** Endoscopy at 1 week after the procedure revealed a 10 mm wide shallow ulcer at the site of the lesion. The O-ring remained on the ulcerated lesion.



**Figure 4** Follow-up colonoscopy (at 2 weeks) showed an ulcer scarring. No further bleeding has occurred during 6 months of follow-up.