


**Figure 1** A 42-year-old woman presented with jaundice, right upper quadrant pain and pruritus. Laboratory test findings were: hemoglobin (Hb) 11.1; bilirubin 4.6; aspartate aminotransferase (AST) 34; alanine aminotransferase (ALT) 38; prothrombin time (PT) 16; and alkaline phosphatase 1083. Abdominal sonography revealed dilatation of the intra- and extrahepatic biliary ducts. Endoscopic retrograde cholangiopancreatography (ERCP) showed dilated bile ducts with a large lobulated filling defect in the proximal portion of the common bile duct toward the intrahepatic ducts. CBD, common bile duct; CHD, common hepatic duct.



**Figure 3** Histological investigation demonstrated a choledochal cystadenoma.

**A. Hedayat<sup>1</sup>, R. Sotoudehmanesh<sup>2</sup>,  
M. Khatibian<sup>2</sup>, H. Mahmoudzadeh<sup>1</sup>,  
M. Sotoudeh<sup>3</sup>**


<sup>1</sup> Department of Surgery, Tehran University of Medical Sciences, Tehran, Iran

<sup>2</sup> Digestive Disease Research Center, Tehran University of Medical Sciences, Tehran, Iran

<sup>3</sup> Department of Pathology, Shariati Hospital, Tehran University of Medical Sciences, Tehran, Iran

### Corresponding Author

R. Sotoudehmanesh, M.D.  
Digestive Disease Research Center,  
Shariati Hospital  
North Kagar Avenue, 14114 Tehran, Iran  
Fax: +98-21-8012992  
E-mail: setoodeh@ams.ac.ir



**Figure 2** During surgery a pedunculated mass lesion was seen extending from the left hepatic duct to the mid-portion of the CBD.