

A rare cause of facial purpura: endoscopy



Figure 1 Pin-head-sized red macules on patient's face and neck.

A 43-year-old woman who underwent stomach endoscopy due to gastritis was referred to our department because of an eruption on her face, which had started during the procedure. Due to the frightening appearance of the lesions, the patient and the endoscopist failed to complete the procedure. On dermatologic examination, hundreds of pin-head-sized red macules on the face, oral mucosa, and neck were seen (● **Figure 1**). Subconjunctival hemorrhage was also observed (● **Figure 2**). There was no pruritus or constitutional symptoms. The patient was otherwise healthy. Hematologic causes were excluded, with normal complete blood count, prothrombin time, and partial thromboplastin time. The rash started to fade spontaneously within 24 hours, and had disappeared completely after 4 days without treatment.

Several factors may contribute towards purpura on the face and neck. The most challenging step in identifying the cause of purpura is to decide whether the purpura presents a sign of an underlying life-threatening disorder or not. The differential diagnosis of purpura can be confusing, and consists of a long list of factors including rheumatologic, dermatologic, infectious, and traumatic entities [1].



Figure 2 Subconjunctival hemorrhage.

In some circumstances, facial purpura is caused by a valsalva maneuver that raises intrathoracic or abdominal pressure and results in rupture of capillaries within the dermis. Alcalay et al. [2] suggested the name “mask phenomenon” to describe this condition. Rare occurrence of purpura on the face is thought to be related to the rich anastomotic vascular supply to face, or the lack of end-artery vessels in this area [1]. Giving birth, emesis, and vigorous coughing are examples of this kind, and are well-known causes of facial purpura [1,2]. Prolonged hand-standing [3] and epileptic seizure [4,5] are also other benign conditions that lead to facial purpura, as reported in the literature. In our patient, the endoscopy probe triggered the retch reflex and caused vigorous coughing.

We want to remind physicians of this benign cause of facial purpura in order to avoid unnecessary investigation. Endoscopic procedures are not easy to tolerate, and internists should be aware of this complication of the procedure. This kind of eruption should be added to the list of procedure-related complications of endoscopy.

Endoscopy_UCTN_Code_CPL_1AH_2AB

E. Adışen, N. Eroglu, M. Öztaş, M. A. Gürer

Gazi University Faculty of Medicine, Department of Dermatology, Ankara, Turkey

References

- 1 Goldman AC, Govindaraj S, Franco RA Jr et al. Facial purpura. *Laryngoscope* 2001; 111: 207–212
- 2 Alcalay J, Ingber A, Sandbank M. Mask phenomenon: postemesis facial purpura. *Cutis* 1986; 38: 28
- 3 Bartunek C, Brodell RT, Brodell LP. Self-assessment quiz. Valsalva purpura. *J Ophthalmic Nurs Technol* 1998; 17: 23–24
- 4 Reis JJ, Kaplan PW. Postictal hemifacial purpura. *Seizure* 1998; 7: 337–339
- 5 Rubegni P, Fimiani M, De Aloe G et al. Thoracocervicofacial purpura as the single manifestation of epileptic seizure. *J Neurol Neurosurg Psychiatry* 1998; 65: 365

Bibliography

DOI 10.1055/s-2007-966317

Endoscopy 2007; 39: E216

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

E. Adışen, MD

Gazi University Faculty of Medicine,

Department of Dermatology

06500 Beşevler

Ankara

Turkey

Fax: +90-312-212-9018

eozsoy@gazi.edu.tr