

Metastatic endometrial cancer; a rare intestinal localization



Figure 1 Tumor in the small bowel.

Endometrial cancer can give rise to haemogenous metastases, most often in lungs, brain, and bones. We describe a patient with metastases of endometrial cancer in the small bowel.

An 85-year-old female patient was admitted with severe anemia and melena. She had endometrial cancer (Figo stage IIIB), 2 years previously, for which she underwent an operation and received adjuvant radiotherapy. She was admitted with severe anemia (Hb 3.8 mmol/L) and melena. Upper endoscopy did not reveal any abnormalities. Colonoscopy was normal.

Video capsule endoscopy was carried out. With this examination, two tumors were discovered with signs of recent hemorrhage (adherent clot) (● Figure 1). During laparotomy, a large tumor in the small bowel was encountered. A large part of the small bowel was resected. An end-to-end enteroenterostomy was carried out. Histological examination showed a carcinosarcoma with extensive angioinvasive growth. The immunohistochemical picture was identical to the endometrial tumor, which was removed in 2004. As well as features of a carcinoma, signs of a sarcoma were also seen, compatible with tumor differentiation (● Figure 2a and ● 2b).




Figure 2 a Adenocarcinoma in the resection specimen. b In the same tumor, signs of a sarcoma were found, indicating tumor differentiation.

On the basis of the initial stage of the endometrial carcinoma (Figo IIIB), this patient was at risk for developing metastases. In this group of patients, the majority develops metastases within 3 years, most often in the peritoneal cavity, lungs, liver, bones, or brain.

Intestinal metastases of endometrial cancer have been described in the literature, but are rarely seen. Biegel et al. describe bleeding colonic metastases in a patient with endometrial cancer [1]. Metastases in the small bowel have also been described [2], sometimes requiring segmental small-bowel resection because of bleeding complications [3].

The small-bowel metastases caused massive bleeding and were diagnosed by means of videocapsule endoscopy. With the introduction of this diagnostic tool, access to the small bowel and diagnostic yield have improved significantly.

Endoscopy_UCTN_Code_CCL_1AC_2AC

W. J. Thijs¹, A. Karrenbeld²,
L. van der Zouwen³, L. de Haan¹

¹ Department of Internal Medicine and Gastroenterology, Scheperziekenhuis, Emmen, The Netherlands

² Department of Pathology, University Medical Hospital, Groningen, The Netherlands

³ Department of Surgery, Scheperziekenhuis, Emmen, The Netherlands

References

- 1 Biegel E. Scintigraphic demonstration of bleeding colonic metastases in a patient with endometrial cancer. *Ugeskr Laeger* 1995; 157: 3337–3338
- 2 Bosscher J, Barnhill D, O'Connor D, Park R. Clinical stage IB endometrial adenocarcinoma with an isolated small bowel metastasis. *Gynaecol Oncol* 1994; 52: 99–101
- 3 Schneider JJ, Shroff S, Moser AJ. Palliative segmental duodenectomy for bleeding, erosive endometrial cancer. *Gynaecol Oncol* 2005; 97: 246–248

Bibliography

DOI 10.1055/s-2007-966372

Endoscopy 2007; 39: E131

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

W. J. Thijs, MD

Department of Internal Medicine and Gastroenterology
Scheperziekenhuis Emmen
Boermarkeweg 60
7824 AA, Emmen
The Netherlands
Fax: +31-591-691361
w.thijs@sze.nl