


Y. TANAKA, S. SAITO, S. MORI, N. ARATANI, H. SHINOKUBO, N. SHIBATA, Y. HIGUCHI, Z. S. YOON, K. S. KIM, S. B. NOH, J. K. PARK, D. KIM,\* A. OSUKA\* (YONSEI UNIVERSITY, SEOUL, KOREA; KYOTO UNIVERSITY, UNIVERSITY OF HYOGO AND THE RIKEN SPRING-8 CENTER, HYOGO, JAPAN)

Metalation of Expanded Porphyrins: A Chemical Trigger Used To Produce Molecular Twisting and Möbius Aromaticity

*Angew. Chem. Int. Ed.* **2008**, *47*, 681-684.

## Metalation-Triggered Möbius Aromaticity



**Significance:** A new way to achieve complexes that exhibit Möbius aromaticity is described. Conjugated oligopyrrolic macrocycles (also known as expanded porphyrins) were locked into a twisted Möbius strip conformation using group 10 metals. Experimental and numerical studies reveal compounds **3**, **5**, and **7–9** as conformationally locked Möbius aromatic molecules.

**Comment:** Structures and aromaticity were confirmed via XRD (**2**, **3**, **5**),  $^1\text{H-NMR}$  and UV/Vis. Significantly negative values of calculated NICS (Nucleus-Independent Chemical Shift) supported Möbius-type aromaticity for **3**, **5**, and **7–9**. A new method for the quantitative analysis of aromaticity based on the measurement of the TPA (two-photon absorption) cross-section ( $\sigma^{(2)}$ ) by the open-aperture Z-scan method showed consistent results.

**SYNFACTS Contributors:** Timothy M. Swager, Jose M. Lobe  
 Synfacts 2008, 4, 0363-0363 Published online: 19.03.2008  
 DOI: 10.1055/s-2008-1042849; Reg-No.: S01508SF