

Editorial

Gastric Lymphoma and Epstein-Barr Virus

Epstein-Barr virus (EBV) is the major biological cofactor contributing to a number of human cancers including B-cell neoplasms (Burkitt's lymphoma, Hodgkin's lymphoma (HL and immunoblastic lymphomas, certain forms of T-cell lymphoma, and some epithelial tumours (nasopharyngeal carcinomas and gastric carcinomas). Recent studies have revealed the association of EBV with about 10% of gastric carcinoma cases worldwide. EBV DNA in cancer biopsies shows monoclonality, indicating that carcinoma arises from a single EBV-infected cell.¹

Primary malignant gastric lymphomas account for less than 15% of gastric malignancies. Most gastric lymphomas are B-cell lineage NHL. They usually arise from mucosal associated lymphoid tissue (MALT). Other types of gastric lymphomas include diffuse large B-cell lymphomas, mantle cell lymphomas and T cell lymphomas.²

Seventy to 100% of MALT lymphoma cases are associated with *Helicobacter pylori*, a bacterium proven to be the cause of gastritis and peptic ulcer and the progression to gastric carcinoma. Lower *H. pylori* infection rates have been reported in gastric lymphoma in Asian populations (50%-60%).³

Gastric lymphoma association with EBV has not been extensively studied. The pilot study by Sood *et al* published in this issue of Journal reports EBV detection in neoplastic cells in 36% of 11 cases,⁴ higher than reports from Hong Kong, Korea and China (2% to 8% in series of 53, 33 and 49 cases respectively).^{5,5} However, the small sample size does not allow any meaningful conclusion.

Detection of EBV in tumour cells by immunohistochemistry or in situ hybridization is no proof of its implication in the pathogenesis. EBV might either play a role in the development of gastric lymphoma, particularly in *H. pylori* negative cases, or be present in neoplastic cells as a secondary phenomenon. Demonstration of EBV clonality in EBV-associated gastric lymphomas would be able to give a definitive answer.

REFERENCES:

1. Imai S, Koizumi S, Sugiura M, et al. Gastric carcinoma: monoclonal epithelial malignant cells expressing Epstein-Barr virus latent infection protein. *Proc Natl Acad Sci U S A* 1994;91(19):9131-9135.
2. WHO classification of tumours of the digestive system: Hamilton SR, Aaltonen LA, eds. *Tumours of the digestive system. World Health Organization classification of tumours.* Lyon: ARC Press, 2000.
3. Xu WS, Ho FC, Ho J, et al. Pathogenesis of gastric lymphoma: the enigma in Hong Kong. *Ann Oncol.* 1997;8 Suppl 2:41-44.
4. Sood A, Braganza A & T Rajalakshmi, Epstein-Barr Virus in gastric lymphoma-An Indian Perspective *Ind. J Med & Paed Oncol* 2008;29(3):5-9.
5. Liu Q, Ohshima K, Masuda Y, Kikuchi M. Detection of the Epstein-Barr virus in primary gastric lymphoma by in situ hybridization. *Pathol Int* 1995;45(2):131-136.
6. Yang WI, Cho MS, Tomita Y, et al. Epstein-Barr virus and gastrointestinal lymphomas in Korea. *Yonsei Med J* 1998;39(3):268-276.

Veronique Dinand,
Pediatric Haematology Oncology
Centre for Child Health
Sir Ganga Ram Hospital
New Delhi 110 060
E-mail: verodinand@fastem.com

