

Geriatric oncology: The need for a separate subspecialty

In India, we are slowly moving toward an increased ratio of geriatric population. Statistics predict that in India, by 2021, the population in the age group 0–14 would drop by 7.6%, whereas those above 60 would increase by 102% when compared to the population in 2001^[1] [Table 1].

In Kerala, there is a 12% fall in the under 14 age group with a corresponding increase of 73% in the above 60 age group^[2] [Table 2].

This would automatically translate into more geriatric problems in the health-care sector. Cancer is 11 times more likely to develop in people above 65 years compared to younger people.^[3] Some of the reasons why cancer is more common in the elderly include telomere shortening and defective deoxyribo nucleic acid (DNA) repair mechanisms, age-related alterations in the immune system, accumulation of random genetic mutations leading to oncogene activation, amplification or decreased tumor suppressor gene activity, life-time carcinogen exposure, and hormonal alterations or exposure.

For the proper management of these patients, a better understanding of the physiology of aging is required. In geriatric age group, there is increase in the body fat and decreased total body water and muscle mass secondary to hormone dysfunction and decreased protein synthesis, loss of sweat glands, decreased gastrointestinal (GI) motility, and absorption and decreased hepatic and renal function. There is decrease in weight of the brain and also there is a decline in the amount of neurotransmitters with age. There is also immunologic dysregulation and compromised marrow reserve.

The assessment of these patients requires a different approach and comprehensive geriatric assessment (CGA) is recommended for older patients with cancer.^[4,5] This adds information to the presently used measures of functional status in oncology, such as the Karnofsky or Eastern Cooperative Oncology Group Performance Status.^[6] The domains of a CGA include assessment of functional status, co-morbidities, cognitive function, socioeconomic status, medications, and nutrition. Each of these domains that are described below predicts the risk of morbidity and mortality in older adults.^[6]

Functional status is the ability of the patient to perform

routine daily activities. Functional disability is common in older patients with cancer.^[7] Activities of daily living (ADL) (skills that are necessary for basic living, includes feeding, grooming, transferring, and toileting) and Instrumental activities of daily living (IADL) (activities such as shopping, managing finances, housekeeping, preparing meals, and taking medications) scales can be used for a better understanding of the functional status in geriatric patients. According to Repetto *et al.*^[8], patients who were ADL dependant or IADL dependant had a nearly twofold higher probability of having an elevated CGA score than independent patients.

Comorbidity represents the extent of an individual's physical and psychological disease in addition to the disease for which he or she is seeking treatment.^[8] Co morbid medical conditions are high in geriatric population^[9] and they adversely affect the life expectancy and treatment tolerance. Repetto *et al.*^[10] noticed that, patients with two or more comorbidities too were found to have a good PS (<2). This should be taken into consideration while estimating the risks and benefits of treatment.^[11]

In general geriatric population, dementia is an independent prognostic factor for survival. The presence of dementia influences the likelihood of both cancer diagnosis and treatment.^[12] Patients with dementia were less likely to have a biopsy-proven diagnosis of cancer and were less likely to receive surgery or chemotherapy.^[13,14]

In general, in older population as well as in patients with cancer, weight loss or low body mass index was found to be associated with increase in mortality.^[15–17] Lower socioeconomic status has an adverse effect on survival due to the effects on nutrition, social support, and access to better surgical or chemotherapeutic options. Social isolation

Table 1: Population changes in India

Age	Population 2001	Population 2021	Change (%)
0–14	364,582,000	336,906,000	–7.6
60+	70,686,000	143,244,000	+102

Table 2: Population changes in Kerala

Age	Population 2001	Population 2021	% Change
0–14	8,303,000	7,319,000	–12
60+	3,362,000	5,833,000	+73

is also an independent predictor of mortality in the geriatric population.^[18] A careful assessment of the medications taken by the patient is necessary to remove any unnecessary medicines and also to look for potential drug interactions.^[19-21]

Even though the majority of cancer patients are in the older age group, participation of older patients in cancer treatment trials had been very low. In Canada, it was found that patients who were 65 years or older accounted only for 22% of patients in the National Cancer Institute of Canada Clinical Trials Group compared with 58% of the Canadian population with cancer in 1993–1996.^[22] Also in India, the number of older cancer patients enrolled in trials will be much less compared to the other age groups. The enrolment of older people in adjuvant trials was lower than their enrolment in advanced clinical trials. This may be because many older patients present in advanced stage^[23–25] and have poor physical condition. It may also be due to reluctance on the part of the physicians or patients to enroll in adjuvant trials.^[22] But now there are a few randomized controlled trials specifically for the geriatric population, like the Intergroupe Francophone de Cancérologie Thoracique 0501 for advanced nonsmall cell carcinoma of the Lung.^[26]

In India, according to registry data about 45–50% of all the cancers in males and 35–41% of all cancer in females occur above the age of 60.^[27] The age specific rate of all cancers affecting males and females in India is represented in Figures 1 and 2.

Based on the incidence rates published in the Indian Council of Medical Research population-based cancer registry report^[27] and Dindigul Ambilikai Cancer registry,^[28] there will be more than 200,000 males and nearly 150,000 females newly diagnosed as cancer every year in India by 2021. The prevalence of cancer patients in India above the age of 60 is estimated to reach more than 1 million by 2021.

Kerala though a small state, has three population-based cancer registries; two in the rural area and one in the urban area (Trivandrum rural and Karunagapally *vs* Trivandrum urban).

Based on the incidence rates of Karunagapally and Trivandrum^[29,30] it can be estimated that there will be around 20,171 people above 65 years newly diagnosed as cancer every year in Kerala by 2021. In addition, in Kerala the prevalence of cancer patients above the age of 60 is estimated to be more than 60,000 by 2021.

As these patients have special needs and a different approach to treatment, there is a strong case for the




Figure 1: Age specific rates of cancer in males




Figure 2: Age specific rates of cancer in females

emergence of geriatric oncology as a sub specialty in oncology.

*D. K. Vijaykumar, R. Anupama, Tejal Kishor Gorasia,
T. R. Haleema Beegum, P. Gangadharan¹*

*Department of Surgical Oncology, and ¹Cancer Registry,
Amrita Institute of Medical Sciences and Research Centre,
Amrita Viswavidyapeetham, Kochi, Kerala, India
E-mail: anupamar@aims.amrita.edu*

REFERENCES


1. Census of India 2001. Population projection for India and States 2001-2026. p. 139-40.
2. Census of India 2001. Population projection for India and States 2001-2026. p. 239-40.
3. Ries LAG, Eisner MP, Kosary CL, Hankey BF, Miller BA, Clegg LX, *et al.* editors. SEER cancer statistics review. In: National Cancer Institute. Bethesda, MD; 1975-2000.
4. National Comprehensive Cancer Network (NCCN) guidelines. Available from: <http://www.nccn.org>. [cited in 2010].
5. Extermann M, Aapro M, Bernabei R, Cohen HJ, Droz JP, Lichtman S, *et al.* Use of comprehensive geriatric assessment in older cancer patients: Recommendations

- from the task force on CGA of the International Society of Geriatric Oncology (SIOG). *Crit Rev Oncol Hematol* 2005;55:241.
6. Pal SK, Katheria V, Hurria A. Evaluating the older patient with cancer: Understanding frailty and the geriatric assessment. *CA Cancer J Clin* 2010;60:120-32.
 7. Stafford RS, Cyr PL. The impact of cancer on the physical function of the elderly and their utilization of health care. *Cancer* 1997;80:1973-80.
 8. Extermann M, Aapro M. Assessment of the older cancer patient. *Hematol Oncol Clin North Am* 2000;14:63.
 9. Yancik R. Cancer burden in the aged: An epidemiologic and demographic overview. *Cancer* 1997;80:1273.
 10. Repetto L, Fratino L, Audisio RA, Venturino A, Gianni W, Vercelli M, *et al.* Comprehensive geriatric assessment adds information to Eastern Cooperative Oncology Group performance status in elderly cancer patients: An Italian Group for Geriatric Oncology Study. *J Clin Oncol* 2002;20:494-502.
 11. Welch HG, Albertsen PC, Nease RF, Bubolz TA, Wasson JH. Estimating treatment benefits for the elderly: The effect of competing risks. *Ann Intern Med* 1996;124:577-84.
 12. Wolfson C, Wolfson DB, Asgharian M, M'Lan CE, Ostbye T, Rockwood K, *et al.* A re-evaluation of the duration of survival after the onset of dementia. *N Engl J Med* 2001;344:1111-6.
 13. Gupta SK, Lamont EB. Patterns of presentation, diagnosis, and treatment in older patients with colon cancer and comorbid dementia. *J Am Geriatr Soc* 2004;52:1681.
 14. Gorin SS, Heck JE, Albert S, Hershman D. Treatment for breast cancer in patients with Alzheimer's disease. *J Am Geriatr Soc* 2005;53:1897.
 15. Newman AB, Yanez D, Harris T, Duxbury A, Enright PL, Fried LP, *et al.* Weight change in old age and its association with mortality. *J Am Geriatr Soc* 2001;49:1309-18.
 16. Grabowski DC, Ellis JE. High body mass index does not predict mortality in older people: Analysis of the Longitudinal Study of Aging. *J Am Geriatr Soc* 2001;49:968.
 17. Dewys WD, Begg C, Lavin PT, Band PR, Bennett JM, Bertino JR, *et al.* Prognostic effect of weight loss prior to chemotherapy in cancer patients: Eastern Cooperative Oncology Group. *Am J Med* 1980;69:491-7.
 18. Seeman TE, Berkman LF, Kohout F, Lacroix A, Glynn R, Blazer D. Intercommunity variations in the association between social ties and mortality in the elderly: A comparative analysis of three communities. *Ann Epidemiol* 1993;3:325-35.
 19. Avorn J, Gurwitz JH. *Geriatric Medicine*: Springer-Verlag; 1997.
 20. Lichtman SM, Villani G. Chemotherapy in the elderly: Pharmacologic considerations. *Cancer Control* 2000;7:548.
 21. Vestal RE. Aging and pharmacology. *Cancer* 1997;80:1302.
 22. Yee KW, Pater JL, Pho L, Zee B, Siu LL. Enrollment of older patients in cancer treatment trials in Canada: Why is age a barrier? *J Clin Oncol* 2003;21:1618-23.
 23. Bergman L, Kluck HM, van Leeuwen FE, Crommelin MA, Dekker G, Hart AA, *et al.* The influence of age on treatment choice and survival of elderly breast cancer patients in south-eastern Netherlands: A population-based study. *Eur J Cancer* 1992;28a:1475-80.
 24. Busch E, Kemeny M, Fremgen A, Osteen RT, Winchester DP, Clive RE. Patterns of breast cancer care in the elderly. *Cancer* 1996;78:101-11.
 25. Goodwin JS, Samet JM, Hurt WC. Determinants of survival in older cancer patients. *J Natl Cancer Inst* 1996;88:1031-7.
 26. Quoix E, Zalcman G, Oster JP, Westeel V, Pichon E, Lavolé A, *et al.* Weekly paclitaxel combined with monthly carboplatin versus single-agent therapy in patients age 70 to 89: IFCT-0501 randomized phase III study in advanced non-small cell lung cancer (NSCLC). *J Clin Oncol* 2010;28:15s Abstract 2.
 27. National cancer registry Program, ICMR Population based cancer registry reports - 2 year report 2004-2005.
 28. Cancer Incidence and mortality in Dindigul district, Tamil Nadu -2003; Dindigul ambilikai Cancer registry, Cancer Institute (W.I.A) Chennai.
 29. Population based cancer registry, Thiruvananthapuram (Urban and Rural) 2 year report 2003-2004; RCC Trivandrum Aug 2007.
 30. Cancer morbidity and Mortality in Karunagappally, Natural Background Radiation Registry 2002-2003, published in 2005.

How to cite this article: Vijaykumar DK, Anupama R, Gorasia TK, Haleema Beegum TR, Gangadharan P. Geriatric oncology: The need for a separate subspecialty. *Indian J Med Paediatr Oncol* 2012;33:134-6.
Source of Support: Nil, **Conflict of Interest:** None declared.

Announcement

iPhone App



Download FREE
**iPhone, iPad
 application**

A free application to browse and search the journal's content is now available for iPhone/iPad. The application provides "Table of Contents" of the latest issues, which are stored on the device for future offline browsing. Internet connection is required to access the back issues and search facility. The application is Compatible with iPhone, iPod touch, and iPad and Requires iOS 3.1 or later. The application can be downloaded from <http://itunes.apple.com/us/app/medknow-journals/id458064375?ls=1&mt=8>. For suggestions and comments do write back to us.