

Radiation oncology: Colors and hues

Treating cancer involves a multidisciplinary approach. Nearly 60-65% of all cancer patients will require radiotherapy during the course of their disease. This issue of the journal presents three articles on diverse aspects of radiation oncology. Over the past few decades, radiation oncology has taken huge leaps in technology.^[1] The traditional low-energy orthovoltage machines have become a curiosity. Orthovoltage machines were replaced by Cobalt machines, the latter in turn being replaced by elegant and sophisticated linear accelerators (LAs). LAs themselves have undergone serial metamorphoses. As a result, the initial machines capable of delivering simple square or rectangular 4 MV/6 MV beams have given way to the present C-arm LA, which boasts of multileaf collimators, arc treatment, asymmetric jaws and dynamic motion capabilities. Taking inspiration from computerized tomography (CT) scanners, helical rotational machines have been developed. Intensity-modulated radiotherapy and image-guided radiotherapy are considered modern radiotherapy approaches in treatment.^[2,3]

These technologies allow us to create otherwise challenging patterns of dose fluence across the target area. This allows the radiation oncologist to deliberately inhomogenize the dose intensity depending on the extent of gross or microscopic disease burden in a particular area.^[4] Treatment planning for patients with metallic hip prostheses composed of high-Z materials, however, can still be challenging due to the presence of streak artifacts from prosthetic hips in the CT dataset and inhomogeneous dose distribution within the target volume. Rana *et al.* present a dosimetric study of volumetric modulated arc therapy planning techniques for treatment of low-risk prostate cancer in patients with bilateral hip prostheses.^[5] The authors have compared three treatment plans created using rapid arc (RA) techniques. They utilized 2 arcs (2-RA), 3 arcs (3-RA) and 4 arcs (4-RA) for the 6 MV photon beam in an Eclipse treatment planning system. All three RA plans were calculated with the anisotropic analytical algorithm. They found that the plan conformity index was highest in the 2-RA plan, but the 4-RA technique reduced the doses to rectum by up to 18.8% and to bladder by up to 7.8% as compared with the 2-RA technique. This study demonstrates the intricacies of modern day radiation planning and the need to be cognizant of its nuances.

A major stride in radiation oncology has been the transition from two-dimensional to three-dimensional radiotherapy. The second article in this issue of the journal reveals the dosimetric variations one encounters while traversing from X-ray based (2 D) to CT-based (3 D) radiotherapy.^[6] The authors of this study compared the prescribed dose versus the calculated dose of spinal cord in standard head and neck irradiation assessed by the 3 D plan. They selected 42 patients with histologically proven squamous cell carcinoma at the head neck region and distributed them in two treatment protocols. In group A, 46 Gy was given in 23 fractions, followed by a tumor boost with off-cord field received 24 Gy in 12 fractions. In group B, 50 Gy was prescribed in 25 fractions initially, then off-cord field given 20 Gy in 10 fractions to analyze the outcome. The maximum dose to cord was 52.6 Gy (range 48.1-49.7 Gy) in group A and 54.3 Gy (range 51.48-52.33 Gy) in group B initially. Off-cord fields received a mean dose of 8.07 Gy (85.85% of maximum) in group A and 5.47 Gy (86.84% of maximum) in group B. At the end of 6 months from the last date of radiotherapy, grade 1 spinal cord toxicity was noted in two patients in group A and one patient in group B ($P = 0.55$). Both the groups received an additional dose that was higher than the prescribed dose, but no patient showed significant spinal cord toxicity after 6 months of follow-up. This study sends out the message of being aware of the possible pitfalls in 2 D planning.

Radiotherapy has been used in cancer patients both for radical treatments as well as for achieving palliation. The third study in the journal dwells on an important aspect of treatment in cancer care, which relates to palliation of their symptoms.^[7] Hemibody irradiation is an economical way of palliation of pain in bone metastasis in advanced cancer.^[8] The authors have reported the outcome of their patients of Hemibody irradiation with respect to pain and quality of life in cancer patients with extensive bone metastases. A total of 21 patients received lower or upper Hemibody irradiation at their center. Evaluations were performed before and 2, 4, 8, 16 and 24 weeks after treatment. Pain evaluation was performed using the Visual Analogue Scale (VAS) (10 points VAS), Verbal Rating Scale of patients at 4 points scale, Percentage of Pain Relief and Global Pain Score. Response (control of pain) was partial in 67% and complete in 22% of the patients. As such, the quality of life was better due to decreased pain and also due to lowering of the dose of analgesics. The toxicity was within acceptable limits. The average total cost of treatment, including hospital stay, medicines and radiation charges, was around INR 400.00. Such cost-effective treatments of cancer care need to be recognized and more commonly used in a resource-constrained country such as ours.

To summarize, modern day radiotherapy can be used in cancer patients for a wide variety of indications ranging

Access this article online	
Quick Response Code: 	Website: www.sajc.org
	DOI: 10.4103/2278-330X.126505

from fully curative aims to palliative treatments. What is needed is judicious use of this very powerful anticancer modality in the most cost-effective manner.^[9]

Acknowledgment

The author acknowledges the critical inputs received from Dr. B. K. Mohanti during the preparation of this manuscript.

Anusheel Munshi

Department of Radiation Oncology, Fortis Memorial Research Institute, Gurgaon, Haryana, India

Correspondence to: Dr. Anusheel Munshi,
E-mail: anusheel8@hotmail.com

References

1. Munshi A, Agarwal JP. Evolution of radiation oncology: Sharp gun, but a blurred target. *J Cancer Res Ther* 2010;6:3-4.
2. Carol MP, Peacock TM. A system for planning and rotational delivery of intensity modulated fields. *Int J Imag Sys Tech* 1995;45:21-32.
3. Vesper J, Bølle B, Wille C, Gerber PA, Matuschek C, Peiper M. Current concepts in stereotactic radiosurgery: A neurosurgical and radiooncological point of view. *Eur J Med Res* 2009;14:93-101.
4. Nicolini G, Ghosh-Laskar S, Shrivastava SK, Banerjee S, Chaudhary S,

- Agarwal JP, *et al.* Volumetric modulation arc radiotherapy with flattening filter-free beams compared with static gantry IMRT and 3D conformal radiotherapy for advanced esophageal cancer: A feasibility study. *Int J Radiat Oncol Biol Phys* 2012;84:553-60.
5. Rana SB, Pokharel S. A dosimetric study of volumetric modulated arc therapy planning techniques for treatment of low-risk prostate cancer in patients with bilateral hip prostheses. *South Asian J Cancer* 2014;3: 18-21.
6. Majumder D, Patra NB, Chatterjee D, Mallick SK, Kabasi AK, Majumdar A. Prescribed dose versus calculated dose of spinal cord in standard head and neck irradiation assessed by 3-D plan. *South Asian J Cancer* 2014;3: 22-7.
7. Pal S, Dutta S, Adhikary SS, Bhattacharya B, Ghosh B, Patra NB. Hemi body irradiation: An economical way of palliation of pain in bone metastasis in advanced cancer. *South Asian J Cancer* 2014;3: 28-32.
8. Miszczyk L, Tukiendorf A, Gaborek A, Wydmanski J. An evaluation of half-body irradiation in the treatment of widespread, painful metastatic bone disease 2008;94:813-21.
9. Munshi A. Resource-sparing and cost-effective strategies in current management of breast cancer. *J Cancer Res Ther* 2009;5: 116-20.

How to cite this article: Munshi A. Radiation oncology: Colors and hues. *South Asian J Cancer* 2014;3:16-7.

Source of Support: Nil. **Conflict of Interest:** None declared.

News

Best Of ASCO India 2014.
Organizing Chairman : Prof. Raghunadhras D.,
Organizing Secretary & Program Director : Dr. Senthil Rajappa
27th - 29th June, 2014
Web: www.bestofascoindia.com
Mail: indiabestofasco@gmail.com

News

31st ICON Conference.
Organizing Secretary : Dr. T. P. Sahoo,
12th - 14th September, 2014
Web: www.iconconferences.com
Mail: 31sticon@gmail.com

Author Help: Reference checking facility

The manuscript system (www.journalonweb.com) allows the authors to check and verify the accuracy and style of references. The tool checks the references with PubMed as per a predefined style. Authors are encouraged to use this facility, before submitting articles to the journal.

- The style as well as bibliographic elements should be 100% accurate, to help get the references verified from the system. Even a single spelling error or addition of issue number/month of publication will lead to an error when verifying the reference.
- Example of a correct style
Sheahan P, O'leary G, Lee G, Fitzgibbon J. Cystic cervical metastases: Incidence and diagnosis using fine needle aspiration biopsy. *Otolaryngol Head Neck Surg* 2002;127:294-8.
- Only the references from journals indexed in PubMed will be checked.
- Enter each reference in new line, without a serial number.
- Add up to a maximum of 15 references at a time.
- If the reference is correct for its bibliographic elements and punctuations, it will be shown as CORRECT and a link to the correct article in PubMed will be given.
- If any of the bibliographic elements are missing, incorrect or extra (such as issue number), it will be shown as INCORRECT and link to possible articles in PubMed will be given.