

Letter to the Editor

Late effects of cancer treatment in breast cancer survivors

DOI: 10.4103/2278-330X.175956

Dear Editor,

We have read the recent review on “late effects of cancer treatment in breast cancer survivors” with great interest. Agrawal *et al.* noted that cardiac toxicity, reproductive dysfunction, pneumonitis, arm lymphedema, neuropathy, and skin changes are examples of the wide range of complications associated with adjuvant treatment of breast cancer. Although many co-morbidities are discussed in the review by Agrawal; we would like to draw attention to a neglected complaint after breast cancer treatment, namely breast edema. Breast edema is a morbidity of breast cancer treatment which is often underdiagnosed in clinical practice. Currently, there is no consensus on the definition of breast edema and standardized assessment criteria. Common criteria found in the literature are an increased volume of the breast, peau d’orange, heaviness of the breast, redness of the skin, breast pain, skin thickening, hyperpigmented skin pores, and a positive pitting sign.^[1,2] In some cases, the breast size can increase by more than one cup size. It is demonstrated that breast edema is a common morbidity in women who underwent breast-conserving surgery and radiotherapy. The breast edema incidence is very broad, namely 0-90.4%.^[3,4] Several factors are responsible for this broad range such as no standard assessment method, no uniform definition of breast edema, different types of radiotherapy, and different follow-up times or measuring intervals. Breast edema can occur because of either the breast surgery or the radiation therapy. Both treatments can disturb the lymphatic circulation of the breast. In most patients breast edema develops during radiation therapy. Some studies, however, describe late-onset breast edema, although these cases are rare.^[5] Late breast edema occurs about 20 months after breast

surgery and/or radiation therapy.^[5] Some patients still suffer from this complaint years after their initial treatment and breast edema has a negative impact on the quality of life. Therefore, further research is warranted. The information on breast edema is additional to the late morbidities found by Agrawal *et al.*

Hanne Verbelen^{1,2}, Nick Gebruers^{1,2}, Wiebren Tjalma^{2,3}

¹Department of Rehabilitation Sciences and Physiotherapy, Faculty of Medicine and Health Sciences, University of Antwerp, Universiteitsplein 1, 2610 Antwerp, ²Multidisciplinary Breast Clinic, Gynecological Oncology Unit, Antwerp University Hospital, Wilrijkstraat 10 2650 Edegem, ³Faculty of Medicine and Health Sciences, University of Antwerp, Universiteitsplein 1, 2610 Antwerp, Belgium

Correspondence to: Dr. Nick Gebruers,
E-mail: nick.gebruers@uantwerpen.be

References

1. Adriaenssens N, Verbelen H, Lievens P, Lamote J. Lymphedema of the operated and irradiated breast in breast cancer patients following breast conserving surgery and radiotherapy. *Lymphology* 2012;45:154-64.
2. Pezner RD, Patterson MP, Hill LR, Desai KR, Vora N, Lipssett JA. Breast edema in patients treated conservatively for stage I and II breast cancer. *Int J Radiat Oncol Biol Phys* 1985;11:1765-8.
3. Mayo C, Lo YC, Fitzgerald TJ, Urie M. Forward-planned, multiple-segment, tangential fields with concomitant boost in the treatment of breast cancer. *Med Dosim* 2004;29:265-70.
4. Adriaenssens N, Belsack D, Buyl R, Ruggiero L, Breucq C, De Mey J, *et al.* Ultrasound elastography as an objective diagnostic measurement tool for lymphoedema of the treated breast in breast cancer patients following breast conserving surgery and radiotherapy. *Radiol Oncol* 2012;46:284-95.
5. Rönkä RH, Pamilo MS, von Smitten KA, Leidenius MH. Breast lymphedema after breast conserving treatment. *Acta Oncol* 2004;43:551-7.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

How to cite this article: Verbelen H, Gebruers N, Tjalma W. Late effects of cancer treatment in breast cancer survivors. *South Asian J Cancer* 2015;4:182.