

Knowledge about imaging modalities, risks, and protection in radiology among medical students at the University of Hail

Meshael N. Alreshidi¹, Dalal Alshubrmi², Fayez Alreshidi², Khaled Soliman³, Ibrahim Alrashidi⁴

¹College of Medicine, University of Hail, Hail, Kingdom of Saudi Arabia, ²Assistant professor, Department of Family and Community Medicine, College of Medicine, University of Hail, Hail, Kingdom of Saudi Arabia, ³Consultant medical Physicist, Medical Physics Department, Prince Sultan Military Medical City, Riyadh, Kingdom of Saudi Arabia, ⁴Interventional Radiologist, Radiology, Prince Sultan Military Medical City, Riyadh, Kingdom of Saudi Arabia

Access this article online

Website: www.avicennajmed.com

DOI: 10.4103/ajm.ajm_49_19

Quick Response Code:



ABSTRACT

Aim: The aim of this study was to evaluate awareness and knowledge about radiation risks and safety principles among medical students at the College of Medicine, University of Hail, Hail, Saudi Arabia, in their clinical years. **Materials and Methods:** In this cross-sectional study, an anonymous electronic questionnaire was sent to 174 randomly selected students in clinical years 4–6. The questionnaire contained 38 questions. The respondents' answers to these questions were used to classify them according to their demographic characteristics and to evaluate their knowledge about common imaging modalities, radiation risks, and safety measures. The data were analyzed using the Statistical Package for the Social Sciences (SPSS) software, version 22. **Results:** Seventy-five (51.7%) of 145 respondents were female and 70 (48.3%) were male. Fifty-five respondents (37.9%) were in year 4, 38 (26.2%) were in year 5, and 52 (35.9%) were in year 6. The mean score for knowledge about common imaging modalities was 4.10 ± 2.030 of 10, that for knowledge about the risks of radiation was 3.17 ± 1.954 (range, 0–8) of 13, and that for knowledge about radiation protection measures was low at 0.79 ± 0.922 (range, 0–4) of 8. Overall, there was an improvement in knowledge about the imaging modalities and the risks of radiation as the number of clinical years increased ($P = 0.000$), but it was still unsatisfactory. **Conclusion:** The results of this study indicate that the medical students at the University of Hail have very limited knowledge about radiation risks and safety measures. These findings highlight the need for urgent action to improve students' knowledge of these topics.

Key words: Awareness, medical students, patient safety, radiation risk, radiation protection

INTRODUCTION

Medical imaging has become an important diagnostic tool in the medical field. However, the radiation dose received from some imaging modalities increases the lifetime risk of cancer.^[1] Moreover, as mentioned in the Biological Effects of Ionizing Radiation Report, the exposure to a radiation dose of 100 mSv has an estimated lifetime risk of radiation-induced cancer of one per 100,^[2] as well as tissue reactions, including hair loss, skin rashes, and ulceration as a result of accidental exposure to high doses.^[3] Furthermore, *in utero* exposure

to radiation can lead to generalized growth retardation, mental disability, and seizures, as well as decreased school performance.^[4] Fortunately, these risks can be minimized by a principle called optimization and by understanding and implementing radiation protection principles.^[5]

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Alreshidi MN, Alshubrmi D, Alreshidi F, Soliman K, Alrashidi I. Knowledge about imaging modalities, risks, and protection in radiology among medical students at the University of Hail. *Avicenna J Med* 2020;10:15-21.

Address for correspondence: Meshael Alreshidi, Intern, College of Medicine, University of Hail, Hail, Kingdom of Saudi Arabia.
E-mail: me.shaal2010@hotmail.com

Many studies conducted worldwide, but only three in Saudi Arabia, have concluded that medical students and even radiology residents lack adequate knowledge about the principles of radiation protection.^[6-11] However, no studies on this subject have been performed in the Hail region, Saudi Arabia. The aim of this study was to evaluate awareness and knowledge of the radiation risks and safety principles among medical students at the University of Hail in their clinical years.

MATERIALS AND METHODS

This study was performed at the University of Hail in 2018, at which time, 244 medical students in their clinical years of study (years 4–6) were enrolled. Hail is a city in Saudi Arabia that has a population of 699,774.^[12] Seventy-nine students (39 female and 40 male) were in year 4, 83 (39 female and 44 male) were in year 5, and 82 (38 female and 44 male) were in year 6. We estimated the study sample size that needed to be enrolled from the total student population of 244 by Raosoft sample size calculator, Raosoft inc, Seattle, USA. It was determined that 150 students would be an appropriate sample size with a confidence interval of 95%. We added 16% ($n = 24$) to this figure to allow for nonresponders.^[13] We used random numbers generated online by a research randomizer website.^[14] One hundred and seventy-four students were selected at random and approached to participate in this prospective cross-sectional study, which involved completing a one-time, self-administered anonymous online questionnaire between August 2 and August 5, 2018.

Ethical approval to conduct this study was obtained from the ethics committee at the College of Medicine, University of Hail (approval number EC-00025/CM/UOH.11/18). Agreement to complete the anonymized online questionnaire was accepted as willingness to participate in the study.

Data collection

The questionnaire contained 38 multiple-choice and true or false questions and was divided into four sections. The first section contained items on demographic characteristics (i.e., gender, clinical year, nationality, and marital status) and three questions regarding the respondent's opinion of their knowledge of this topic. The second section contained items that tested the respondent's knowledge about the various imaging modalities and the differences between them, with additional questions regarding effective radiation doses. The third section contained items that tested knowledge about radiation risks and the contraindications of some imaging modalities. The fourth section tested knowledge about radiation safety principles. The validity of the questionnaire was confirmed by a radiologist and a medical physicist [Appendix 1].

We calculated the score for each respondent by awarding one mark for each correctly answered question, for a total possible score of 31. The second section was scored of 10, the third section of 13, and the fourth section of 8.

Statistical analysis

The data were shown as frequency and percentage. The categorical data were compared between subgroups using the chi-square test. The mean scores were compared between the students in the three clinical years using analysis of variance (ANOVA). All statistical analyses were performed using the Statistical Package for the Social Sciences (SPSS) software, for Windows (version 22; IBM, Armonk, New York). P values <0.05 were considered statistically significant.

RESULTS

One hundred and forty-five students (75 [51.7%] female and 70 [48.3%] male) returned completed questionnaires. One hundred and forty students (96.6%) were single, four (2.8%) were married, one (0.7%) was divorced, and all were Saudi nationals. The sample distribution according to clinical year and gender is shown in Table 1.

Fifteen respondents (10.3%) had attended a radiation protection course. Two (13.3%) of these 15 students were in year 4, two (13.3%) were in year 5, and 11 (73.3%) were in year 6. A statistically significant difference was observed in the likelihood of attending a radiation protection course as students moved from clinical year 5 to 6 ($P = 0.006$, chi-square test) but not according to gender ($P = 0.337$, chi-square test). The students who attended a radiation protection course had significantly higher scores in the radiation safety principles section ($P = 0.015$, independent samples t test, two-tailed) but not in the other sections [Table 2].

Forty-five respondents (31%) agreed that they had adequate information about the risks of radiation, 18 (12.4%) disagreed, and 82 (56.6%) were unsure. A statistically significant difference was observed in the responses to this question according to the clinical year of study ($P = 0.010$,

Table 1: Sample distribution according to clinical year and gender

Clinical year	Gender	Participants, n	Percentage
4	Female	24	16.6
	Male	31	21.4
	Total	55	37.9
5	Female	22	15.2
	Male	16	11.0
	Total	38	26.2
6	Female	29	20.0
	Male	23	15.9
	Total	52	35.9

chi-square test) but not according to gender ($P = 0.396$, chi-square test). Students in year 6 were more confident that they had adequate information about radiation risks than those in the earlier clinical years.

Sixteen respondents (11%) agreed that they were adequately informed about radiation protection measures, 71 (49%) disagreed, and 58 (40%) were unsure. Of 75 female students, 6 (8%) agreed that they were adequately informed, 46 (61.3%) disagreed, and 23 (30.7%) were unsure. Of 70 male students, 10 (14.3%) agreed that they were adequately informed, 25 (35.7%) disagreed, and 35 (50%) were unsure. A statistically significant difference was observed in the responses to this question according to gender ($P = 0.009$, chi-square test) but not according to the clinical year of study ($P = 0.309$, chi-square test).

A statistically significant difference was observed in the overall scores between students in years 4, 5, and 6 ($F_{2,142} = 29.170$, $P = 0.000$, ANOVA). *Post hoc* comparisons using Tukey's honestly significant difference test indicated that the mean score for year 6 was significantly higher than that for years 4 and 5 ($P = 0.000$). However, no significant difference was observed in the scores between years 4 and 5 ($P = 0.629$). The scores are shown in Table 3.

DISCUSSION

This cross-sectional study evaluated the level of awareness and knowledge about common imaging modalities, the

risks of radiation, and radiation protection measures among medical students in their clinical years of study. According to its results, only 10.3% of students had attended a radiation protection course. Statistically significant differences were observed in course attendance rates between the three clinical years, with the highest attendance rate of 73.3% reported by students in year 6. The students who attended a radiation protection course had significantly higher scores in the radiation safety principles section ($P = 0.015$, independent samples *t* test, two-tailed). This result indicates that attendance at a radiation protection course can significantly improve students' knowledge about radiation safety and is consistent with the finding of a study by Hagi and Khafaji.^[15]

In general, the level of awareness among these students was poor, with statistically significant differences in mean scores between the clinical years of study. Overall, 31% considered that they had adequate knowledge about the risks of radiation and 11% reported that they had adequate knowledge about radiation protection measures.

We found that awareness of the imaging modalities and the risks of radiation improved as the number of clinical years increased, which is in agreement with a report by O'Sullivan *et al.*,^[16] who found that medical students' awareness of radiation exposure improved in their final years of medical school but it was still unsatisfactory. The study by Hagi and Khafaji^[15] yielded similar results, and the authors concluded that the subject of radiation safety needed to be addressed at the undergraduate level.

Table 2: Relationship between attendance of a radiation protection course and knowledge about radiation risks and safety principles

Section	Course attendance	Mean	95% Confidence interval		Mean difference	P value
			Lower	Upper		
Overall results	Yes	8.80	7.16	10.44	0.831	0.450
	No	7.97	7.25	8.68		
Knowledge about different imaging modalities	Yes	3.87	2.84	4.89	0.264	0.635
	No	4.13	3.77	4.49		
Knowledge about radiation risks	Yes	3.60	2.69	4.51	0.485	0.365
	No	3.12	2.77	3.46		
Knowledge about radiation safety principles	Yes	1.33	0.75	1.91	0.610	0.015
	No	0.72	0.57	0.88		

Table 3: Mean scores according to clinical years

Clinical year	Overall results	Knowledge about imaging modalities	Knowledge about radiation risks	Knowledge about radiation safety principles
4	Mean	6.18	3.44	2.24
	SD	3.59	1.98	1.66
5	Mean	6.84	3.63	2.66
	SD	3.37	1.71	1.77
6	Mean	10.92	5.15	4.52
	SD	3.22	1.88	1.60

SD = standard deviation

In our study, the lowest score was found in the section on knowledge about safety principles in radiology; the mean score of 0.79 ± 0.922 of 8 indicates that the students had a very low level of knowledge regarding how to protect themselves and their patients in the radiology field, which is similar to previous reports, and signals a very serious patient and occupational safety issue.^[6-11]

An important limitation of this study was that it targeted knowledge about radiation protection among medical undergraduates in their clinical years of study. Further studies are required to assess the knowledge of interns.

In summary, the findings of this study indicate that medical students at the University of Hail, Hail, Saudi Arabia, have very limited knowledge about radiation risks and safety measures. These results highlight the need for urgent action to improve students' knowledge about these important issues.

Acknowledgement

We are grateful to Ahmed Hameed Alreshidi for his assistance with the collection of the data used in this study.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Sheppard JP, Nguyen T, Alkhalid Y, Beckett JS, Salamon N, Yang I. Risk of brain tumor induction from pediatric head CT procedures: A systematic literature review. *Brain Tumor Res Treat* 2018;6:1-7.
- National Research Council. *Health Risks from Exposure to Low Levels of Ionizing Radiation: BEIR VII*. Washington, DC: The National Academies Press; 2006.
- Parikh JR, Geise RA, Bluth EI, Bender CE, Sze G, Jones AK, *et al.* Potential radiation-related effects on radiologists. *Am J Roentgenol* 2017;208:595-602.
- Verreet T, Verslegers M, Quintens R, Baatout S, Benotmane MA. Current evidence for developmental, structural, and functional brain defects following prenatal radiation exposure. *Neural Plast* 2016;2016:1243527.
- Do K-H. General principles of radiation protection in fields of diagnostic medical exposure. *J Korean Med Sci* 2016;31:6-9.
- Algohani KA, Aldahhasi AA, Algarni AH, Amrain KY, Marouf MA. Awareness of radiation protection measures among radiologists and non-radiologists. *Egypt J Hosp Med* 2018;70:371-5.
- Salih S, Zeidan ZA, Alzabalani A, Albadrani MS, Yousef M. Awareness and knowledge towards ionizing radiation hazard among medical students, interns and residents in Al-Madinah Al-Munawarah, KSA. *Life Sci J* 2014;11:6-10.
- Zhou GZ, Wong DD, Nguyen LK, Mendelson RM. Student and intern awareness of ionising radiation exposure from common diagnostic imaging procedures. *J Med Imaging Radiat Oncol* 2010;54:17-23.
- Lee RK, Chu WC, Graham CA, Rainer TH, Ahuja AT. Knowledge of radiation exposure in common radiological investigations: A comparison between radiologists and non-radiologists. *Emerg Med J* 2012;29:306-8.
- AlDossari HM, Aldosary A, Alrashed A, Mohammed H. Assessing the level of knowledge about radiation dose in common. *Int J Pharm Res Allied Sci* 2018;7:114-8.
- Ahmad A, Saleh J, Alhawiti AE, Aljehani EA, Alblowi A, Albalawi SM, *et al.* Radiation awareness among sixth-year medical students and interns at University of Tabuk, Saudi Arabia. *BRJMCS* 2017;6:24-9.
- The General Population and Housing Census. *Population Characteristics Survey 2017*. Available from: https://www.stats.gov.sa/sites/default/files/population_characteristics_surveysar.pdf. [Last accessed on 2018 Jun 5].
- Sample Size Calculator by Raosoft. Available from: <http://www.raosoft.com/samplesize.html>. [Last accessed on 2018 Jun 5].
- Research Randomizer. Available from: <https://www.randomizer.org/>. [Last accessed on 2018 Jun 5].
- Hagi SK, Khafaji MA. Medical students' knowledge of ionizing radiation and radiation protection. *Saudi Med J* 2011;32:520-4.
- O'Sullivan J, O'Connor OJ, O'Regan K, Clarke B, Burgoyne LN, Ryan MF, *et al.* An assessment of medical students' awareness of radiation exposures associated with diagnostic imaging investigations. *Insights Imaging* 2010;1:86-92.

APPENDIX 1

Awareness and Knowledge of Clinical Phase Medical Students at Hail University about Common Imaging Modalities, Radiation Risk, and Protection Applied in Radiology

We are medical students in the college of medicine at University of Hail, conducting a research study about the awareness and knowledge of Hail University clinical years' medical students about radiation risks and safety measures. Our objective is evaluation of awareness and knowledge of these students to assess the need for basic radiation safety course.

You have been chosen randomly through research randomizer website although this, your participation in this study, is voluntary. Your participation is very important to us and remember your response will be kept confidential, only the research team involved in this study will read your response. Filling out the questionnaire will take only few minutes.

For any inquiry, please contact us via me.shaal2010@hotmail.com

I have read the above information and I consent to take part in the study

1st section

- Q1. Gender:
- Female
 - Male
- Q2. Which clinical year you're in?
- 4th year
 - 5th year
 - 6th year
- Q3. Nationality:
- Saudi
 - Non-Saudi
- Q4. Marital status:
- single
 - married
 - divorced
 - widow
- Q5. Have you attended a radiation protection course?
- Yes, I have
 - No, I have not
- Q6. Do think that you have adequate information about radiation risks?
- yes

- No
- Maybe

Q7. Do think that you have adequate information about radiation protection measures?

- yes
- No
- Maybe

2nd section

Q8. A medical imaging technology that combines multiple X-ray projections taken from different angles to produce detailed cross-sectional images of areas inside the body

- CT
- MRI
- Ultrasound
- X-ray
- I do not know

Q9. A medical imaging technology that uses radio waves and a magnetic field to create detailed images of organs and tissues

- CT
- MRI
- Ultrasound
- X-ray
- I do not know

Q10. A medical imaging technology that uses high-frequency sound waves to create images of the inside of the body

- Nuclear medicine
- MRI
- Ultrasound
- X-ray
- I do not know

Q11. The oldest and most commonly used form of medical imaging that uses ionizing radiation to produce images of the internal structure

- Nuclear medicine
- MRI
- Ultrasound
- X-ray
- I do not know

Q12. A medical imaging technology that uses a radioactive material (radiopharmaceutical) to produce images of the internal structure

- CT
- MRI
- Ultrasound
- Nuclear medicine
- I do not know

- Q13. All the following imaging modalities use ionizing radiation except:
- CT
 - X-ray
 - Ultrasound
 - Mammography
 - I do not know
- Q14. What is the approximate effective radiation dose from a chest x-ray?
- 0.1 mSv
 - 0.9 mSv
 - 1 mSv
 - 3 mSv
 - I do not know
- Q15. Chest X-ray when compared to natural background radiation is equal to:
- 10 days
 - 2 months
 - 1 year
 - 3 years
 - I do not know
- Q16. What is the approximate effective radiation dose of abdomen and pelvis computed tomography (CT) scan?
- 10 mSv
 - 20 mSv
 - 30 mSv
 - 50 mSv
 - I do not know
- Q17. What is the approximate effective radiation dose from a Computed Tomography (CT) exam of the Head?
- 1mSv
 - 2 mSv
 - 3 mSv
 - 4 mSv
 - I do not know
- Q19. Which of the following tissues is more radiosensitive to ionizing radiation damage?
- Kidney
 - Breast
 - Liver
 - Muscle
 - I do not know
- Q20. Which of the following diseases may be a result of stochastic effects of exposure to ionizing radiation?
- Dermatitis
 - Leukemia
 - Cataract
 - All of the above
 - I do not know
- Q21. As a result of eye exposure to radiation the patient might be at risk of which of the following:
- Glaucoma
 - cataract
 - optic neuritis
 - corneal ulcers
 - I do not know
- Q22. At which of the following periods the fetus is more sensitive to radiation:
- 2nd week until 18th week
 - 4th week until the 20th week
 - 5th week until 14th week
 - 6th week until 16th week
 - I do not know
- Q23. Deterministic effects of radiation exposure during pregnancy depend only on the radiation dose
- True
 - False
 - I do not know
- Q24. Pregnant women should avoid **all types of medical imaging**
- true
 - false
 - I do not know
- Q25. Is it safe for pregnant women to have a mammography?
- yes
 - No
 - I do not know
- Q26. Hair loss is one of the localized short-term injuries that might happen after radiation exposure. At which dose you will expect it to be a result of the radiation exposure?
- 1 Gy
 - 2 Gy
 - 3 Gy

3rd section

- Q18. Regarding the risk of cancer as long-term effect of radiation exposure, which statement of the following is true:
- cancer will not always occur, but its likelihood is proportional to the radiation dose
 - cancer will not always occur, and its likelihood is **not** proportional to the radiation dose
 - Cancer risk is one of the deterministic effects of radiation exposure
 - Cancer after radiation exposure occurs in children only
 - I do not know

- 4 Gy
 - I do not know
- Q27. Cutaneous necrosis is one of the localized short-term injuries that might happen after radiation exposure. At which dose you will expect it to be a result of the radiation exposure?
- 30 Gy
 - 40 Gy
 - 50 Gy
 - 60 Gy
 - I do not know
- Q28. The dose threshold for acute radiation syndrome is about:
- 1 Sv
 - 3 Sv
 - 6 Sv
 - 10 Sv
 - I do not know
- Q29. Any metal device is considered as a contraindication to use in which of the followings medical imaging modalities?
- CT scan
 - MRI
 - ultrasonography
 - X-ray
 - I do not know
- Q30. The metformin should be withheld at the time of intervenous contrast administration in patients who are known to have kidney disease and discontinued for
- 12 h afterward
 - 24 h afterward
 - 48 h afterward
 - 72 h afterward
 - I do not know
- 4th section**
- Q31. Which of the following substances are used to coat the walls of a CT scan room for radiation shielding?
- Tungsten
 - Glass
 - Lead
 - Iron
 - I do not know
- Q32. Dosimeter is:
- a device that measures exposure to ionizing radiation
 - a device that measures the distance from the source of ionizing radiation
 - a device that provides physical protection from ionizing radiation
 - a device that measures the safe area
 - I do not know
- Q33. Which one of the International Commission on Radiological Protection (ICRP) principles can't be applied in medical radiation exposure?
- justification
 - Dose limit
 - optimization
 - All of the above
 - I do not know
- Q34. As low as reasonably achievable (ALARA) principle is equivalent to which one of the following ICRP principles?
- justification
 - Dose limit
 - optimization
 - None of the above
 - I do not know
- Q35. All the following are methods used to reduce the amount of exposure to ionizing radiation except:
- Time
 - Distance
 - Source
 - None of the above
 - I do not know
- Q36. How far from the X-ray, should you stand without any protection during the radiological-guided procedure (e.g.,C-arm)?
- 1 m
 - 2 m
 - 5 m
 - 8 m
 - I do not know
- Q37. What is the annual effective dose limit for occupational exposure?
- 10 mSv
 - 20 mSv
 - 30 mSv
 - 40 mSv
 - I do not know
- Q38. The use of diagnostic reference levels is an important tool in which of the following ICRP radiation protection principles
- justification
 - Dose limit
 - optimization
 - None of the above
 - I do not know