Letters to Editor

Re: A novel method to insert drain atraumatically after liposuction in gynecomastia

Sir.

We read with great interest the article titled, 'A novel method to insert drain atraumatically after liposuction in gynaecomastia' by Sunil Gaba, and we congratulate the authors.^[1]

We agree with the authors that, also liposuction, even when combined with glandular resection, has an important role in the surgical treatment of gynaecomastia, for flattening the thorax, eliminating fat residuals and creating a controlled shrinking of the cutaneous flat to recontour the male chest area. Seromas is one of the most common complications when liposuction is included in gynaecomastia surgical correction.^[2]

In our practice, with more than 600 patients undergoing gynaecomastia surgical correction and long-term evaluation for complications and recurrences, mainly based on direct excision of the gland, the inferior emi-periareolar access, allowing a direct view of the surgical field, ensures a valid control of haemostasis, reducing the incidence of haematomas significantly.^[3]

Furthermore, we retain that quilting stitches, firmly connecting the adipo-cutaneous thoracic flap to the fascial plane, with a compressive medical dressing, effectively contributes in reducing seromas and haematomas. Moreover, the compression is maintained, for at least 1 month postoperatively, using a compressive jersey.^[4,5]

The use of drains should be suggested only in patients with the removal of large amount of breast gland or with a personal history of coagulation disorder, although the final choice for their insertions should be made directly in the operation room and last for not longer than 48 h. In fact, we do not retain the use of drains mandatorily for this type of surgery, even because the use of suction drains themselves could be responsible for complications such as surgical-site infections and pathological scarring which might be the cause of patients' complaints, especially in aesthetic surgery.

Financial support and sponsorship Nil.

Conflicts of interest

There are no conflicts of interest.

Alessandro Innocenti, Dario Melita, Marco Innocenti

Plastic and Reconstructive Microsurgery, Careggi University Hospital, Florence, Italy

Address for correspondence:

Dr. Alessandro Innocenti, Viale Giacomo Matteotti 42, 50132 Firenze, Italia. E-mail: innocentialessandro@alice.it

REFERENCES

- Gaba S, Ajai K S, Sharma RK, Chetan L. A novel method to insert drain atraumatically after liposuction in gynaecomastia. Indian J Plast Surg 2018;51:102-3.
- Innocenti A, Melita D, Mori F, Ciancio F, Innocenti M.Management of Gynecomastia in Patients With Different Body Types: Considerations on 312 Consecutive Treated Cases. Ann Plast Surg. 2017;78:492-6.
- Innocenti A, Ciancio F, Portincasa A, Parisi D. Discussion: Surgical Management of Gynecomastia-Subcutaneous Mastectomy and Liposuction. Aesthetic Plast Surg. 2017;41:983-8.
- Innocenti A, Andretto Amodeo C, Ciancio F. Wide-undermining neck liposuction: tips and tricks for good results. Aesthetic Plast Surg. 2014;38:662-9.
- Innocenti A, Ciancio F, Francesco M, Melita D, Innocenti M. Comment to "No-Drain Single Incision Liposuction Pull-Through Technique for Gynecomastia". Aesthetic Plast Surg. 2017;41:990-1.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Access this article online Quick Response Code: Website: www.ijps.org DOI: 10.4103/ijps.IJPS_106_18

How to cite this article: Innocenti A, Melita D, Innocenti M. Re: A novel method to insert drain atraumatically after liposuction in gynecomastia. Indian J Plast Surg 2018;51:342-3.

© 2019 Indian Journal of Plastic Surgery | Published by Wolters Kluwer - Medknow