

# Sleep and the aging brain. A multifaceted approach

Angeliki Tsapanou \*  
Nikolaos Scarmeas  
Yaakov Stern

Cognitive Neuroscience Division,  
Department of Neurology, Columbia  
University Medical Center - New York -  
New York.

## ABSTRACT

In the current review we provide a theoretical background on studies examining the association between sleep and brain function. We focus on the association between sleep and cognitive performance, cognitive changes over time and incident dementia as well. We then present some data on the link between sleep and subjective cognitive complaints, in participants without any objective clinical cognitive decline. We conclude with investigating the association between sleep and brain biomarkers, by highlighting the importance of specific genes and specific brain regions' morphometry. The role of sleep is vital in maintaining a healthy aging brain, and multiple factors should be taken under account when investigating this association.

**Keywords:** Sleep; Cognition; Dementia; Brain; Aging.

## \*Corresponding author:

Angeliki Tsapanou  
E-mail: at2859@cumc.columbia.edu

Received: October 13, 2019;  
Accepted: November 7, 2019.

DOI: 10.5935/1984-0063.20190128

Changes in sleep patterns are common among the aging population. Approximately 5% of older adults meet the criteria for clinically significant insomnia disorders and 20% for sleep apnea syndromes<sup>1</sup>. Self-reported sleep problems seem to reflect poor overall quality of sleep, which in turn has been associated with changes in cognition<sup>2-6</sup>. Some cross-sectional studies have reported that sleep problems, especially daytime sleepiness, are related to poor cognitive function<sup>7-9</sup>. For example, excessive daytime sleepiness has been associated with impaired memory, orientation, and attention<sup>8</sup>. Narcolepsy also, which is linked to daytime sleepiness, has been associated with executive control deficits in different studies, however, as most of the narcoleptic patients are under treatment, it is difficult to clarify the directionality and the exact power of the results<sup>10</sup>.

Although a number of cross-sectional studies have investigated the relationship between sleep problems and cognitive functioning in older adults, there is a relative paucity of longitudinal research on this relationship<sup>11-16</sup>. From the few longitudinal studies that do exist, findings suggest that self-reported daytime sleepiness, in particular, is a major risk factor for<sup>11</sup>, and a possible early marker of cognitive decline<sup>12</sup>. Some results of the existing longitudinal literature, however, are contradictory. No longitudinal association between self-reported sleep duration, difficulty sleeping, and snoring with cognitive function in older women was reported in one study<sup>17</sup>. Also, on a different study, results about the association between sleep problems and cognitive decline did not survive after controlling for depression<sup>11</sup>.

Many of the previous longitudinal studies have some apparent research gaps. For example, it is unclear whether sleep problems are related to specific cognitive abilities, as most of the existing studies used short global measures of cognitive function<sup>13,16,18</sup> or a telephone-based screening battery<sup>19,21</sup>. One large longitudinal study that did investigate the association of sleep with a specific cognitive domain (i.e. memory) used a single memory test (i.e. Delayed Word Recall Test) in addition to the Mini Mental State Examination (MMSE)<sup>22</sup>, which may not provide a complete picture of cognitive function. Furthermore, many of the previous longitudinal studies did not include a comprehensive clinical evaluation. Some of the studies had also a relatively small sample size<sup>13,23,24</sup> or used specific sex-groups, i.e. only male participants<sup>15,25</sup>, limiting the generalizability of the results.

Our group examined the association between self-reported sleep problems and cognition, both cross-sectionally and longitudinally in a large sample of older adults. Sleep was measured by the Medical Outcomes Study-Sleep Scale (MOS-SS), while cognition was examined through an extensive neuropsychological evaluation. Results indicated that daytime sleepiness was linked to slow speed of processing cross-sectionally but also after a 3.2 years follow-up<sup>26</sup>. This relationship did not change even after excluding demented participants at baseline, indicating that specific sleep problems such as increased daytime sleepiness can be an early sign of cognitive decline in the elderly. We also examined the above cross-sectional association in a different group of older adults, with a different geographical background

(Greece) but with a similar study design. Poor sleep quality, as a general sleep problems variable, and long sleep duration appeared to be associated with poor memory performance in older adults, even after excluding the participants with dementia or Mild Cognitive Impairment (MCI)<sup>27</sup>.

After examining the association between sleep and cognition/cognitive changes over time, a question that arises is what is the relationship between sleep problems and incident dementia. Daytime sleepiness has been associated with incident dementia in several longitudinal studies<sup>19,28-30</sup>. Another study found that daytime sleepiness predicted vascular dementia in a sample of older men<sup>30</sup>. In addition, 'sleep fragmentation' (high activity during sleep) has been linked to incident Alzheimer's disease (AD)<sup>23</sup> in older adults. Prolonged sleep duration has been also linked to an increased risk of dementia in a large population based study<sup>31</sup>. A specific meta-analysis suggests insomnia to be a risk factor for AD, while Sleep-Disordered Breathing (SDB) to be a risk factor for not only AD, but also vascular and all-cause dementia<sup>32</sup>. Examination of the above association in our large multiethnic group of cognitively normal older adults, showed that 7.2% of the participants became demented in a 3-year follow-up. Daytime sleepiness and sleep inadequacy appeared to be significant risk factors for developing dementia, even after adjusting for multiple demographic factors and comorbidities<sup>33</sup>.

Based on the existing literature, we conclude that sleep problems are associated with cognition, cognitive changes over time and incident dementia as well. Multiple clinical, neurobiological, and demographic factors could play a significant role to these links. Disrupted circadian rhythms could affect both sleep regulation and cognitive performance<sup>34,35</sup>. Amyloid- $\beta$  deposition, especially in cortical areas and the precuneus, is another factor which could affect the relationship between sleep and cognition/ risk for dementia<sup>36,37</sup>. We could speculate that sleep problems are causing cognitive decline, however, reverse causality could be also true; sleep dysregulation could be an early symptom of an upcoming neurodegeneration.

After examining the relationship between sleep and objective cognitive decline, it is plausible to examine any possible association between sleep and subjective cognitive decline (SCD), which might be an early indicator of upcoming cognitive changes. Of the studies that have examined sleep in relation to SCD, one showed that sleep loss was associated with increased severity of SCD<sup>38</sup>. Poor sleep quality and daytime sleepiness were also linked to SCD in older adults<sup>39,40</sup>. Our group examined this relationship in two different cohorts—the ones mentioned before showing the association between sleep and objective cognitive changes-, and results revealed links between sleep and SCD even after controlling for objective cognitive status<sup>41</sup>. At any given level of objective cognition, sleep disturbance is accompanied by subjective cognitive impairment. Personality, mood, and attitudinal factors could significantly affect this association.

As we could now consider the exploration of the links between sleep and cognition quite established, the main interest transfers to the possible neurobiological connections underlying these associations. Several studies have suggested

a familial aggregation of obstructive sleep apnea/ hypopnea (OSA/H)<sup>42,43</sup>, snoring, and daytime hypersomnolence<sup>44</sup>. Results from a large longitudinal twin study showed that insomnia is moderately heritable (14%-38%)<sup>45</sup>, suggesting that there might be a strong genetic contribution to sleep disturbances. Apolipoprotein E gene (*ApoE*) is known as a major risk factor for late-onset AD<sup>46,47</sup>.

However, studies assessing whether *ApoE* gene may contribute to the risk of disturbances in sleep have been scarce, and results are often contradictory. One study suggests that apnea/hypopnea is linked to chromosome 19, a region that contains the *ApoE* gene<sup>48</sup>, while different studies suggest that individuals carrying the *ApoE*- $\epsilon$ 4 locus have an increased risk of developing OSA/H<sup>49</sup>. At the same time, OSA *ApoE*- $\epsilon$ 4 carriers appear to have demonstrated an increased risk of impaired spatial working memory<sup>50</sup>. *ApoE*- $\epsilon$ 4 has also been associated with SDB<sup>51-53</sup>. On the other hand, studies with OSA patients showed that the frequency of the *ApoE*- $\epsilon$ 4 allele is the same as in a random population<sup>54-56</sup>. According to a study which compared individuals with different degrees of SDB, there was no association between SDB and *ApoE*- $\epsilon$ 4<sup>57</sup>. More existing studies found no association between *ApoE*- $\epsilon$ 4 and OSAS<sup>55,58</sup>. A different study investigating the association among napping, *ApoE*- $\epsilon$ 4, and dementia revealed that napping for up to 60 minutes had an apparently protective effect against developing AD, especially for the *ApoE*- $\epsilon$ 4 carriers<sup>59</sup>. Interestingly, our data analyses in a large group of non-demented older adults indicated that *ApoE*- $\epsilon$ 4 carriers had less snoring and subjective sleep apnea, even after controlling for multiple covariates<sup>60,61</sup>. A complex etiology combining multiple environmental and genetic causes could affect the association between sleep and *ApoE*- $\epsilon$ 4, leaving space for more research on the field.

Different genes have been also related to sleep regulation. The most common genes linked to sleep regulations are *circadian locomotor output cycles kaput* (*CLOCK*) which is mostly associated with insomnia and circadian rhythms, *PERIOD* (*PER*), linked to sleep homeostasis and circadian alertness, *Brain and Muscle Arnt-Like 1* (*BMAL1*) linked to sleep deprivation, and *crystal* (*CRY*) linked to sleep homeostasis<sup>62</sup>. Associations have been found between most of these genes and their mutations with cognitive phenotypes such as memory formation, consolidation, and cognitive alertness<sup>62,63</sup>. A different review reports that the rhythmic expression of specific sleep genes (*BMAL1*, *CRY1*, and *PER1*) is lost in neurodegenerative diseases such as MCI and AD<sup>64</sup>. Combining sleep, genes, and cognition, a review reports that specific molecular clocks in different brain regions and their circadian phases and anatomical relationships to the central brain pacemakers indicate new ways for understanding the mechanisms of interaction between circadian clocks, sleep and cognition<sup>62</sup>. A recent study suggests that a Polygenic Risk Score (PRS), which is calculated based on variation in multiple genetic loci and their associated weights, can identify associations between sleep duration and specific diseases such as congestive heart failure and obesity<sup>65</sup>. Thus, the association

among sleep, genes, and cognition remains to be explored further, with studies combining all the three factors and including multiple genes.

Apart from genetic factors, most recent studies examining sleep regulation and the brain focus their research on brain morphology. Age-related medial prefrontal cortex gray matter atrophy has been associated with reduced non-REM slow-wave activity in older adults, mediating the impairment of overnight sleep-dependent memory retention<sup>66</sup>. According to a different study, short sleep duration was associated with greater age-related brain atrophy and cognitive decline in older adults<sup>67</sup>. Based on a review, sleep deprivation and hippocampal vulnerability could cause changes in both neuronal plasticity, neurogenesis, and cognitive function<sup>68</sup>. Sleep deprivation has also been found to impair memory and frontal lobe function in insomnia patients<sup>69</sup>. Cortical thinning in the superior frontal lobe has been noted in female OSA patients<sup>70</sup>, while the hippocampus also shows lateralized and sex-specific, OSA-related regional volume differences<sup>71</sup>. Among cognitively normal older adults, both short and long sleep duration can increase the rate of subsequent frontotemporal gray matter atrophy<sup>72</sup>.

Preliminary results from our elderly group revealed significant associations between self-reported daytime sleepiness and total gray matter and cortical volume, while longer sleep duration was associated with low left entorhinal thickness<sup>73</sup>. From the fewer studies on sleep and functional connectivity, insomnia and more precisely difficulty in falling asleep has been linked to increased functional connectivity between sensory-motor brain regions<sup>74</sup>. Self-reported sleep duration has been also negatively correlated with prefrontal-amygdala connectivity in young adults<sup>75</sup>. Highlighting the role of sleep in brain function, a recent study reports that sleep-wake cycle regulates brain interstitial fluid tau in mice, and cerebro-spinal fluid tau in humans<sup>76</sup>. Hence, exploration of the links between sleep and brain morphometric factors can lead us to a better understanding of the neurobiology of sleep, and subsequently to a better understanding of how cognition works.

The importance of maintaining a normal sleep regulation is highlighted through the extensive literature. Sleep deprivation affects circadian clock gene expression in the cerebral cortex parallel electroencephalographic differences in an animal study<sup>77</sup>, promoting the need for similar research which combines information about sleep, genes, and brain morphometry, in humans. As sleep problems, especially short and long sleep duration, and obstructive sleep apnea as well, have been linked to even increased risk for mortality<sup>78,79</sup>, limiting the factors affecting negatively sleep regulation is of vital importance. Although in the current review we focused on sleep and cognition, as well as factors affecting this relationship, different factors could also play an important role to both sleep regulation and the brain function. Diet, light exposure, overall lifestyle, and multiple environmental elements could also contribute significantly to the association between sleep and brain<sup>80-82</sup>, indicating the need for further investigation, combining sleep, genes, brain biomarkers, and cognitive factors.

## ACKNOWLEDGEMENTS

The present work was supported by the “Bodossakis foundation”.

## REFERENCES

- Gooneratne NS, Vitiello MV. Sleep in older adults: normative changes, sleep disorders, and treatment options. *Clin Geriatr Med*. 2014;30(3):591-627.
- Basner M, Fomberstein KM, Razavi FM, Banks S, William JH, Rosa RR, et al. American time use survey: sleep time and its relationship to waking activities. *Sleep*. 2007;30(9):1085-95.
- Ohayon MM. Sleep and the elderly. *J Psychosom Res*. 2004;56(5):463-4.
- Ohayon MM, Carskadon MA, Guilleminault C, Vitiello MV. Meta-analysis of quantitative sleep parameters from childhood to old age in healthy individuals: developing normative sleep values across the human lifespan. *Sleep*. 2004;27(7):1255-73.
- Ramos AR, Dong C, Elkind MS, Boden-Albala B, Sacco RL, Rundek T, et al. Association between sleep duration and the mini-mental score: the Northern Manhattan study. *J Clin Sleep Med*. 2013;9(7):669-73.
- Miyata S, Noda A, Iwamoto K, Kawano N, Okuda M, Ozaki N. Poor sleep quality impairs cognitive performance in older adults. *J Sleep Res*. 2013;22(5):535-41.
- Merlino G, Piani A, Gigli GL, Cancelli I, Rinaldi A, Baroselli A, et al. Daytime sleepiness is associated with dementia and cognitive decline in older Italian adults: a population-based study. *Sleep Med*. 2010;11(4):372-7.
- Ohayon MM, Vecchierini MF. Daytime sleepiness and cognitive impairment in the elderly population. *Arch Intern Med*. 2002;162(2):201-8.
- Dealberto MJ, Pajot N, Courbon D, Alperovitch A. Breathing disorders during sleep and cognitive performance in an older community sample: the EVA Study. *J Am Geriatr Soc*. 1996;44(11):1287-94.
- Naumann A, Bellebaum C, Daum I. Cognitive deficits in narcolepsy. *J Sleep Res*. 2006;15(3):329-38.
- Jelicic M, Bosma H, Ponds RW, Van Boxtel MP, Houx PJ, Jolles J. Subjective sleep problems in later life as predictors of cognitive decline. Report from the Maastricht Ageing Study (MAAS). *Int J Geriatr Psychiatry*. 2002;17(1):73-7.
- Jaussent I, Bouyer J, Ancelin ML, Berr C, Foubert-Samier A, Ritchie K, et al. Excessive sleepiness is predictive of cognitive decline in the elderly. *Sleep*. 2012;35(9):1201-7.
- Cohen-Zion M, Stepnowsky C, Marler, Shochat T, Kripke DF, Ancoli-Israel S. Changes in cognitive function associated with sleep disordered breathing in older people. *J Am Geriatr Soc*. 2001;49(12):1622-7.
- Yaffe K, Laffan AM, Harrison SL, Redline S, Spira AP, Ensrud KE, et al. Sleep-disordered breathing, hypoxia, and risk of mild cognitive impairment and dementia in older women. *JAMA*. 2011;306(6):613-9.
- Blackwell T, Yaffe K, Laffan A, Ancoli-Israel S, Redline S, Ensrud KE, et al.; Osteoporotic Fractures in Men (MrOS) Study Group. Associations of objectively and subjectively measured sleep quality with subsequent cognitive decline in older community-dwelling men: The MrOS sleep study. *Sleep*. 2014;37(4):655-63.
- Keage HA, Banks S, Yang KL, Morgan K, Brayne C, Matthews FE. What sleep characteristics predict cognitive decline in the elderly? *Sleep Med*. 2012;13(7):886-92.
- Twoogor SS, Lee S, Schernhammer ES, Grodstein F. The association of self-reported sleep duration, difficulty sleeping, and snoring with cognitive function in older women. *Alzheimer Dis Assoc Disord*. 2006;20(1):41-8.
- Potvin O, Lorrain D, Forget H, Dubé M, Grenier S, Préville M, et al. Sleep quality and 1-year incident cognitive impairment in community-dwelling older adults. *Sleep*. 2012;35(4):491-9.
- Virta JJ, Heikkilä K, Perola M, Koskenvuo M, Riihä I, Rinne JO, et al. Midlife sleep characteristics associated with late life cognitive function. *Sleep*. 2013;36(10):1533-41, 1541A.
- Loerbroks A, Debling D, Amelang M, Stürmer T. Nocturnal sleep duration and cognitive impairment in a population-based study of older adults. *Int J Geriatr Psychiatry*. 2010;25(1):100-9.
- Devore EE, Grodstein F, Duffy JF, Stampfer MJ, Czeisler CA, Schernhammer ES. Sleep duration in midlife and later life in relation to cognition. *J Am Geriatr Soc*. 2014;62(6):1073-81.
- Xu L, Jiang CQ, Lam TH, Zhang WS, Cherny SS, Thomas GN, et al. Sleep duration and memory in the elderly Chinese: longitudinal analysis of the Guangzhou Biobank Cohort Study. *Sleep*. 2014;37(11):1737-44.
- Lim AS, Kowgier M, Yu L, Buchman AS, Bennett DA. Sleep Fragmentation and the Risk of Incident Alzheimer's Disease and Cognitive Decline in Older Persons. *Sleep*. 2013;36(7):1027-32.
- Nebes RD, Buysse DJ, Halligan EM, Houck PR, Monk TH. Self-reported sleep quality predicts poor cognitive performance in healthy older adults. *J Gerontol B Psychol Sci Soc Sci*. 2009;64(2):180-7.
- Foley DJ, Monjan AA, Masaki KH, Enright PL, Quan SF, White LR. Associations of symptoms of sleep apnea with cardiovascular disease, cognitive impairment, and mortality among older Japanese-American men. *J Am Geriatr Soc*. 1999;47(5):524-8.
- Tsapanou A, Gu Y, O'Shea D, Eich T, Tang MX, Schupf N, et al. Daytime somnolence as an early sign of cognitive decline in a community-based study of older people. *Int J Geriatr Psychiatry*. 2016;31(3):247-55.
- Tsapanou A, Gu Y, O'Shea DM, Yannakoulia M, Kosmidis M, Dardiotis E, et al. Sleep quality and duration in relation to memory in the elderly: Initial results from the Hellenic Longitudinal Investigation of Aging and Diet. *Neurobiol Learn Mem*. 2017;141:217-25.
- Foley D, Monjan A, Masaki K, Ross W, Havlik R, White L, et al. Daytime sleepiness is associated with 3-year incident dementia and cognitive decline in older Japanese-American men. *J Am Geriatr Soc*. 2001;49(12):1628-32.
- Schlosser Covell GE, Dhawan PS, Lee Iannotti JK, Hoffman-Snyder CR, Wellik KE, Caselli RJ, et al. Disrupted daytime activity and altered sleep-wake patterns may predict transition to mild cognitive impairment or dementia: a critically appraised topic. *Neurologist*. 2012;18(6):426-9.
- Elwood PC, Bayer AJ, Fish M, Pickering J, Mitchell C, Gallacher JE. Sleep disturbance and daytime sleepiness predict vascular dementia. *J Epidemiol Community Health*. 2011;65(9):820-4.
- Benito-León J, Bermejo-Pareja F, Vega S, Louis ED. Total daily sleep duration and the risk of dementia: a prospective population-based study. *Eur J Neurol*. 2009;16(9):990-7.
- Shi L, Chen SJ, Ma MY, Bao YP, Han Y, Wang YM, et al. Sleep disturbances increase the risk of dementia: A systematic review and meta-analysis. *Sleep Med Rev*. 2018;40:4-16.
- Tsapanou A, Gu Y, Manly J, Schupf N, Tang MX, Zimmerman M, et al. Daytime Sleepiness and Sleep Inadequacy as Risk Factors for Dementia. *Dement Geriatr Cogn Dis Extra*. 2015;5(2):286-95.
- Gehrman P, Marler M, Martin JL, Shochat T, Corey-Bloom J, Ancoli-Israel S. The relationship between dementia severity and rest/activity circadian rhythms. *Neuropsychiatr Dis Treat*. 2005;1(2):155-63.
- Tranah GJ, Blackwell T, Stone KL, Ancoli-Israel S, Paudel ML, Ensrud KE, et al.; SOF Research Group. Circadian activity rhythms and risk of incident dementia and mild cognitive impairment in older women. *Ann Neurol*. 2011;70(5):722-32.
- Cavanna AE, Trimble MR. The precuneus: a review of its functional anatomy and behavioural correlates. *Brain*. 2006;129(Pt 3):564-83.
- Spira AP, Gamaldo AA, An Y, Wu MN, Simonsick EM, Bilgel M, et al. Self-reported sleep and  $\beta$ -amyloid deposition in community-dwelling older adults. *JAMA Neurol*. 2013;70(12):1537-43.
- Stocker RPJ, Khan H, Henry L, Germain A. Effects of Sleep Loss on Subjective Complaints and Objective Neurocognitive Performance as Measured by the Immediate Post-Concussion Assessment and Cognitive Testing. *Arch Clin Neuropsychol*. 2017;32(3):349-68.
- Tardy M, Gonthier R, Barthelemy JC, Roche F, Crawford-Achour E. Subjective sleep and cognitive complaints in 65 year old subjects: a significant association. The PROOF cohort. *J Nutr Health Aging*. 2015;19(4):424-30.
- Kang SH, Yoon IY, Lee SD, Kim T, Lee CS, Han JW, et al. Subjective memory complaints in an elderly population with poor sleep quality. *Aging Ment Health*. 2017;21(5):532-6.
- Tsapanou A, Vlachos GS, Cosentino S, Gu Y, Manly JJ, Brickman AM, et al. Sleep and subjective cognitive decline in cognitively healthy elderly: Results from two cohorts. *J Sleep Res*. 2019;28(5):e12759.
- Gaultier C. [Obstructive sleep apnea syndrome and genetics]. *Rev Neurol (Paris)*. 2003;159(11 Suppl):6S98-101. [Article in French]
- Buxbaum SG, Elston RC, Tishler PV, Redline S. Genetics of the apnea hypopnea index in Caucasians and African Americans: I. Segregation analysis. *Genet Epidemiol*. 2002;22(3):243-53.
- Holberg CJ, Natrajan S, Cline MG, Quan SF. Familial Aggregation and Segregation Analysis of Snoring and Symptoms of Obstructive Sleep Apnea. *Sleep Breath*. 2000;4(1):21-30.
- Barclay NL, Gehrman PR, Gregory AM, Eaves LJ, Silberg JL. The heritability of insomnia progression during childhood/adolescence: results from a longitudinal twin study. *Sleep*. 2015;38(1):109-18.
- Strittmatter M, Hamann G, Cramer H, Reuner C, Kuntzmann F, Strubel D. Neurochemical parameters in senile dementia of the Alzheimer type. A longitudinal study in four cases. *Eur Arch Psychiatry Clin Neurosci*. 1996;246(2):110-1.
- Raber J. Androgens, apoE, and Alzheimer's disease. *Sci Aging Knowledge Environ*. 2004;2004(11):re2.

48. Palmer LJ, Buxbaum SG, Larkin E, Patel SR, Elston RC, Tishler PV, et al. A whole-genome scan for obstructive sleep apnea and obesity. *Am J Hum Genet.* 2003;72(2):340-50.
49. Gottlieb DJ, DeStefano AL, Foley DJ, Mignot E, Redline S, Givelber RJ, et al. APOE epsilon4 is associated with obstructive sleep apnea/hypopnea: the Sleep Heart Health Study. *Neurology.* 2004;63(4):664-8.
50. Cosentino FI, Bosco P, Drago V, Prestianni G, Lanuzza B, Iero I, et al. The APOE epsilon4 allele increases the risk of impaired spatial working memory in obstructive sleep apnea. *Sleep Med.* 2008;9(8):831-9.
51. Kadotani H, Kadotani T, Young T, Peppard PE, Finn L, Colrain IM, et al. Association between apolipoprotein E epsilon4 and sleep-disordered breathing in adults. *JAMA.* 2001;285(22):2888-90.
52. Osorio RS, Ayappa I, Mantua J, Gumb T, Varga A, Mooney AM, et al. Interaction between sleep-disordered breathing and apolipoprotein E genotype on cerebrospinal fluid biomarkers for Alzheimer's disease in cognitively normal elderly individuals. *Neurobiol Aging.* 2014;35(6):1318-24.
53. Lemoine P, Sassolas A, Lestra C, Laforest L, Chamba G. [Is there an interaction between sleep-disordered breathing, depression and apolipoprotein E phenotype?]. *L'Encephale.* 2004;30(4):360-2. [Article in French]
54. Larkin EK, Patel SR, Redline S, Mignot E, Elston RC, Hallmayer J. Apolipoprotein E and obstructive sleep apnea: evaluating whether a candidate gene explains a linkage peak. *Genet Epidemiol.* 2006;30(2):101-10.
55. Saarelainen S, Lehtimäki T, Kallonen E, Laasonen K, Poussa T, Nieminen MM. No relation between apolipoprotein E alleles and obstructive sleep apnea. *Clin Genet.* 1998;53(2):147-8.
56. Thakre TP, Mamtani MR, Kulkarni H. Lack of association of the APOE epsilon 4 allele with the risk of obstructive sleep apnea: meta-analysis and meta-regression. *Sleep.* 2009;32(11):1507-11.
57. Foley DJ, Masaki K, White L, Redline S. Relationship between apolipoprotein E epsilon4 and sleep-disordered breathing at different ages. *JAMA.* 2001;286(12):1447-8.
58. Uyrum E, Balbay O, Annakkaya AN, Gulec Balbay E, Silan F, Arbak P. The relationship between obstructive sleep apnea syndrome and apolipoprotein E genetic variants. *Respiration.* 2015;89(3):195-200.
59. Asada T, Motonaga T, Yamagata Z, Uno M, Takahashi K. Associations between retrospectively recalled napping behavior and later development of Alzheimer's disease: association with APOE genotypes. *Sleep.* 2000;23(5):629-34.
60. Tsapanou A, Scarmeas N, Gu Y, Manly J, Schupf N, Stern Y, et al. Examining the association between Apolipoprotein E (APOE) and self-reported sleep disturbances in non-demented older adults. *Neurosci Lett.* 2015;606:72-6.
61. Tsapanou A, Scarmeas N, Gu Y, Manly J, Schupf N, Stern Y, et al. Data from a cross-sectional study on Apolipoprotein E (APOE-ε4) and snoring/sleep apnea in non-demented older adults. *Data Brief.* 2015;5:351-3.
62. Kyriacou CP, Hastings MH. Circadian clocks: genes, sleep, and cognition. *Trends Cogn Sci.* 2010;14(6):259-67.
63. Sakai T, Tamura T, Kitamoto T, Kidokoro Y. A clock gene, period, plays a key role in long-term memory formation in *Drosophila*. *Proc Natl Acad Sci U S A.* 2004;101(45):16058-63.
64. Guarnieri B, Sorbi S. Sleep and Cognitive Decline: A Strong Bidirectional Relationship. It Is Time for Specific Recommendations on Routine Assessment and the Management of Sleep Disorders in Patients with Mild Cognitive Impairment and Dementia. *Eur Neurol.* 2015;74(1-2):43-8.
65. Dashti HS, Redline S, Saxena R. Polygenic risk score identifies associations between sleep duration and diseases determined from an electronic medical record biobank. *Sleep.* 2019;42(3). pii: zsy247.
66. Mander BA, Rao V, Lu B, Saletin JM, Lindquist JR, Ancoli-Israel S, et al. Prefrontal atrophy, disrupted NREM slow waves and impaired hippocampal-dependent memory in aging. *Nature Neurosci.* 2013;16(3):357-64.
67. Lo JC, Loh KK, Zheng H, Sim SK, Chee MW. Sleep duration and age-related changes in brain structure and cognitive performance. *Sleep.* 2014;37(7):1171-8.
68. Kreutzmann JC, Havekes R, Abel T, Meerlo P. Sleep deprivation and hippocampal vulnerability: changes in neuronal plasticity, neurogenesis and cognitive function. *Neuroscience.* 2015;309:173-90.
69. Noh HJ, Joo EY, Kim ST, Yoon SM, Koo DL, Kim D, et al. The relationship between hippocampal volume and cognition in patients with chronic primary insomnia. *J Clin Neurol.* 2012;8(2):130-8.
70. Macey PM, Haris N, Kumar R, Thomas MA, Woo MA, Harper RM. Obstructive sleep apnea and cortical thickness in females and males. *PLoS One.* 2018;13(3):e0193854.
71. Macey PM, Prasad JP, Ogren JA, Moiyadi AS, Aysola RS, Kumar R, et al. Sex-specific hippocampus volume changes in obstructive sleep apnea. *Neuroimage Clin.* 2018;20:305-17.
72. Spira AP, Gonzalez CE, Venkatraman VK, Wu MN, Pacheco J, Simonsick EM, et al. Sleep Duration and Subsequent Cortical Thinning in Cognitively Normal Older Adults. *Sleep.* 2016;39(5):1121-8.
73. Tsapanou A, Gu Y, Manly J, Schupf N, Scarmeas N, Barral S, et al. Association Between Sleep Problems and Brain Volume in the Elderly. *Neurology.* 2016;86(16 Suppl):19.001.
74. Killgore WD, Schwab ZJ, Kipman M, Deldonna SR, Weber M. Insomnia-related complaints correlate with functional connectivity between sensory-motor regions. *Neuroreport.* 2013;24(5):233-40.
75. Killgore WD. Self-reported sleep correlates with prefrontal-amygdala functional connectivity and emotional functioning. *Sleep.* 2013;36(11):1597-608.
76. Holth JK, Fritsch SK, Wang C, Pedersen NP, Cirrito JR, Mahan TE, et al. The sleep-wake cycle regulates brain interstitial fluid tau in mice and CSF tau in humans. *Science.* 2019;363(6429):880-4.
77. Wisor JP, Pasumarthi RK, Gerashchenko D, Thompson CL, Pathak S, Sancar A, et al. Sleep deprivation effects on circadian clock gene expression in the cerebral cortex parallel electroencephalographic differences among mouse strains. *J Neurosci.* 2008;28(28):7193-201.
78. Cappuccio FP, D'Elia L, Strazzullo P, Miller MA. Sleep duration and all-cause mortality: a systematic review and meta-analysis of prospective studies. *Sleep.* 2010;33(5):585-92.
79. Tsapanou A, Gu Y, O'Shea D, Manly J, Schupf N, Scarmeas N, et al. Self-Reported Sleep Disordered Breathing as Risk Factor for Mortality in the Elderly. *J Stroke Cerebrovasc Dis.* 2016;25(6):1524-31.
80. Anastasiou CA, Yannakoulia M, Kosmidis MH, Dardiotis E, Hadjigeorgiou GM, Sakka P, et al. Mediterranean diet and cognitive health: Initial results from the Hellenic Longitudinal Investigation of Ageing and Diet. *PLoS One.* 2017;12(8):e0182048.
81. Ancoli-Israel S, Gehrman P, Martin JL, Shochat T, Marler M, Corey-Bloom J, et al. Increased light exposure consolidates sleep and strengthens circadian rhythms in severe Alzheimer's disease patients. *Behav Sleep Med.* 2003;1(1):22-36.
82. Muzet A. Environmental noise, sleep and health. *Sleep Med Rev.* 2007;11(2):135-42.